

```
In [1]: #Tuple
```

```
In [2]: t1 = {1,2,3}
t1
```

```
Out[2]: {1, 2, 3}
```

```
In [3]: t1.add(10)
t1
```

```
Out[3]: {1, 2, 3, 10}
```

```
In [4]:
```

```
-----
AttributeError                                Traceback (most recent call last)
Cell In[4], line 1
----> 1 t1.append(11)
      2 t1

AttributeError: 'set' object has no attribute 'append'
```

```
In [5]: # SET remaining
```

```
In [6]: s7={1,2,3,4,5,}
s8={4,5,6,7,8}
s9={8,9,10}
```

```
In [7]: s7.intersection(s8)
```

```
Out[7]: {4, 5}
```

```
In [8]: s7.intersection(s9)
```

```
Out[8]: set()
```

```
In [9]: s8.intersection(s9)
```

```
Out[9]: {8}
```

```
In [10]: s7.difference(s8)
```

```
Out[10]: {1, 2, 3}
```

```
In [11]: s8 - s9
```

```
Out[11]: {4, 5, 6, 7}
```

```
In [12]: s7.symmetric_difference(s8)
```

```
Out[12]: {1, 2, 3, 6, 7, 8}
```

```
In [20]: s7={1,2,3,4,5,}  
s8={4,5,6,7,8}  
s9={8,9,10}
```

```
In [17]: s7.symmetric_difference(s8)
```

```
Out[17]: {1, 2, 3, 6, 7, 8}
```

```
In [16]: s7
```

```
Out[16]: {1, 2, 3, 4, 5}
```

```
In [21]: s7.symmetric_difference_update(s8) # will update the difference value to the s7
```

```
In [22]: s7
```

```
Out[22]: {1, 2, 3, 6, 7, 8}
```

```
In [23]: a={1,2,3,4,5,6,7,8,9}  
b= {3,4,5,6,7,8}  
c={10,20,30,40}
```

```
In [24]: a.issubset(b)
```

```
Out[24]: False
```

```
In [25]: a.issuperset(b)
```

```
Out[25]: True
```

```
In [26]: c.isdisjoint(a)
```

```
Out[26]: True
```

```
In [27]: b.issubset(a)
```

```
Out[27]: True
```

```
In [28]: a1={1,2,3,4,5,6,7,8,9}  
b1= {30,40,50,60,70,80}  
c1={10,20,30,40}
```

```
In [29]: a1.issubset(b1)
```

```
Out[29]: False
```

```
In [30]: b1.isdisjoint(a1)
```

```
Out[30]: True
```

```
In [31]: c1.isdisjoint(a1)
```

```
Out[31]: True
```

```
In [ ]:
```

In []:

In []: