



The Role of Nonadiabatic Couplings in Spectroscopic Calculations of Aromatic Molecules

Soumya Mukherjee¹, Saikat Hazra¹, Satyam Ravi^{1,2},
Subhankar Sardar^{1,3} and Satrajit Adhikari^{1,*}

¹School of Chemical Sciences, Indian Association for the Cultivation of Science,
Jadavpur, Kolkata - 700 032, West Bengal, India;

²School of Advance Science and Languages, VIT Bhopal University, Bhopal – 466114, India

³Department of Chemistry, Bhatler College, Paschim Midnapore, West Bengal, India

*Email: pcsa@iacs.res.in

Poster Presenter's Email: pcsm5@iacs.res.in, soumyamukherjee093@gmail.com

Objective and key features of my work

- Born-Oppenheimer (BO) approximation^{1,2} : No coupling between fast-moving electrons and slow-moving nuclei (huge differences in masses).
- At degenerate points, nonadiabatic coupling terms (NACTs) (electron-nuclei coupling) become infinite (Hellmann-Feynman theorem)^{3,4}. At other points also, NACT may have non-negligible value due to substantial coupling between electronic and nuclear motion.
- A paradigm shift from adiabatic to diabatic framework is needed.
- In diabatic representation, electron-nuclear couplings appear in form of diabatic couplings which are smooth, continuous as well as single-valued.
- Diabatic wavefunctions are formed by orthogonal rotation of the adiabatic wavefunction. The elements of such rotation matrix are obtained by employing the Adiabatic-to-Diabatic Transformation (ADT) condition^{5,6}.
- Finally, diabatic potential energy is calculated using the similarity transformation of adiabatic potential energy matrix by the rotation (ADT) matrix.
- We have developed a generalized code, 'ADT' to formulate analytic expressions of ADT quantities and compute their numerical values.⁷

1. Born M. and Oppenheimer J. R., Ann. Phys. 1927, **84**, 457.
2. Born M. and Huang K., Dynamical Theory of Crystal Lattices, Oxford University Press: Oxford, U.K., (1954)
3. Hellmann, H. Einfuhrang in die Quantenchemie; Franz Duetliche: Leipzig, Germany, 1937.
4. Feynman, R. Forces in Molecules. Phys. Rev. 1939, **56**, 340–343.
5. Baer, M. Phys. Rep. 2002, **358**, 75–142.
6. Sarkar, B.; Adhikari, S. J. Chem. Phys. 2006, **124**, 074101.
7. Naskar K., Mukherjee S., Mukherjee B., Ravi S., Mukherjee S., Sardar S. and Adhikari S. J. Chem. Theory Comput. 2020, **16**, 1666.

Origination of NACT

❖ BO expansion :

$$|\Psi(s_e, s_n)\rangle = \sum_i \psi_i(s_n) |\xi_i(s_e | s_n)\rangle$$

where, $|\Psi(s_e, s_n)\rangle \longrightarrow$ molecular wavefunction

$|\xi_i(s_e | s_n)\rangle \longrightarrow$ electronic wavefunctions

$\psi_i(s_n) \longrightarrow$ nuclear wavefunctions

❖ Total time-independent Schrödinger equation :

$$\hat{H}(s_e, s_n) |\Psi(s_e, s_n)\rangle = E |\Psi(s_e, s_n)\rangle$$

where, $\hat{H}(s_e, s_n) = \hat{T}_n(s_n) + \hat{H}_e(s_e | s_n) \longrightarrow$ sum of nuclear kinetic energy operator and electronic Hamiltonian

$$\hat{T}_n = -\frac{1}{2} \sum_i \left(\frac{\nabla_{s_n, i}^2}{m_i} \right) \longrightarrow \text{nuclear kinetic energy operator}$$

❖ Electronic Schrödinger equation :

$$\hat{H}_e(s_e | s_n) |\xi_i(s_e | s_n)\rangle = u_i(s_n) |\xi_i(s_e | s_n)\rangle \quad [u_i \text{ s are adiabatic potential energy}]$$

- ❖ Substituting the wavefunction and projecting this equation with $|\xi_j(s_e | s_n)\rangle$,

$$\left[-\frac{1}{2}(\vec{\nabla}_n + \vec{\tau})^2 + \mathbf{U} - \mathbf{E} \right] \psi = 0$$

where $\mathbf{U} \longrightarrow$ adiabatic potential energy matrix (diagonal matrix)

$\vec{\tau} \longrightarrow$ nonadiabatic coupling matrix (NACM). (skew-symmetric for real wavefunctions)

- ❖ Here, **NACT** :

$$\vec{\tau}_{ij} = \langle \xi_i(s_e | s_n) | \vec{\nabla} \xi_j(s_e | s_n) \rangle$$

- ❖ In BO approximation this is considered as 0 (i.e. no electron-nuclear coupling) and the above Schrödinger equation becomes,

$$\left[-\frac{1}{2} \nabla_n^2 + \mathbf{U} - \mathbf{E} \right] \psi = 0$$

Diabatic Representation

❖ Adiabatic wavefunctions :

$$\Psi = \mathbf{A}\phi^{\text{d}}$$

where, $\mathbf{A} \longrightarrow$ rotation matrix (orthogonal)

$\phi^{\text{d}} \longrightarrow$ diabatic nuclear wavefunction matrix.

❖ **Adiabatic to Diabatic Transformation (ADT) Condition :**

$$\vec{\nabla}_n \mathbf{A} + \vec{\tau} \mathbf{A} = 0$$

❖ Modified Schrödinger equation :

$$-\frac{1}{2} \nabla_n^2 \phi^{\text{d}} + (\mathbf{W} - \mathbf{E}) \phi^{\text{d}} = 0$$

where,

$$\mathbf{W} = \mathbf{A}^\dagger \mathbf{U} \mathbf{A}$$

Here \mathbf{W} is diabatic potential energy matrix (off diagonal terms are diabatic couplings and **they are not singular**)

Example: Diabatization for six state sub-Hilbert space ^[1]

$$\vec{\tau} = \begin{pmatrix} 0 & \vec{\tau}_{12} & \vec{\tau}_{13} & \vec{\tau}_{14} & \vec{\tau}_{15} & \vec{\tau}_{16} \\ -\vec{\tau}_{12} & 0 & \vec{\tau}_{23} & \vec{\tau}_{24} & \vec{\tau}_{25} & \vec{\tau}_{26} \\ -\vec{\tau}_{13} & -\vec{\tau}_{23} & 0 & \vec{\tau}_{34} & \vec{\tau}_{35} & \vec{\tau}_{36} \\ -\vec{\tau}_{14} & -\vec{\tau}_{24} & -\vec{\tau}_{34} & 0 & \vec{\tau}_{45} & \vec{\tau}_{46} \\ -\vec{\tau}_{15} & -\vec{\tau}_{25} & -\vec{\tau}_{35} & -\vec{\tau}_{45} & 0 & \vec{\tau}_{56} \\ -\vec{\tau}_{16} & -\vec{\tau}_{26} & -\vec{\tau}_{36} & -\vec{\tau}_{46} & -\vec{\tau}_{56} & 0 \end{pmatrix} \quad U = \begin{pmatrix} u_1 & 0 & 0 & 0 & 0 & 0 \\ 0 & u_2 & 0 & 0 & 0 & 0 \\ 0 & 0 & u_3 & 0 & 0 & 0 \\ 0 & 0 & 0 & u_4 & 0 & 0 \\ 0 & 0 & 0 & 0 & u_5 & 0 \\ 0 & 0 & 0 & 0 & 0 & u_6 \end{pmatrix}$$

$$\mathbf{A} = \mathbf{A}_{12}(\Theta_{12}).\mathbf{A}_{13}(\Theta_{13}).\dots\dots\dots\mathbf{A}_{45}(\Theta_{45}).\mathbf{A}_{16}(\Theta_{16}).\mathbf{A}_{26}(\Theta_{26}).\mathbf{A}_{36}(\Theta_{36}).\mathbf{A}_{46}(\Theta_{46}).\mathbf{A}_{56}(\Theta_{56})$$

❖ One of the rotation matrix :

$$\mathbf{A}_{12} = \begin{pmatrix} \cos \Theta_{12} & \sin \Theta_{12} & 0 & 0 & 0 & 0 \\ -\sin \Theta_{12} & \cos \Theta_{12} & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$


❖ One of the ADT equations :

$$\begin{aligned} \vec{\nabla} \Theta_{12} = & -\vec{\tau}_{12} - \sin \Theta_{12} \left\{ \vec{\tau}_{13} \tan \Theta_{13} + \sec \Theta_{13} \right. \\ & \left. (\vec{\tau}_{14} \tan \Theta_{14} + \vec{\tau}_{15} \sec \Theta_{14} \tan \Theta_{15} + \vec{\tau}_{16} \sec \Theta_{14} \sec \Theta_{15} \tan \Theta_{16}) \right\} \\ & - \cos \Theta_{12} \left\{ \vec{\tau}_{23} \tan \Theta_{13} + \sec \Theta_{13} (\vec{\tau}_{24} \tan \Theta_{14} \right. \\ & \left. + \vec{\tau}_{25} \sec \Theta_{14} \tan \Theta_{15} + \vec{\tau}_{26} \sec \Theta_{14} \sec \Theta_{15} \tan \Theta_{16}) \right\} \end{aligned}$$





BBO Treatment for $C_6H_6^+$

Ab Initio Calculations :



1. Completely symmetric breathing mode (Q_2).
2. Ring deformation mode (Q_8).
3. Degenerate C–C asymmetric stretching pair (Q_{16x} and Q_{16y} modes).
4. Degenerate in plane bending (Q_{18x} and Q_{18y} modes).
5. One component of degenerate out of plane bending (Q_{19x}).
6. Lowest five doublet states are \tilde{X}^2E_{1g} , \tilde{B}^2E_{2g} and \tilde{C}^2A_{2u} .
7. At equilibrium, it has D_{6h} symmetry.
8. cc-pVDZ/MRCI level energy calculation.
9. CAS is 9 electrons in 5 orbitals.
10. Nonadiabatic Coupling Terms (NACTs) are calculated by CP-MCSCF method.
11. $Q_i = \rho \cos \phi$ $Q_j = \rho \sin \phi$.



$Q_{18x}-Q_{19y}$ Pair: Adiabatic Energies and NACTs





Q_{18x} - Q_{19y} Pair: ADT Quantities




$$W_{ij}(Q_k, Q_l) = \sum_{n=0}^N \sum_{m=0}^M c_{nm} Q_k^n Q_l^{2m} \quad n, m \equiv \{0, 1, 2, \dots\}$$

Non-Traditional JT Model



1. Mukherjee S., Ravi S., Naskar K., Sardar S. and Adhikari S. J. Chem. Phys., 2021, **154**, 094306.

Photoelectron Spectra




$E(\text{eV})$ (experiment of Baltzer)	$E(\text{eV})$ (MCTDH dynamics)	$E(\text{eV})$ (present work)
9.2431	9.2544	9.2424 *
9.3265	9.3361	9.3120
9.3628	9.3766	9.3620 *
9.4002	9.4092	9.4170
9.477	9.4852	9.4793 *
9.537	9.5443	9.5384 *

1. Baltzer P. et al. Chem. Phys. 1997, **224**, 95.
2. Köppel H., Bâldea I., and Szalay P. G., Adv. Quantum Chem. 2003, **44**, 199.
3. Mukherjee S., Ravi S., Naskar K., Sardar S. and Adhikari S. J. Chem. Phys., 2021, **154**, 094306.



BBO Treatment for 1,3,5- $C_6H_3F_3^{\pm}$

Ab Initio Calculations :

1. C-F symmetric stretching mode (Q_2).
2. Degenerate C-C asymmetric stretching pair (Q_{9x} and Q_{9y} modes).
3. Degenerate C-C symmetric stretching pair (Q_{10x} and Q_{10y} modes).
4. Degenerate scissoring motion of C-C-C moiety (Q_{12x} and Q_{12y} modes).
5. Degenerate in-plane bending of C-C-C moiety (Q_{13x} and Q_{13y} modes)
6. Degenerate C-C twisting (Q_{18x} and Q_{18y} modes).
7. Lowest six doublet states are \tilde{X}^2E'' , \tilde{A}^2A_2'' , \tilde{B}^2E' and \tilde{C}^2A_2' .
8. At equilibrium, it has D_{3h} symmetry.
9. cc-pVDZ/MRCI level energy calculation.
10. CAS is 11 electrons in 9 orbitals.
11. Nonadiabatic Coupling Terms (NACTs) are calculated by CP-MCSCF method.
12. $Q_i = \rho \cos \phi$ $Q_j = \rho \sin \phi$.





Q_{9y} - Q_{18y} Pair: ADT Quantities




$$W_{ij}(Q_k, Q_l) = \sum_{n=0}^N \sum_{m=0}^M c_{nm} Q_k^n Q_l^{2m} \quad n, m \equiv \{0, 1, 2, \dots\}$$

Non-Traditional JT Model



Photoelectron Spectra




\tilde{X}^2E'' state		
$E(\text{eV})$ (experiment of Gilbert <i>et al.</i>)	$E(\text{eV})$ (MCTDH)	$E(\text{eV})$ (present work)
9.6383	9.6402	9.6324 *
9.7086	9.7069	9.7238
9.7696	9.7708	9.7600
9.8281	9.8101	9.8233 *
9.8917	9.8768	9.8834 *
9.9382	9.9408	9.9090
9.9939	9.9800	9.9895 *
10.0642	10.0477	10.0545*

Q_{13x} , Q_{13y} , Q_{18x} , Q_{18y} , Q_2 , Q_{9x} and Q_{9y}
normal modes are used

1. Gilbert, R.; Sauvageau, P.; Sandorfy, C. Chem. Phys. Lett. 1972, **17**, 465-470.
2. Mondal, T.; Mahapatra, S.; Phys. Chem. Chem. Phys. 2009, **11**, 10867-10880.
3. Mukherjee S., Ravi S., Dutta J., Sardar S. and Adhikari S. Phys. Chem. Chem. Phys., 2022, **22**, 2185.

Including Q_{10x} , Q_{10y} ,
 Q_{12x} , Q_{12y} normal modes



Future Plans

- Investigation of non-adiabatic interactions of o-difluorobenzene and pyridine cations by scanning the pairs of normal modes.
- Calculation of theoretical photoabsorption spectra of the neutral counterparts by TDDVR dynamics.

Acknowledgements

Soumya Mukherjee (File No: SPM-07/080(0250)/2016-EMR-I) acknowledges CSIR India and IACS for research fellowship. Saikat Hazra and Satyam Ravi thank IACS for research fellowship. Subhankar Sardar acknowledges Principal, Bhatler College, Dantan for providing research facility in his institution. Satrajit Adhikari acknowledges DST, India, through project no. File No. CRG/2019/000793 for the research funding. Satrajit Adhikari also acknowledges IACS for CRAY supercomputing facility.

Thank You