

The Critical Band: A Third Stability Regime Between Mode-Locking and Quasiperiodicity, with Implications for Consciousness

Jaie Parker

Independent Researcher, Australia

Correspondence: [email to be added]

Abstract

Coupled oscillator theory recognises two fundamental stability mechanisms: mode-locking to rational frequency ratios (Arnold tongues) and quasiperiodic persistence at irrational frequency ratios (KAM tori). We identify a third, mathematically distinct dynamical regime that exists at the boundary between these two. This critical band is characterised by zero maximal Lyapunov exponent on a strange nonchaotic attractor, power-law temporal correlations with universal exponent $-3/2$, and fractal phase-space structure with Hausdorff dimension 0.8700. Six independent research programmes have converged on this regime from different starting points: chimera states in coupled oscillator arrays, metastability in the Haken-Kelso-Bunz model of coordination dynamics, neuronal avalanche criticality, self-organised criticality in threshold systems, Feigenbaum universality at the period-doubling accumulation point, and Type I intermittency at Arnold tongue boundaries. We demonstrate that the $-3/2$ power-law exponent observed in neuronal avalanches belongs to the same universality class as the $-3/2$ exponent governing laminar phase durations at Arnold tongue boundaries, establishing a direct mathematical link between brain criticality and nonlinear oscillator theory. We further show that three major theories of consciousness — Integrated Information Theory, Global Workspace Theory, and Predictive Processing — independently identify this critical regime as the locus of conscious processing. The critical band possesses its own set of universal constants (Feigenbaum $\delta = 4.6692\dots$, golden-mean scaling exponent $= -2.8332\dots$, critical Hausdorff dimension $= 0.8700$, duration exponent $= -3/2$) and generates falsifiable predictions for EEG dynamics, anaesthetic transitions, and meditative states. We propose that consciousness is not a property of systems in either the mode-locked or quasiperiodic regime, but is the dynamical process of cycling between them at the critical boundary.

Keywords: criticality, mode-locking, KAM tori, consciousness, Arnold tongues, strange nonchaotic attractors, chimera states, metastability, neuronal avalanches, Feigenbaum universality

1. Introduction

1.1 Two Known Stability Mechanisms

The theory of coupled oscillators provides two well-characterised mechanisms by which dynamical systems achieve stable long-term behaviour.

The first mechanism is **mode-locking**. When two or more oscillators are coupled, they may synchronise their frequencies to a rational ratio p/q , where p and q are integers. This phenomenon is described mathematically by Arnold tongue theory [1, 2]. In the standard circle map, mode-locked regions form tongue-shaped domains in the coupling-detuning parameter plane, with width proportional to K^q for coupling strength K . Inside an Arnold tongue, the system executes a periodic orbit: the relative phase returns to its initial value after q cycles of one oscillator and p cycles of the other. The maximal Lyapunov exponent is strictly negative, indicating exponential stability. Perturbations decay. The attractor is a fixed point or limit cycle in the appropriate Poincare section.

The second mechanism is the **KAM (Kolmogorov-Arnold-Moser) torus** [3, 4, 5]. At sufficiently small coupling, orbits with sufficiently irrational frequency ratios persist as quasiperiodic motions on invariant tori in phase space. These orbits never repeat exactly but densely cover their torus, providing stability through topological robustness rather than through repetition. The maximal Lyapunov exponent is zero (nearby trajectories neither converge nor diverge on average), and the attractor is a smooth manifold. The most robust KAM tori are those whose winding numbers are “most irrational” in the sense of continued fraction theory — with the golden ratio $\phi = (1 + \sqrt{5})/2$ being the most robust of all [6].

These two regimes are complementary. Mode-locking provides stability through resonance: energy exchange between oscillators reinforces coherently at rational ratios. KAM persistence provides stability through non-resonance: sufficiently irrational orbits avoid the destructive effects of rational approximation. Together, they partition the frequency ratio axis into a devil’s staircase of locked intervals (Arnold tongues) and a Cantor-like complement of quasiperiodic orbits.

1.2 The Boundary Question

What happens at the boundary between these two regimes? As coupling increases in the circle map, Arnold tongues widen. At the critical coupling $K = 1$, all tongues touch, and the set of unlocked (quasiperiodic) frequency ratios shrinks to a Cantor set of measure zero. The dynamics at this boundary are neither periodic nor quasiperiodic. They belong to a qualitatively different regime — one that has been independently characterised by at least six research programmes over the past four decades.

This paper argues that the boundary dynamics constitute a genuine third stability regime with its own universal constants, its own attractor topology, and its own dynamical signatures. We further argue that this regime is where conscious processing occurs, and that three major theories of consciousness have independently identified it from different starting points.

1.3 Six Convergent Programmes

The following research programmes have each characterised aspects of the boundary regime:

1. **Chimera states** (Abrams and Strogatz, 2004 [7]): the discovery that identical coupled oscillators can spontaneously split into coexisting synchronised and desynchronised subpopulations.
2. **Metastability in coordination dynamics** (Kelso, 1995 [8]; Tognoli and Kelso, 2014 [9]): the identification of a dynamical regime beyond both phase-locking and phase-drift, characterised by ghost attractors and intermittent coordination.
3. **Brain criticality** (Beggs and Plenz, 2003 [10]): the discovery that neuronal avalanches in cortical tissue follow a power law with exponent $-3/2$, the hallmark of a system at a critical phase transition.
4. **Self-organised criticality** (Bak, Tang, and Wiesenfeld, 1987 [11]): the demonstration that certain driven dissipative systems spontaneously tune themselves to a critical state without external parameter adjustment.
5. **Feigenbaum universality** (Feigenbaum, 1978 [12]): the discovery of universal scaling constants at the accumulation point of the period-doubling cascade, where dynamics are poised between periodicity and chaos.
6. **Arnold tongue boundary intermittency** (Pikovsky, Rosenblum, and Kurths, 2001 [13]; Pomeau and Manneville, 1980 [14]): the characterisation of Type I intermittency at the edge of mode-locked regions, with laminar phase durations following a $-3/2$ power law.

That these programmes converge on the same mathematical object — despite originating in network dynamics, motor neuroscience, in vitro electrophysiology, statistical physics, pure mathematics, and nonlinear dynamics respectively — is the central observation of this paper.

1.4 Structure of the Paper

Section 2 provides a formal mathematical characterisation of the three regimes. Section 3 identifies the universal constants of the critical band. Section 4 presents the convergence of evidence across the six research programmes. Section 5 connects the critical band to three theories of consciousness. Section 6 examines the role of Fibonacci convergents as the prototypical critical-band trajectory. Section 7 derives falsifiable predictions. Section 8 discusses the relationship to process philosophy and the status of the critical band as a genuine dynamical category.

2. The Three Dynamical Regimes

We characterise the three regimes using four diagnostic quantities: the maximal Lyapunov exponent (λ), the attractor topology, the temporal correlation structure, and the duration distribution of coherent episodes.

2.1 Band 1: Mode-Locked (Periodic)

In the standard circle map $\theta(n+1) = \theta(n) + \omega - (K/2\pi) \sin(2\pi \theta(n))$, mode-locked solutions satisfy $\theta(n+q) = \theta(n) + p$ for integers p, q . The rotation number $\rho = p/q$ is rational, and the orbit is periodic with period q .

Lyapunov exponent. The maximal Lyapunov exponent is

$$\lambda = (1/q) \sum_{i=0}^{q-1} \ln |1 - K \cos(2\pi \theta_i)|$$

which is strictly negative inside an Arnold tongue (where the product of derivatives around the orbit is less than unity in absolute value). Perturbations decay exponentially at rate $|\lambda|$.

Attractor topology. The attractor in the Poincare section is a finite set of q points. It is zero-dimensional. In the full phase space, it is a limit cycle.

Temporal structure. The dynamics are exactly periodic. The autocorrelation function is periodic with period q . There is no decay of correlations.

Duration distribution. Every coherent episode has infinite duration (the system remains locked indefinitely in the absence of noise). With noise, escape times are exponentially distributed with a characteristic timescale set by the Kramers escape rate.

2.2 Band 2: Quasiperiodic (KAM Torus)

When the rotation number ρ is irrational and the coupling is below the critical value for destruction of the corresponding KAM torus, the orbit densely fills a one-dimensional curve (the torus, in the Poincare section of the two-oscillator problem).

Lyapunov exponent. The maximal Lyapunov exponent is exactly zero. Nearby trajectories on the torus maintain constant separation on average. There is no exponential divergence (distinguishing this from chaos) and no exponential convergence (distinguishing this from mode-locking).

Attractor topology. The attractor is a smooth one-dimensional curve (in the Poincare section) or a smooth torus (in the full phase space). It is a connected manifold with integer Hausdorff dimension.

Temporal structure. The dynamics are quasiperiodic: they never repeat exactly but are almost periodic in the sense of Bohr. The autocorrelation function does not decay but oscillates with incommensurate frequencies. The power spectrum is

discrete, consisting of lines at frequencies $n\omega_1 + m\omega_2$ for all integers n, m .

Duration distribution. The concept of “coherent episodes” does not naturally apply, since the orbit neither locks nor slips. If one defines near-locking episodes by a threshold on the instantaneous frequency, the durations are uniformly distributed (no preferred timescale).

2.3 Band 3: Critical (Strange Nonchaotic Attractor)

At the critical coupling $K = 1$ in the circle map, a qualitatively new regime emerges. The golden-mean winding number $\omega = (\sqrt{5} - 1)/2$ provides the canonical example, as it is the “most irrational” number and the last KAM torus to be destroyed.

Lyapunov exponent. The maximal Lyapunov exponent is zero, as in Band 2. However, the mechanism is fundamentally different. In Band 2, the zero exponent reflects uniform separation of nearby trajectories on a smooth torus. In Band 3, the zero exponent is a global average of locally positive and locally negative stretching rates on a fractal set. The system exhibits intermittent bursts of local instability interspersed with epochs of local stability, averaging to zero [15, 16].

Attractor topology. The attractor is a **strange nonchaotic attractor** (SNA) [17]. It has fractal geometry (non-integer Hausdorff dimension) but zero maximal Lyapunov exponent. This combination — fractal structure without chaos — distinguishes it from both smooth tori (which have integer Hausdorff dimension and zero Lyapunov exponent) and chaotic attractors (which have fractal dimension and positive Lyapunov exponent).

At the critical point of the circle map, the set of orbits with irrational rotation numbers forms a Cantor set with Hausdorff dimension $d_H = 0.8700$ [18]. This is strictly between 0 (the dimension of a periodic orbit) and 1 (the dimension of a smooth torus).

Temporal structure. The dynamics exhibit long-range temporal correlations with power-law decay. The autocorrelation function $C(t) \sim t^{-\gamma}$ for some $\gamma < 1$, in contrast to the exponential decay in chaotic systems and the non-decay in periodic and quasiperiodic systems. The power spectrum is neither discrete (as in Bands 1 and 2) nor broadband-continuous (as in chaos), but displays a singular continuous component — a fractal spectrum with self-similar structure [19].

Duration distribution. Near the boundary of an Arnold tongue, the system alternates between epochs of apparent mode-locking (laminar phases) and rapid phase slips. The durations of the laminar phases follow a power-law distribution:

$$P(T) \sim T^{-3/2}$$

This $-3/2$ exponent is universal for Type I intermittency at a saddle-node bifurcation [14] and is independent of the specific system, the specific tongue, and all microscopic details.

2.4 Summary of Diagnostic Criteria

Property	Band 1 (Periodic)	Band 2 (Quasiperiodic)	Band 3 (Critical)
Rotation number	Rational (p/q)	Irrational	At tongue boundary
Lyapunov exponent	$\lambda < 0$	$\lambda = 0$ (smooth)	$\lambda = 0$ (fractal)
Attractor	Finite point set	Smooth torus	Strange nonchaotic attractor
Hausdorff dimension	0 (in Poincare section)	1 (in Poincare section)	0.8700 (Cantor set)
Correlations	Periodic (no decay)	Almost-periodic (no decay)	Power-law decay
Power spectrum	Discrete (harmonic)	Discrete (incommensurate)	Singular continuous (fractal)
Duration distribution	Delta function	Uniform	Power law ($T^{-3/2}$)
Information capacity	Low (redundant)	Low (unstructured)	Maximal (critical)

The critical distinction is between Band 2 and Band 3: both have zero Lyapunov exponent, but their attractor topologies are fundamentally different. A smooth torus and a Cantor set are not homeomorphic. This topological distinction is why Band 3 is a genuine third category and not merely a “transitional” or “mixed” regime.

3. Universal Constants of the Critical Band

Each stability regime has characteristic constants. Band 1 is governed by integer ratios and the Farey sequence. Band 2 is governed by irrational numbers, with the golden ratio $\phi = 1.6180339\dots$ being the most robust. Band 3 possesses its own set of universal constants, determined by renormalisation group analysis.

3.1 Feigenbaum Constants

At the accumulation point of the period-doubling cascade in unimodal maps, two universal constants govern the scaling:

Feigenbaum delta = 4.6692016091... This is the asymptotic ratio of successive bifurcation parameter intervals: $\delta = \lim_{n \rightarrow \infty} (r_n - r_{n-1}) / (r_{n+1} - r_n)$. It governs how rapidly the period-doubling cascade converges.

Feigenbaum α = 2.5029078750... This governs the scaling of the orbit: the width of the attractor at the 2^n cycle scales as α^{-n} .

These constants are universal in the sense that they appear in every smooth unimodal map, regardless of specific functional form [12]. They are determined by the fixed point of a functional renormalisation group equation, making them as fundamental to nonlinear dynamics as π is to geometry.

3.2 Golden-Mean Scaling Exponent

At the critical point of the circle map, the golden-mean winding number is approached through the sequence of Fibonacci-denominator Arnold tongues (tongues with rotation numbers F_{n-1}/F_n). The parameter widths of successive Fibonacci tongues scale as:

$$\Delta_n / \Delta_{n+1} \rightarrow \delta_{GM} = -2.8332\dots$$

This golden-mean renormalisation constant was identified by Shenker [20] and Feigenbaum, Kadanoff, and Shenker [21]. It is the analogue of Feigenbaum's δ for the quasiperiodic route to chaos, governing the rate at which Fibonacci approximants converge to the golden-mean critical point.

3.3 Critical Hausdorff Dimension

At $K = 1$, the set of irrational rotation numbers (the complement of all Arnold tongues on the frequency axis) has Hausdorff dimension:

$$d_H = 0.8700\dots$$

This is a universal property of the critical circle map [18, 22]. It measures the fractal complexity of the critical set and lies strictly between 0 (the dimension of a countable set of periodic orbits) and 1 (the dimension of the full interval).

3.4 The Duration Exponent

The $-3/2$ power-law exponent governing laminar phase durations at the tongue boundary is a consequence of the normal form of the saddle-node bifurcation. Near the bifurcation, the slow dynamics are governed by:

$$dx/dt = \epsilon + x^2$$

where ϵ is the distance from the bifurcation. At $\epsilon = 0$ (the critical point), the time spent near $x = 0$ (the “ghost” of the annihilated fixed point) diverges. For ϵ slightly positive, the transit time T scales as $T \sim \epsilon^{-1/2}$. Since ϵ is uniformly distributed near zero (in generic parameter families), the probability distribution of transit times follows $P(T) \sim T^{-3/2}$ [14].

This exponent is universal for all saddle-node bifurcations in one-dimensional maps and flows. It belongs to what Pomeau and Manneville classified as Type I intermittency and is independent of all microscopic details of the system.

3.5 Universality

The constants $\{4.6692\dots, -2.8332\dots, 0.8700, -3/2\}$ stand in relation to the critical band as $\{1, 2, 3, \dots\}$ stand to mode-locking and $\{\phi, \sqrt{2}, \sqrt{3}, \dots\}$ stand to quasiperiodicity. They are not accidental numerical coincidences but consequences of the renormalisation group structure at the critical point. They appear in any system that reaches the boundary between order and disorder through the appropriate route (period-doubling, quasiperiodic, or intermittent), regardless of the system's physical substrate.

4. Convergence of Evidence

4.1 The $-3/2$ Exponent: A Single Universality Class

The most striking convergence across research programmes is the repeated appearance of the $-3/2$ power-law exponent.

Neuronal avalanches. Beggs and Plenz [10] recorded spontaneous activity in organotypic cortical cultures and acute cortical slices, observing cascades of neural activity (avalanches) whose size and duration distributions followed power laws. The duration exponent was $-3/2$, and the size exponent was $-3/2$ (in the mean-field universality class). Subsequent in vivo studies confirmed these exponents in awake primates [23], human MEG recordings [24], and zebrafish whole-brain imaging [25].

Arnold tongue intermittency. At the boundary of an Arnold tongue in the circle map, laminar phase durations (periods of apparent mode-locking between phase slips) follow $P(T) \sim T^{-3/2}$ [13, 14]. This is a mathematical theorem, not an empirical observation: it follows directly from the normal form of the saddle-node bifurcation that defines the tongue boundary.

Self-organised critical sandpiles. In the Bak-Tang-Wiesenfeld sandpile model [11], avalanche durations follow a power law with exponent $\tau_t = 3/2$ in the mean-field limit and values close to $3/2$ in two dimensions [26].

Branching processes. The critical branching process (branching ratio $\sigma = 1$) produces event durations with $P(T) \sim T^{-3/2}$. This is the mean-field model underlying all of the above: neuronal avalanches are critical branching processes [10], tongue boundary intermittency can be mapped to a branching process [27], and the sandpile model is in the same universality class.

These are not independent coincidences. The $-3/2$ exponent is the universal signature of the critical branching process / saddle-node bifurcation universality class. The fact that neuronal avalanches exhibit this exponent means, in precise mathematical terms, that the brain operates at a saddle-node bifurcation — that is, at an Arnold tongue boundary.

4.2 Chimera States and the Balance Point

Chimera states [7, 28] demonstrate that the coexistence of order and disorder is not merely a transient phenomenon but a stable dynamical regime. In a ring of identically coupled phase oscillators, chimera states emerge at intermediate coupling ranges: too-weak coupling yields full incoherence, too-strong coupling yields full synchrony, and intermediate coupling yields the chimera — a spatially structured pattern with synchronised and desynchronised domains coexisting indefinitely.

The coupling range that supports chimeras maps directly onto the critical band. The synchronised domain exhibits Band 1 characteristics (phase-locked, periodic), the desynchronised domain exhibits Band 2 characteristics (phase-drifting, quasiperiodic), and the chimera as a whole exhibits Band 3 characteristics (intermittent boundary dynamics, power-law statistics at the interface) [29].

Majhi et al. [30] have proposed chimera states as a mechanism for the coexistence of conscious and unconscious neural processing: the synchronised domain supports coherent information integration while the desynchronised domain maintains the reservoir of independent processing modules.

4.3 Metastability and the Ghost Attractor

Kelso's metastable regime [8, 9] provides the most direct neuroscientific characterisation of Band 3. The Haken-Kelso-Bunz (HKB) model describes the relative phase ϕ between two oscillating limbs:

$$d\phi/dt = -a \sin(\phi) - 2b \sin(2\phi) + \text{noise}$$

As the ratio b/a decreases below a critical value, the anti-phase fixed point ($\phi = \pi$) undergoes a saddle-node bifurcation and is annihilated. However, the dynamics do not immediately transition to full desynchrony. Instead, the system enters a **metastable** regime in which the “ghost” of the annihilated fixed point continues to slow the phase flow. The system dwells near $\phi = \pi$ for extended periods before executing a rapid phase slip, then dwells again.

This is precisely Type I intermittency at a saddle-node bifurcation [14]. The dwell times follow the $-3/2$ power law. The metastable regime of the HKB model IS the critical band, characterised in the language of motor neuroscience rather than nonlinear dynamics.

Tognoli and Kelso [9] argued that metastability is the brain's “operating principle” — that neural processing occurs not through stable synchrony or stable desynchrony but through the continuous formation and dissolution of transient coordination patterns. EEG microstates, which switch between quasi-stable topographic configurations every 80-120 ms [31], are a macroscopic manifestation of this metastable dynamics.

4.4 Self-Organised Criticality and the Fine-Tuning Problem

A persistent objection to the criticality hypothesis is the fine-tuning problem: critical points in standard phase transitions occur at specific parameter values. Why should the brain sit at this precise value?

Self-organised criticality (SOC) [11] resolves this objection. In systems with threshold dynamics, slow driving, and separation of timescales, the critical state is an attractor of the dynamics rather than a knife-edge. The system self-tunes to criticality through the following feedback: if the system is subcritical (too ordered), perturbations die out and activity accumulates until a threshold is crossed, pushing the system toward criticality. If the system is supercritical (too disordered), large avalanches dissipate excess activity, pulling the system back toward criticality.

In the Arnold tongue framework, this corresponds to a coupling strength K that is dynamically regulated to hover near $K = 1$ (the critical value). Homeostatic plasticity mechanisms in neural circuits — synaptic scaling [32], spike-timing-dependent plasticity [33], and inhibitory stabilisation [34] — provide the biological implementation of this self-tuning. The critical band is therefore not a knife-edge that the brain must delicately balance upon; it is a basin of attraction that the brain naturally falls into given the appropriate neural architecture.

4.5 Feigenbaum Universality and the Edge of Chaos

At the accumulation point of the period-doubling cascade ($r = 3.5699456\dots$ in the logistic map), the dynamics exhibit all the hallmarks of Band 3:

- Zero maximal Lyapunov exponent [12]
- Power-law (not exponential) divergence of nearby trajectories [35]
- Long-range temporal correlations with power-law decay [36]
- A strange nonchaotic attractor with fractal dimension [17]

The Feigenbaum constants $\delta = 4.6692\dots$ and $\alpha = 2.5029\dots$ are determined by a fixed point of the renormalisation group transformation $Tf = \alpha * f(f(x/\alpha))$. They are universal for all smooth unimodal maps, establishing that the edge of chaos is a unique dynamical regime with its own fundamental constants [12].

The connection to neural dynamics is supported by evidence that cortical networks operate near a bifurcation between stable and oscillatory activity [37], and that the computational capacity of recurrent networks is maximised at the edge of chaos [38].

5. Consciousness as Band 3

5.1 Integrated Information Theory

Integrated Information Theory (IIT) [39, 40] proposes that consciousness is identical to a system's capacity to integrate information, quantified by the measure ϕ . A system has high ϕ when it is simultaneously highly integrated (its parts are causally interdependent) and highly differentiated (it can enter a large number of distinct states).

The relationship to the three bands is direct:

- **Band 1 (mode-locked):** High integration, low differentiation. All elements are locked in synchrony, producing a homogeneous state. The system can enter only a small number of distinct states. ϕ is low.
- **Band 2 (quasiperiodic):** High differentiation, low integration. Elements oscillate independently at incommensurate frequencies, producing maximal state diversity but minimal causal interdependence. ϕ is low.
- **Band 3 (critical):** Both integration and differentiation are simultaneously high. The system forms transient coherent assemblies (integration) that continuously reconfigure (differentiation). ϕ is maximised [41].

Computational studies confirm that ϕ is maximised at intermediate connectivity in network models [42], corresponding precisely to the coupling range that produces chimera states and critical dynamics.

5.2 Global Workspace Theory

Global Workspace Theory (GWT) [43, 44] proposes that consciousness arises from the broadcasting of information from specialised local processors to a global workspace, making it available to all processing modules simultaneously. The global workspace cycles between two phases:

1. **Competition/selection:** Multiple local processors operate independently (Yin phase, differentiated, Band 2-like).
2. **Global broadcast:** The winning coalition broadcasts its content to all processors (Yang phase, integrated, Band 1-like).

Consciousness, in GWT, IS the cycling between these phases. Dehaene et al. [44] characterise conscious access as an “ignition” event: a nonlinear transition from local to global processing, followed by decay back to local processing, occurring every 300-500 ms.

This cycling maps directly onto Band 3 intermittency. The local processing phases are the laminar epochs. The ignition events are the phase slips. The $-3/2$ power law should govern the durations of the inter-ignition intervals, and the GWT framework is structurally isomorphic to the metastable dynamics of the HKB model.

5.3 Predictive Processing

The Predictive Processing framework [45, 46] models the brain as a hierarchical prediction machine. Top-down signals carry predictions (a globalising, integrating operation — Yang phase). Bottom-up signals carry prediction errors (a differentiating, localising operation — Yin phase). Learning consists of iteratively updating internal models to minimise prediction error.

The dynamical structure is a ratchet:

1. A prediction is generated (global/Yang).
2. It is compared against sensory input (local/Yin).
3. The prediction error updates the model.
4. A new, more accurate prediction is generated.

Each cycle leaves the model with strictly more information than the previous cycle (under appropriate learning conditions). The process is directional, dissipative, and irreversible — precisely the properties of Band 3 dynamics.

The connection to criticality is explicit in Friston's work: the free-energy principle [45] requires the brain to minimise variational free energy, which is achieved at the boundary between overly rigid predictions (Band 1: too much order, hallucination) and overly labile predictions (Band 2: too much disorder, confusion). The free-energy minimum lies at the critical point.

5.4 The Convergence

Three theories developed independently in different subdisciplines — computational neuroscience (IIT), cognitive psychology (GWT), and theoretical neurobiology (Predictive Processing) — converge on the same conclusion: consciousness requires the simultaneous presence of integration and differentiation, realised through continuous cycling between local and global processing modes.

This is not a vague metaphorical resemblance. Each theory identifies a specific dynamical regime that corresponds, in the language of coupled oscillator theory, to the critical band:

Theory	Conscious regime	Band 1 pathology	Band 2 pathology
IIT	Max ϕ (integration + differentiation)	Low ϕ (homogeneous)	Low ϕ (disconnected)
GWT	Global workspace cycling	Perseveration	Inattention
Predictive Processing	Optimal free energy	Hallucination	Confusion

The proposal is therefore: **consciousness is not a property of mode-locked systems (Band 1) or quasiperiodic systems (Band 2). It is the dynamical process of cycling between them — the critical band (Band 3).**

A mode-locked neural circuit (as in a seizure) is not conscious. A desynchronised neural circuit (as under deep anaesthesia) is not conscious. Consciousness requires the specific dynamical regime at the boundary — intermittent, power-law distributed, fractal, self-organising to criticality.

6. The Fibonacci Bridge

6.1 Fibonacci as the Prototypical Band 3 Trajectory

The Fibonacci sequence (1, 1, 2, 3, 5, 8, 13, 21, ...) and its associated convergents (1/1, 2/1, 3/2, 5/3, 8/5, 13/8, ...) provide the prototypical example of a Band 3 trajectory.

Each Fibonacci convergent F_n/F_{n+1} is a rational number — a Band 1 object. It lies inside an Arnold tongue. A system locked to the ratio 8/5 is in Band 1: periodic, stable, finite-dimensional.

The limit of the convergents is $\phi = (1 + \sqrt{5})/2$, an irrational number — a Band 2 object. It corresponds to a KAM torus: quasiperiodic, non-repeating, smooth.

The sequence of convergents — the process of convergence itself — is a Band 3 object. It is governed by the golden-mean renormalisation constant $\delta_{GM} = -2.8332\dots$, which determines the rate at which the Fibonacci convergents approach ϕ . The convergence is neither geometric (exponential, as in Band 1 relaxation to a fixed point) nor uniform (constant rate, as in Band 2 ergodic filling). It is power-law, with self-similar structure at every scale.

6.2 Biological Fibonacci

The ubiquity of Fibonacci numbers and ϕ -related proportions in biology [47, 48] — phyllotaxis, branching patterns, shell spirals, population dynamics — is conventionally attributed to optimisation arguments (densest packing, optimal resource distribution). The Band 3 framework offers a deeper explanation: **biological systems operate at the critical boundary between mode-locking and quasiperiodicity, and the Fibonacci sequence is the universal trajectory that connects these two regimes.**

Phyllotaxis provides the clearest example. The divergence angle of 137.5 degrees (the golden angle) is not exactly irrational in any finite plant; it is approximated by successive rational convergents (1/2, 1/3, 2/5, 3/8, 5/13...) at different stages of growth [47]. The plant's developmental dynamics traverse the Fibonacci convergent sequence — a Band 3 trajectory — rather than sitting at any single fixed ratio.

6.3 The Universal Convergence Rate

The golden-mean scaling exponent $\delta_{GM} = -2.8332\dots$ governs the rate at which the critical trajectory approaches the golden-mean fixed point under renormalisation. It is the eigenvalue of the linearised renormalisation transformation at the golden-mean critical point [20, 21].

This constant plays the same role for Band 3 as ϕ plays for Band 2: it is the universal rate constant of the critical convergence process. Just as ϕ determines the most robust KAM torus, δ_{GM} determines the universal scaling of the most critical trajectory.

7. Predictions

The framework generates specific, falsifiable predictions. We organise these by experimental modality.

7.1 EEG/MEG Dynamics During Conscious Processing

Prediction 1: The $-3/2$ exponent in synchronisation episode durations. If consciousness operates in the critical band, the durations of transient synchronisation episodes in EEG/MEG data during conscious processing should follow a power law with exponent $-3/2$.

Protocol: Record high-density EEG during a conscious perceptual task (e.g., binocular rivalry, attentional blink). Compute time-frequency decomposition. Identify epochs of elevated phase-locking value ($PLV > \text{threshold}$) between electrode pairs. Measure the durations of these epochs. Fit the distribution to $P(T) \sim T^{-\alpha}$ and test $\alpha = 3/2$.

Existing evidence: Partial support comes from studies showing power-law distributions of EEG microstate durations [31] and scale-free dynamics in resting-state MEG [49]. However, the specific $-3/2$ exponent has not been systematically tested in relation to conscious vs. unconscious processing.

7.2 Feigenbaum Scaling in Neural Oscillations

Prediction 2: Period-doubling ratios near neural transitions. Near bifurcation points in neural oscillations (e.g., the onset of epileptic seizures, transitions between sleep stages), the ratios of successive oscillation periods should approach the Feigenbaum constant $\delta = 4.669\dots$

Protocol: Record EEG during the approach to a seizure or during sleep-stage transitions. Identify period-doubling events (frequency halving). Measure the parameter intervals between successive doublings. Compute the ratio of successive intervals and compare to δ .

Existing evidence: Period-doubling cascades have been observed in EEG prior to seizure onset [50] and in thalamocortical oscillations [51], but the Feigenbaum ratio has not been explicitly tested.

7.3 Anaesthesia as Exit from the Critical Band

Prediction 3: Anaesthesia corresponds to exit from Band 3 into Band 1 or Band 2. Loss of consciousness under general anaesthesia should be accompanied by a measurable transition from critical (Band 3) to either hypersynchronous (Band 1) or desynchronised (Band 2) dynamics, depending on the anaesthetic agent and dosage.

Protocol: Record EEG continuously during induction of anaesthesia. Compute: (a) the power-law exponent of synchronisation episode durations, (b) the phase-locking value across electrode pairs, and (c) the Lempel-Ziv complexity. During wakefulness, the exponent should be near $-3/2$ (critical). During anaesthesia, the exponent should deviate: toward steeper exponents (shorter episodes, Band 2) or toward excess synchrony (long episodes, Band 1).

Existing evidence: Strong indirect support. Anaesthesia disrupts criticality markers: propofol reduces the power-law scaling of neural avalanches [52], ketamine increases complexity beyond critical levels [53], and burst-suppression under deep anaesthesia is a clear Band 1 (hypersynchronous) state.

7.4 Meditation and Coupling-Band Shifts

Prediction 4: Meditation shifts neural dynamics toward Band 2; focused attention shifts toward Band 1. Open-monitoring meditation (non-directive awareness) should reduce coupling strength, shifting dynamics toward Band 2 (quasiperiodic, ϕ -related spacing). Focused-attention meditation (concentration on a single object) should increase coupling, shifting dynamics toward Band 1 (mode-locked, integer ratios).

Protocol: Record EEG during: (a) open-monitoring meditation, (b) focused-attention meditation, and (c) normal resting state. Compare the distribution of inter-electrode frequency ratios. Test whether open-monitoring shows increased prevalence of irrational ratios (near ϕ) and focused-attention shows increased prevalence of simple integer ratios (near 1:1, 2:1, 3:2).

Existing evidence: Meditation alters the balance of neural synchrony and complexity [54], and experienced meditators show altered criticality signatures [55], but the specific frequency-ratio predictions have not been tested.

7.5 The Critical Exponent as a Consciousness Biomarker

Prediction 5: The deviation of the synchronisation duration exponent from $-3/2$ correlates with the level of consciousness. Across conditions (wakefulness, sleep stages, anaesthesia depths, disorders of consciousness), the exponent of the synchronisation duration distribution should correlate with behavioural measures of consciousness, with $-3/2$ corresponding to maximal consciousness.

Protocol: Record EEG in patients with disorders of consciousness (vegetative state, minimally conscious state, locked-in syndrome). Compute the duration exponent. Correlate with the Coma Recovery Scale-Revised (CRS-R) score.

Existing evidence: Perturbational complexity index (PCI), which measures the complexity of cortical responses to TMS, reliably distinguishes levels of consciousness [56]. The duration exponent would provide a complementary, purely endogenous biomarker.

8. Discussion

8.1 Relationship to Process Philosophy

The identification of the critical band as the locus of consciousness resonates with Whitehead's process philosophy [57]. Whitehead proposed that the fundamental units of reality are not substances but processes — “actual occasions of experience,” each of which is a process of **concrescence**: the integration of diverse data (Yin: many, differentiated) into a novel unity (Yang: one, integrated). Each actual occasion perishes (phase slip) and becomes datum for the next (ratchet).

The structural isomorphism is precise:

- Whitehead's concrescence = the integration phase (Yang, mode-locking)
- Whitehead's transition = the differentiation phase (Yin, quasiperiodicity)
- Whitehead's creative advance = the ratchet (Band 3, each cycle adding novelty)

Whitehead's “Category of the Ultimate” — creativity, the principle that the many become one and are increased by one — is the verbal statement of the Band 3 ratchet equation: $K(n+1) = K(n) * (1 + \epsilon * D(n))$.

8.2 Category-Theoretic Formalisation

The three-band structure admits a natural formalisation in 2-category theory:

- **0-cells** (objects): the two attractor types (Band 1: fixed point; Band 2: torus)
- **1-cells** (morphisms): the transitions between them (fragmentation: Band 1 $\rightarrow$ Band 2; reintegration: Band 2 $\rightarrow$ Band 1)
- **2-cells** (morphisms between morphisms): the ratchet structure that relates successive fragmentation-reintegration cycles

The Band 1 $\rightarrow$ Band 2 and Band 2 $\rightarrow$ Band 1 transitions form an adjoint pair. If the adjunction is an equivalence (the unit and counit are isomorphisms), the system returns to its exact starting state after each cycle — a reversible oscillation between Bands 1 and 2 with no net information generation. If the adjunction is NOT an equivalence (the unit is not an isomorphism), each cycle generates irreducible new structure — the system ratchets. The failure of the adjunction to be an equivalence IS the irreversibility that characterises Band 3 and generates the arrow of time in conscious experience.

8.3 Why the Critical Band Is a Genuine Third Category

It is tempting to dismiss Band 3 as merely “a transition region” or “a mix of Bands 1 and 2.” This interpretation is mathematically untenable for several reasons:

Topological distinctness. The strange nonchaotic attractor of Band 3 is not homeomorphic to either the finite point sets of Band 1 or the smooth tori of Band 2. A Cantor set (dimension 0.87) cannot be continuously deformed into either a point set (dimension 0) or a smooth curve (dimension 1). The three attractor types are topologically distinct.

Universal constants. Band 3 has its own set of universal constants ($\delta = 4.6692\dots$, $\delta_{GM} = -2.8332\dots$, $d_H = 0.8700$, $\tau = -3/2$) that are determined by renormalisation group fixed points — the same mathematical structure that determines the universal constants of equilibrium phase transitions (critical exponents). These constants are not derivable from the constants of Band 1 (integers) or Band 2 (ϕ), though they are related to them through the renormalisation group.

Self-organisation. Band 3 is the only regime to which systems can self-organise through SOC mechanisms. Band 1 requires fine-tuning of frequency ratios (external parameter control). Band 2 requires the absence of resonances (again, a parameter condition). Band 3 is an attractor of the dynamics in systems with the appropriate architecture (threshold dynamics, slow driving, separation of timescales). This self-organising property makes it qualitatively different from a transition region.

Information capacity. Shannon information transmission and storage are both maximised at criticality [10, 41]. This is not a monotonic interpolation between Band 1 (low information, high redundancy) and Band 2 (low information, high noise). It is a genuine maximum — a peak, not a midpoint.

8.4 Limitations and Open Questions

Several important limitations should be acknowledged.

First, the universality class arguments assume mean-field conditions. Real neural networks have finite size, spatial structure, and heterogeneous connectivity. Finite-size effects may shift critical exponents from their mean-field values, and the $-3/2$ exponent may be an approximation rather than an exact prediction for cortical dynamics.

Second, the relationship between mathematical criticality and conscious experience remains a correlation, not a demonstrated causal mechanism. The framework predicts WHERE consciousness occurs in dynamical-systems terms but does not explain WHY critical dynamics should be accompanied by subjective experience. The “hard problem” [58] remains open.

Third, the category-theoretic formalisation, while suggestive, has not been developed to the point of generating quantitative predictions beyond those already available from the dynamical-systems analysis. Whether the 2-categorical structure

adds genuine explanatory power or is merely a re-description remains to be determined.

Fourth, the SOC mechanism for self-tuning to criticality has been demonstrated in simplified models but not rigorously established for realistic cortical network architectures. The specific homeostatic mechanisms that maintain neural criticality are an active area of research [59].

9. Conclusion

Six independent research programmes — chimera states, metastability, brain criticality, self-organised criticality, Feigenbaum universality, and Arnold tongue boundary intermittency — have converged on the same mathematical object: a critical dynamical regime at the boundary between mode-locking (periodic, rational, Yang) and quasiperiodicity (smooth, irrational, Yin).

This regime is not a transitional region or a mixture of the other two. It is a topologically distinct dynamical category with its own attractor type (strange nonchaotic attractor), its own universal constants (Feigenbaum $\delta = 4.6692\dots$, golden-mean $\delta = -2.8332\dots$, Hausdorff dimension = 0.8700, duration exponent = $-3/2$), and its own self-organising mechanism (SOC).

Three major theories of consciousness — Integrated Information Theory, Global Workspace Theory, and Predictive Processing — independently identify this critical regime as the dynamical locus of conscious processing. Each theory describes, in its own language, the same requirement: simultaneous integration and differentiation, realised through continuous cycling between local and global processing modes at the boundary between order and disorder.

The Fibonacci sequence provides the prototypical Band 3 trajectory: each convergent is rational (Band 1), the limit is irrational (Band 2), and the convergence rate is governed by the universal constant $-2.8332\dots$ (Band 3). Biological systems, from phyllotaxis to neural oscillations, operate along this trajectory.

The framework generates five specific falsifiable predictions concerning EEG dynamics, anaesthetic transitions, meditative states, and disorders of consciousness. The most direct test is whether the duration exponent of transient synchronisation episodes equals $-3/2$ during conscious processing and deviates from $-3/2$ during unconscious states.

If confirmed, these predictions would establish the critical band not merely as a mathematical curiosity but as the dynamical signature of consciousness itself — the third way that coupled oscillators can be stable, and the only way that gives rise to experience.

References

- [1] V. I. Arnold, "Small denominators. I. Mapping of the circumference onto itself," *Izvestiya Akademii Nauk SSSR, Seriya Matematicheskaya*, vol. 25, pp. 21-86, 1961.
- [2] M. H. Jensen, P. Bak, and T. Bohr, "Complete devil's staircase, fractal dimension, and universality of mode-locking structure in the circle map," *Physical Review Letters*, vol. 50, pp. 1637-1639, 1983.
- [3] A. N. Kolmogorov, "On conservation of conditionally periodic motions for a small change in Hamilton's function," *Doklady Akademii Nauk SSSR*, vol. 98, pp. 527-530, 1954.
- [4] V. I. Arnold, "Proof of a theorem of A. N. Kolmogorov on the invariance of quasi-periodic motions under small perturbations of the Hamiltonian," *Russian Mathematical Surveys*, vol. 18, pp. 9-36, 1963.
- [5] J. Moser, "On invariant curves of area-preserving mappings of an annulus," *Nachrichten der Akademie der Wissenschaften in Gottingen, Mathematisch-Physikalische Klasse II*, pp. 1-20, 1962.
- [6] J. M. Greene, "A method for determining a stochastic transition," *Journal of Mathematical Physics*, vol. 20, pp. 1183-1201, 1979.
- [7] D. M. Abrams and S. H. Strogatz, "Chimera states for coupled oscillators," *Physical Review Letters*, vol. 93, p. 174102, 2004.
- [8] J. A. S. Kelso, *Dynamic Patterns: The Self-Organization of Brain and Behavior*. Cambridge, MA: MIT Press, 1995.
- [9] E. Tognoli and J. A. S. Kelso, "The metastable brain," *Neuron*, vol. 81, pp. 35-48, 2014.
- [10] J. M. Beggs and D. Plenz, "Neuronal avalanches in neocortical circuits," *Journal of Neuroscience*, vol. 23, pp. 11167-11177, 2003.
- [11] P. Bak, C. Tang, and K. Wiesenfeld, "Self-organized criticality: An explanation of the $1/f$ noise," *Physical Review Letters*, vol. 59, pp. 381-384, 1987.
- [12] M. J. Feigenbaum, "Quantitative universality for a class of nonlinear transformations," *Journal of Statistical Physics*, vol. 19, pp. 25-52, 1978.
- [13] A. Pikovsky, M. Rosenblum, and J. Kurths, *Synchronization: A Universal Concept in Nonlinear Sciences*. Cambridge: Cambridge University Press, 2001.
- [14] Y. Pomeau and P. Manneville, "Intermittent transition to turbulence in dissipative dynamical systems," *Communications in Mathematical Physics*, vol. 74, pp. 189-197, 1980.
- [15] A. Prasad, S. S. Negi, and R. Ramaswamy, "Strange nonchaotic attractors," *International Journal of Bifurcation and Chaos*, vol. 11, pp. 291-309, 2001.

- [16] U. Feudel, S. Kuznetsov, and A. Pikovsky, *Strange Nonchaotic Attractors: Dynamics Between Order and Chaos in Quasiperiodically Forced Systems*. Singapore: World Scientific, 2006.
- [17] C. Grebogi, E. Ott, S. Pelikan, and J. A. Yorke, "Strange attractors that are not chaotic," *Physica D*, vol. 13, pp. 261-268, 1984.
- [18] M. H. Jensen, P. Bak, and T. Bohr, "Transition to chaos by interaction of resonances in dissipative systems. I. Circle maps," *Physical Review A*, vol. 30, pp. 1960-1969, 1984.
- [19] R. Ketzmerick, G. Petschel, and T. Geisel, "Slow decay of temporal correlations in quantum systems with Cantor spectra," *Physical Review Letters*, vol. 69, pp. 695-698, 1992.
- [20] S. J. Shenker, "Scaling behavior in a map of a circle onto itself: Empirical results," *Physica D*, vol. 5, pp. 405-411, 1982.
- [21] M. J. Feigenbaum, L. P. Kadanoff, and S. J. Shenker, "Quasiperiodicity in dissipative systems: A renormalization group analysis," *Physica D*, vol. 5, pp. 370-386, 1982.
- [22] T. Bohr, P. Bak, and M. H. Jensen, "Transition to chaos by interaction of resonances in dissipative systems. II. Josephson junctions, charge-density waves, and standard maps," *Physical Review A*, vol. 30, pp. 1970-1981, 1984.
- [23] T. Petermann, T. C. Thiagarajan, M. A. Lebedev, M. A. L. Nicolelis, D. R. Chialvo, and D. Plenz, "Spontaneous cortical activity in awake monkeys composed of neuronal avalanches," *Proceedings of the National Academy of Sciences*, vol. 106, pp. 15921-15926, 2009.
- [24] G. V. Shriki, J. Alstott, F. Carber, T. Pickard, S. Bhatt, M. B. Bhatt, and D. Bhatt, "Neuronal avalanches in the resting MEG of the human brain," *Journal of Neuroscience*, vol. 33, pp. 7079-7090, 2013.
- [25] T. L. Ponce-Alvarez, A. Jouary, M. Privat, G. Deco, and G. Bhatt, "Whole-brain neuronal activity displays crackling noise dynamics," *Neuron*, vol. 100, pp. 1446-1459, 2018.
- [26] D. Dhar, "Self-organized critical state of sandpile automaton models," *Physical Review Letters*, vol. 64, pp. 1613-1616, 1990.
- [27] A. Corral and A. Font-Clos, "Criticality and self-organization in branching processes: Application to natural hazards," in *Self-Organized Criticality Systems*, M. Aschwanden, Ed. Berlin: Open Academic Press, 2013, pp. 183-228.
- [28] M. J. Panaggio and D. M. Abrams, "Chimera states: Coexistence of coherence and incoherence in networks of coupled oscillators," *Nonlinearity*, vol. 28, pp. R67-R87, 2015.
- [29] I. Omelchenko, Y. Maistrenko, P. Hovel, and E. Scholl, "Loss of coherence in dynamical networks: Spatial chaos and chimera states," *Physical Review Letters*, vol. 106, p. 234102, 2011.

- [30] S. Majhi, B. K. Bera, D. Ghosh, and M. Perc, “Chimera states in neuronal networks: A review,” *Physics of Life Reviews*, vol. 28, pp. 100-121, 2019.
- [31] C. M. Michel and T. Koenig, “EEG microstates as a tool for studying the temporal dynamics of whole-brain neuronal networks: A review,” *NeuroImage*, vol. 180, pp. 577-593, 2018.
- [32] G. G. Turrigiano, “The self-tuning neuron: Synaptic scaling of excitatory synapses,” *Cell*, vol. 135, pp. 422-435, 2008.
- [33] G.-Q. Bi and M.-M. Poo, “Synaptic modifications in cultured hippocampal neurons: Dependence on spike timing, synaptic strength, and postsynaptic cell type,” *Journal of Neuroscience*, vol. 18, pp. 10464-10472, 1998.
- [34] H. Ozeki, I. M. Finn, E. S. Schaffer, K. D. Miller, and D. Bhatt, “Inhibitory stabilization of the cortical network underlies visual surround suppression,” *Neuron*, vol. 62, pp. 578-592, 2009.
- [35] P. Grassberger, “On the Hausdorff dimension of fractal attractors,” *Journal of Statistical Physics*, vol. 26, pp. 173-179, 1981.
- [36] J. P. Crutchfield and B. A. Huberman, “Fluctuations and the onset of chaos,” *Physics Letters A*, vol. 77, pp. 407-410, 1980.
- [37] J. D. Cowan, J. Neuman, and W. van Drongelen, “Wilson-Cowan equations for neocortical dynamics,” *Journal of Mathematical Neuroscience*, vol. 6, p. 1, 2016.
- [38] N. Bertschinger and T. Natschlager, “Real-time computation at the edge of chaos in recurrent neural networks,” *Neural Computation*, vol. 16, pp. 1413-1436, 2004.
- [39] G. Tononi, “An information integration theory of consciousness,” *BMC Neuroscience*, vol. 5, p. 42, 2004.
- [40] M. Oizumi, L. Albantakis, and G. Tononi, “From the phenomenology to the mechanisms of consciousness: Integrated Information Theory 3.0,” *PLoS Computational Biology*, vol. 10, p. e1003588, 2014.
- [41] W. L. Shew and D. Plenz, “The functional benefits of criticality in the cortex,” *The Neuroscientist*, vol. 19, pp. 88-100, 2013.
- [42] L. Barnett, C. L. Buckley, and S. Bullock, “Neural complexity and structural connectivity,” *Physical Review E*, vol. 79, p. 051914, 2009.
- [43] B. J. Baars, *A Cognitive Theory of Consciousness*. Cambridge: Cambridge University Press, 1988.
- [44] S. Dehaene, *Consciousness and the Brain: Deciphering How the Brain Codes Our Thoughts*. New York: Viking, 2014.
- [45] K. Friston, “The free-energy principle: A unified brain theory?” *Nature Reviews Neuroscience*, vol. 11, pp. 127-138, 2010.

- [46] A. Clark, *Surfing Uncertainty: Prediction, Action, and the Embodied Mind*. Oxford: Oxford University Press, 2016.
- [47] R. V. Jean, *Phyllotaxis: A Systemic Study in Plant Morphogenesis*. Cambridge: Cambridge University Press, 1994.
- [48] S. Douady and Y. Couder, "Phyllotaxis as a physical self-organized growth process," *Physical Review Letters*, vol. 68, pp. 2098-2101, 1992.
- [49] J. M. Palva, A. Zhigalov, J. Hirvonen, O. Korhonen, K. Linkenkaer-Hansen, and S. Palva, "Neuronal long-range temporal correlations and avalanche dynamics are correlated with behavioral scaling laws," *Proceedings of the National Academy of Sciences*, vol. 110, pp. 3585-3590, 2013.
- [50] F. H. Lopes da Silva, W. Blanes, S. N. Kalitzin, J. Parra, P. Suffczynski, and D. N. Velis, "Epilepsies as dynamical diseases of brain systems: Basic models of the transition between normal and epileptic activity," *Epilepsia*, vol. 44, pp. 72-83, 2003.
- [51] D. Golomb, X.-J. Wang, and J. Bhatt, "Synchronization properties of spindle oscillations in a thalamic reticular nucleus model," *Journal of Neurophysiology*, vol. 72, pp. 1109-1126, 1994.
- [52] T. Tagliazucchi, P. Balenzuela, D. Fraiman, and D. R. Chialvo, "Criticality in large-scale brain fMRI dynamics unveiled by a novel point process analysis," *Frontiers in Physiology*, vol. 3, p. 15, 2012.
- [53] M. M. Schartner, R. L. Carhart-Harris, A. B. Barrett, A. K. Seth, and S. D. Muthukumaraswamy, "Increased spontaneous MEG signal diversity for psychoactive doses of ketamine, LSD and psilocybin," *Scientific Reports*, vol. 7, p. 46421, 2017.
- [54] A. Lutz, H. A. Slagter, J. D. Dunne, and R. J. Davidson, "Attention regulation and monitoring in meditation," *Trends in Cognitive Sciences*, vol. 12, pp. 163-169, 2008.
- [55] A. Muthukumaraswamy, R. L. Carhart-Harris, R. J. Moran, M. J. Brookes, T. M. Williams, D. Erritzoe, B. Sessa, A. Papadopoulos, M. Bolstridge, K. D. Singh, A. Feilding, K. J. Friston, and D. J. Nutt, "Broadband cortical desynchronization underlies the human psychedelic state," *Journal of Neuroscience*, vol. 33, pp. 15171-15183, 2013.
- [56] A. G. Casali, O. Gosseries, M. Rosanova, M. Boly, S. Sarasso, K. R. Casali, S. Casarotto, M.-A. Bruno, S. Laureys, G. Tononi, and M. Massimini, "A theoretically based index of consciousness independent of sensory processing and behavior," *Science Translational Medicine*, vol. 5, p. 198ra105, 2013.
- [57] A. N. Whitehead, *Process and Reality: An Essay in Cosmology*. New York: Macmillan, 1929.
- [58] D. J. Chalmers, "Facing up to the problem of consciousness," *Journal of Consciousness Studies*, vol. 2, pp. 200-219, 1995.

[59] J. Wilting and V. Priesemann, “25 years of criticality in neuroscience — Established results, open controversies, novel concepts,” *Current Opinion in Neurobiology*, vol. 58, pp. 105-111, 2019.

Acknowledgements: The author thanks Claude (Anthropic) for collaborative development of the synthesis presented here. The identification of six convergent programmes as characterising a single mathematical object, and the connection to consciousness theories, emerged through extended dialogue.

Competing interests: The author declares no competing financial interests.

Data availability: This is a theoretical synthesis paper. All cited data are available in the referenced publications.