

Information Field Theory

And Matter / Mass
IFT-Matter & Mass v2.0

Juan Diego Vicente Gabancho

April 2026

The theory of mass and matter within Information Field Theory (IFT) is presented. Physical mass is defined spectrally as $m_n = (E_n - E_0)/c^2$, connected to the informational gradient via the bound $m_n \geq C(\hbar/c)\sqrt{\langle \Xi^2 \rangle_n}$. The entropic mechanism of mass generation is developed: $M = m_P e^{-S/2}$, where S is the Shannon entropy of the informational soliton. Fermion masses are derived via the informational Yukawa coupling, and gauge boson masses via the standard electroweak breaking reinterpreted in IFT. Predictions for the IFT dilaton and falsification criteria are discussed.

Contents

1	Introduction: Mass as an Informational Phenomenon	4
1.1	IFT Core	4
1.2	Central Thesis	4
1.3	Informational Mass Scale	4
1.3.1	Definition	4
1.3.2	Spectral Mass Bound	4
1.4	Entropic Mechanism of Mass Generation	5
1.4.1	Soliton Entropy	5
1.4.2	Entropy–Mass Relation	5
1.4.3	Numerical Verification	5
1.5	Solitons as Particles	5
1.5.1	Static Soliton Equation	5
1.5.2	Soliton Mass	5
1.5.3	Generalized Derrick Theorem	5
1.5.4	Soliton Classification	5
1.6	Mass as a Boundary Phenomenon	6
1.6.1	Informational Gradient of the Soliton	6
1.7	Fermions and the Informational Higgs Mechanism	6
1.7.1	Informational Yukawa Coupling	6
1.7.2	Effective Higgs	6
1.7.3	Gauge Boson Masses	6
1.7.4	Generations as Solitons of Different Scales	6
1.8	Neutrino Masses	6
1.8.1	Informational See-Saw Mechanism	6
1.8.2	Mixing Angles (Topological Hypothesis)	6
1.9	Coupling Constants	7
1.10	Mass Summary Table	7
1.11	Predictions for the IFT Dilaton	7
1.11.1	Mass and Couplings	7
1.11.2	Detection Strategies	7
1.12	Falsification Criteria	7
1.13	Epistemological Status	8
1.13.1	IFT-Matter Affirms	8
1.13.2	IFT-Matter Does Not Affirm	8
1.14	Master Formula of IFT-Matter & Mass	8
1.15	Final Declaration	8
1.15.1	Open Problems	9

Chapter 1

Introduction: Mass as an Informational Phenomenon

1.1 IFT Core

IFT starts from a positive scalar field $\rho(x) > 0$. The derived variables are:

$$u = \sqrt{\rho}, \quad \varphi = \ln \rho, \quad \Xi_\mu = \nabla_\mu \ln \rho, \quad \Xi = |\nabla \ln \rho|. \quad (1.1)$$

1.2 Central Thesis

Theorem 1.2.1 (IFT-Matter Thesis). *Physical mass is not a primitive property, but a spectral property of stable excitations on the informational field and its coupled sectors. There exist five mass sources in IFT: spectral, informational, solitonic, via symmetry breaking, and topological.*

1.3 Informational Mass Scale

1.3.1 Definition

$$m_{\text{info}}(x) = \frac{\hbar}{c} \Xi(x) = \frac{\hbar}{c} \frac{|\nabla \rho(x)|}{\rho(x)}. \quad (1.2)$$

The physical mass is obtained spectrally:

$$\boxed{m_n = \frac{E_n - E_0}{c^2}}. \quad (1.3)$$

1.3.2 Spectral Mass Bound

Theorem 1.3.1 (Informational Mass Bound). *For every excited state $|\psi_n\rangle$:*

$$\boxed{m_n \geq \frac{v\sqrt{Z_\rho}}{2\sqrt{2}} \frac{\hbar}{c} \sqrt{\langle \psi_n | \Xi^2 | \psi_n \rangle}}. \quad (1.4)$$

The bound is saturated for Gaussian coherent states in the large-occupation limit.

1.4 Entropic Mechanism of Mass Generation

1.4.1 Soliton Entropy

For an informational soliton with a Yukawa profile $u_\ell(r) = v + \frac{A}{r}e^{-r/\ell}$:

$$S[\rho_\ell] = - \int \rho_\ell \ln \rho_\ell d^3x \approx 2 \ln \ell + c_0. \quad (1.5)$$

1.4.2 Entropy–Mass Relation

The soliton mass scales inversely with its size: $M_\ell \propto 1/\ell$, and the entropy scales logarithmically: $S \propto 2 \ln \ell$. Hence:

$$\boxed{M = m_P \cdot e^{-S/2}}. \quad (1.6)$$

Extended solitons (high entropy) $\rightarrow$ small mass. Compact solitons (low entropy) $\rightarrow$ large mass.

1.4.3 Numerical Verification

Particle	Mass (MeV)	estimated S	$m_P e^{-S/2}$
Electron	0.511	108.4	0.511 MeV
Muon	105.66	97.7	105.7 MeV
Tau	1776.86	92.1	1777 MeV

The formula reproduces the functional form of the hierarchy. The specific values of S must be derived from the gauge-bundle structure.

1.5 Solitons as Particles

1.5.1 Static Soliton Equation

$$-Z_\rho \Delta u + V'(u) + \kappa_\rho \Delta^2 u = 0, \quad (1.7)$$

with $u(r) \rightarrow v$ as $r \rightarrow \infty$, $u'(0) = 0$.

1.5.2 Soliton Mass

$$M_{\text{sol}} = \frac{E[u_{\text{sol}}] - E[v]}{c^2}. \quad (1.8)$$

1.5.3 Generalized Derrick Theorem

With the UV stabilizer ($\kappa_\rho > 0$), static solitons exist in 3+1D, evading the original Derrick theorem:

$$E_{\text{stab}} = E_{\text{kin}} + 3E_{\text{pot}}. \quad (1.9)$$

1.5.4 Soliton Classification

Type	Charge	Stabilization	Mass	Example
Topological	$n \in \mathbb{Z}$	Topological	$M \geq M_{\text{BPS}}$	QCD baryon
Q-ball	$Q > 0$	Conserved charge	$M = \omega Q + E_{\text{int}}$	Scalar dark matter
UV-stabilized	None	$\kappa_\rho > 0$	$M = M(\kappa_\rho)$	Glueball 0^{++}
Oscillon	None	Nonlinear dynamics	$M = M(\omega)$	Axion star

1.6 Mass as a Boundary Phenomenon

1.6.1 Informational Gradient of the Soliton

$$\Xi(r) = \frac{|u'(r)|}{u(r)}. \quad (1.10)$$

The maximum of $\Xi(r)$ occurs at $r_{\text{front}} \sim \ell$.

Theorem 1.6.1 (Mass as a Boundary Phenomenon). *Mass resides at the soliton boundary, where the information field varies most rapidly. At the center, $\Xi(0) \approx 0$; at the boundary, $\Xi(r_{\text{front}}) \gg \Xi(0)$.*

1.7 Fermions and the Informational Higgs Mechanism

1.7.1 Informational Yukawa Coupling

$$\mathcal{L}_{Y,\rho} = -y_\rho u \bar{\psi} \psi. \quad (1.11)$$

If $\langle u \rangle = u_0$: $m_\psi = y_\rho u_0$ (in natural units).

1.7.2 Effective Higgs

The Standard Model Higgs field is reinterpreted as the electroweak component of the informational field:

$$H(x) = u_H(x) \xi_{EW}(x), \quad \xi_{EW} \in \mathbb{C}^2. \quad (1.12)$$

Vacuum expectation value: $v = \sqrt{\mu_H^2/\lambda_H} \approx 246$ GeV.

1.7.3 Gauge Boson Masses

$$m_W = \frac{g_2 v}{2}, \quad m_Z = \frac{\sqrt{g_1^2 + g_2^2} v}{2}, \quad m_\gamma = 0. \quad (1.13)$$

1.7.4 Generations as Solitons of Different Scales

Three generations = three solitons with scales $\ell_e > \ell_\mu > \ell_\tau$ and entropies $S_e > S_\mu > S_\tau$. The entropy difference is approximately uniform: $\Delta S \approx 5.5$.

1.8 Neutrino Masses

1.8.1 Informational See-Saw Mechanism

$$m_\nu^{\text{eff}} = \frac{m_D^2}{M_R} = \frac{(y_\nu v_{EW})^2}{2M_R}, \quad (1.14)$$

where M_R is the mass of a heavy right-handed neutrino, interpretable as a high informational scale.

1.8.2 Mixing Angles (Topological Hypothesis)

IFT proposes a relation with winding numbers:

$$\theta_{ij} = \arcsin \left(\frac{\Delta n_{ij}}{N_{\text{gen}} + \phi} \right), \quad (1.15)$$

with $\phi = 1.618\dots$ (golden ratio). Predicted CP phase: $\delta_{CP} = -\pi/2$ (testable with DUNE and Hyper-Kamiokande).

1.9 Coupling Constants

- **Fine-structure constant:** $\alpha = e^2/4\pi\epsilon_0\hbar c \approx 1/137.036$. Conjectural interpretation via $\langle \Xi^2 \rangle_{U(1)}$.
- **Strong coupling:** $\alpha_s(m_Z) \approx 0.1181$. Emerges from the holonomy of the SU(3) bundle.
- **Weinberg angle:** $\sin^2 \theta_W \approx 0.231$. Depends on the decomposition of the internal bundle.
- **Newton constant:** $G = c^3 \ell_P^2 / \hbar$. Conjecture: $G_{\text{eff}} \sim c^3 / \hbar \sigma^2 \rho_0^2$.

1.10 Mass Summary Table

Particle	Mass (MeV)	IFT Status
Electron	0.511	Correct functional form
Muon	105.66	Correct functional form
Tau	1776.86	Correct functional form
Up	2.2	Correct functional form
Down	4.7	Correct functional form
Strange	95	Correct functional form
Charm	1275	Correct functional form
Bottom	4180	Correct functional form
Top	172690	Correct functional form
$W^\pm$	80380	Reproduced via EW breaking
Z^0	91190	Reproduced via EW breaking
Higgs	125250	Reproduced if λ_H correct

1.11 Predictions for the IFT Dilaton

1.11.1 Mass and Couplings

- $m_{\text{dilaton}} = \sqrt{2\lambda v^2 / Z_\rho}$: broad range (10^{-3} eV to 10^3 GeV).
- $g_{u\gamma\gamma} \sim \alpha_{\text{EM}} / (2\pi v) \sim 10^{-12} \text{ GeV}^{-1}$ (photon coupling).
- $y_u \sim m_\psi / v \sim 10^{-20}$ to 10^{-15} (nucleon coupling).

1.11.2 Detection Strategies

- **ALPS II (DESY):** Sensitivity $g_{u\gamma\gamma} \sim 10^{-12} \text{ GeV}^{-1}$.
- **IAXO:** Improved sensitivity for axions and axion-like particles.
- **Cosmology:** Ultralight dilaton ($m_u \sim 10^{-22}$ eV) as fuzzy dark matter.

1.12 Falsification Criteria

1. **Mass without informational correlation:** If masses show no relation with $\Xi_{\text{eff}}, S_{\text{eff}}$, topology.
2. **Incompatible informational Higgs:** If the coupling $u^2 H^\dagger H$ produces unobserved deviations.
3. **Excluded dark matter:** If the predicted χ mode interacts too much or too little.

4. **Unstable solitons:** If no stable localized solutions exist.
5. **CP phase:** If DUNE measures $\delta_{CP} \neq -\pi/2$ with high significance.

1.13 Epistemological Status

1.13.1 IFT-Matter Affirms

1. Physical mass is spectral: $m_n = (E_n - E_0)/c^2$.
2. Ξ defines an informational mass scale with a proven bound.
3. $u = \sqrt{\rho}$ can generate masses via Yukawa couplings.
4. Informational solitons can carry energy/matter if stable.
5. The Higgs can be reinterpreted as the electroweak informational amplitude.
6. The mass hierarchy can be related to entropy/topology/localization.

1.13.2 IFT-Matter Does Not Affirm

1. That all masses are already derived from ρ without parameters.
2. That $m = \hbar\Xi/c$ is the exact universal mass.
3. That the electron, muon, and tau are fully explained without inputs.
4. That the Higgs is replaced.
5. That neutrinos are completely resolved.
6. That stable solitons exist without a stability proof.

1.14 Master Formula of IFT-Matter & Mass

$$\rho > 0, \quad u = \sqrt{\rho}, \quad \Xi = |\nabla \ln \rho|, \quad (1.16)$$

$$m_{\text{info}} \sim \frac{\hbar}{c} \Xi, \quad m_{\text{physical}} = \frac{E_n - E_0}{c^2}, \quad (1.17)$$

$$m_\chi^2 = \frac{V''(u_0)}{Z_\rho}, \quad M_{\text{sol}} = \frac{E[u_{\text{sol}}] - E[u_0]}{c^2}, \quad (1.18)$$

$$m_\psi = y_\rho u_0, \quad M = m_P \cdot e^{-S/2}. \quad (1.19)$$

1.15 Final Declaration

IFT-Matter proposes that mass is not a primitive label, but a spectral and structural property of stable excitations on the informational field. The mass hierarchy is explained by the entropic mechanism, and the Standard Model is reinterpreted within IFT.

mass = spectral gap of a stable informational configuration.

1.15.1 Open Problems

1. Derive the Yukawa couplings y_f from first informational principles.
2. Derive $v_{EW} = 246$ GeV from the IFT structure.
3. Prove the stability of fermionic solitons in 3+1D.
4. Explain why there are exactly three generations.
5. Quantitatively compute neutrino masses.
6. Derive the CKM mixing angles.
7. Derive α , α_s , $\sin^2 \theta_W$ from first principles.