

# Chapter 2

Name: Liu Zhaohong

Student ID: 122431910061

## Exercise 2.1

Given the following 4 data points

$$\{(0, 1), (1, 2), (3, 6), (5, 7)\}$$

find a polynomial in Lagrange form to interpolate these data

we know that the formula for Lagrange interpolation is:

$$L_{n,k}(x) = \prod_{i=0, i \neq k}^n \frac{x - x_i}{x_k - x_i}$$

so the Lagrange polynomial for the 4 points should be:

$$\begin{aligned}\mathcal{L}_0 &= -\frac{-15 + 23x - 9x^2 + x^3}{15} \\ \mathcal{L}_1 &= \frac{15x - 8x^2 + x^3}{4} \\ \mathcal{L}_2 &= -\frac{5x - 6x^2 + x^3}{2} \\ \mathcal{L}_3 &= \frac{7(3x - 4x^2 + x^3)}{40}\end{aligned}$$

so the final polynomial should be sum of the 4 equations:

$$\mathcal{L} = 1 + \frac{29x}{120} + \frac{9x^2}{10} - \frac{17x^3}{120}$$

## Exercise 2.5

Consider a function  $f(x) = \sin(\pi x) + 3x$  at 6 distinct nodes in the interval  $[-1, 1]$  to determine the Newton's divided difference formula. The data are given in Table below:

$x$	-1.0	-0.6	-0.2	0.2	0.6	1.0
$f(x)$	-3.0	-2.7511	-1.1878	1.1878	2.7511	3.0

the whole coefficient matrix can be presented below:

$$\begin{bmatrix} p=0 & p=1 & p=2 & p=3 & p=4 & p=5 \\ -3 & & & & & \\ -2.7511 & 0.62225 & & & & \\ -1.1878 & 3.90825 & 4.1075 & & & \\ 1.1878 & 5.939 & 2.5384375 & -1.30755208 & & \\ 2.7511 & 3.90825 & -2.5384375 & -4.23072917 & -1.82698568 & \\ 3 & 0.62225 & -4.1075 & -1.30755208 & 1.82698568 & 1.82698568 \end{bmatrix}$$


according to the Newton's divided difference formula,

$$P_n(x) = F_{0,0} + \sum_{i=1}^n F_i \prod_{j=0}^{i-1} (x - x_j)$$

so the polynomial should be:

$$\begin{aligned}
 P(x) = & -3 \\
 & + 0.62225(x + 1.0) \\
 & + 4.1075(x + 1.0)(x + 0.6) \\
 & - 1.30755208(x + 1.0)(x + 0.6)(x + 0.2) \\
 & - 1.82698568(x + 1.0)(x + 0.6)(x + 0.2)(x - 0.2) \\
 & + 1.82698568(x + 1.0)(x + 0.6)(x + 0.2)(x - 0.2)(x - 0.6)
 \end{aligned}$$

compare the polynomial we get and the real function  $f(x) = \sin(\pi x) + 3x$



we can see that **inside** the interval  $[-1, 1]$ , the polynomial performs well, but the error will exceed greatly **beyond** the interval.

## Exercise 2.6

Construct an approximating polynomial for the following data using Hermite interpolation with Newton's forward difference

Table 1:

$x$	<b>0.1</b>	<b>0.2</b>	<b>0.3</b>
$f(x)$	-0.29004996	-0.56079734	-0.81401972
$f'(x)$	-2.8019975	-2.6159201	-2.9734038

Table 2:

$x$	<b>-1.0</b>	<b>-0.5</b>	<b>0.0</b>	<b>0.5</b>
$f(x)$	0.86199480	0.95802009	1.0986123	1.2943767
$f'(x)$	0.15536240	0.23269654	0.33333333	0.45186776

for data in Table 1:

$z_0 = 0.1$	$f[z_0] = -0.29004996$	$-0.29004996$					
$z_1 = 0.1$	$f[z_1] = -0.29004996$	$-0.29004996$	$-2.8019975$				
$z_2 = 0.2$	$f[z_2] = -0.56079734$	$-0.56079734$	$-2.7074738$	$0.945237$			
$z_3 = 0.2$	$f[z_3] = -0.56079734$	$-0.56079734$	$-2.6159201$	$0.915537$	$-0.297$		
$z_4 = 0.3$	$f[z_4] = -0.81401972$	$-0.81401972$	$-2.5322238$	$0.836963$	$-0.39287$	$-0.47935$	
$z_5 = 0.3$	$f[z_5] = -0.81401972$	$-0.81401972$	$-2.9734038$	$-4.4118$	$-52.48763$	$-260.4738$	$-1299.97225$

for data in Table 2:

0.8619948							
0.8619948	0.1553624						
0.95802009	0.19205058	0.07337636					
0.95802009	0.23269654	0.08129192	0.01583112				
1.0986123	0.28118442	0.09697576	0.01568384	$-0.00014728$			
1.0986123	0.33333333	0.10429782	0.01464412	$-0.00103972$	$-0.00089244$		
1.2943767	0.3915288	0.11639094	0.01209312	$-0.002551$	$-0.00100752$	$-0.00007672$	
1.2943767	0.45186776	0.12067792	0.00857396	$-0.00351916$	$-0.00096816$	$0.00002624$	$0.00006864$

so the polynomial 1 should be:

$$\begin{aligned}
 P(x) = & -0.29004996 \\
 & -2.8019975(x - 0.1) \\
 & + 0.945237(x - 0.1)^2 \\
 & - 0.297(x - 0.1)^2(x - 0.2) \\
 & - 0.47935(x - 0.1)^2(x - 0.2)^2 \\
 & - 1299.97225(x - 0.1)^2(x - 0.2)^2(x - 0.3)
 \end{aligned}$$

the polynomial 2 should be:


$$\begin{aligned}
 P(x) = & 0.8619948 \\
 & + 0.1553624(x + 1) \\
 & + 0.07337636(x + 1)^2 \\
 & + 0.01583112(x + 1)^2(x + 0.5) \\
 & - 0.00014728(x + 1)^2(x + 0.5)^2 \\
 & - 0.00089224(x + 1)^2(x + 0.5)^2x \\
 & - 0.00007672(x + 1)^2(x + 0.5)^2x^2 \\
 & + 0.00006864(x + 1)^2(x + 0.5)^2x^2(x - 0.5)
 \end{aligned}$$

## Exercise 2.7

Choose equally-spaced 15 points in  $[-5, 5]$  and get the data points

$$\{x_i, f(x_i)\}_{i=0}^{14}$$

by  $f(x) = 1/(1 + x^2)$ . Using piecewise linear interpolation to get the approximation of  $f(x)$  in  $[-5, 5]$  and, compare the result with that obtained by Lagrange interpolation.



It can be seen that Lagrange interpolation keeps a good tracing trend in the center area; Piecewise interpolation keeps a good trending in all area;