




Calculating theoretical probabilities from two-way tables (two events)

- 1 If this spinner is spun twice, what is the probability that the sum of both results would be 11? Tick **1** correct answer



- ☐ $\frac{1}{32}$
- ☐ $\frac{1}{64}$
- ☐ $\frac{1}{16}$
- ☐ $\frac{11}{64}$


- 2 If this spinner is spun twice, what is the probability of not losing either time? Tick **1** correct answer



- ☐ $\frac{1}{9}$
- ☐ $\frac{1}{3}$
- ☐ $\frac{2}{9}$
- ☐ $\frac{4}{9}$



- 3 If this spinner is spun twice, what is the probability that the sum of both results would be odd? Tick **1** correct answer




- ☐ $\frac{1}{2}$
- ☐ $\frac{1}{4}$
- ☐ $\frac{1}{3}$

- 4 If spun twice what is the probability this spinner would spell 'pea' then 'nut' ? Tick **0** correct answers



- ☐ $\frac{1}{5}$
- ☐ $\frac{1}{25}$
- ☐ $\frac{2}{5}$


- 5 If each of these spinners is spun once, what is the probability of getting H and 6? Tick **1** correct answer



- ☐ $\frac{1}{10}$
- ☐ $\frac{1}{5}$
- ☐ 0



- 6 If this spinner is spun twice, what is the probability of the sum of the results being less than 10? Tick **1** correct answer



- ☐ $\frac{19}{64}$
- ☐ $\frac{23}{64}$
- ☐ $\frac{17}{64}$

