02-10-'20 13:37 FR0M- T-977 Created with a trial version of Syncfusion Essential PDF



FAX TRANSMISSION

Date 2 10 20 20 Time:	From:
Receiver's Name: Infusion Inbound Phone Front Desk:845-333-1150	Department: RAVI RAM 43 WAMS Phone: (805) 342 777 4 Fax: 845 3423229
Stacy phone : 845-333-1905	1 an. 375 27 25 227
Inbound Fax : 845-333-9400	

Information Transmitted: Appointment Date Needed: Name of Patient: ______MIKAL, (If Applicable) Diagnosis written on Order: Authorization Number:

Thank you

The documents accompanying this transmission contain confidential information belonging to the sender, which is legally privileged. This information is intended only for he use of the individuals or entity named above. The authorized recipient of the information is prohibited from disclosing the information to any other party and is required to destroy the information after it's stated need has been fulfilled. If you are not the intended recipient, you are hereby notified that any disclosure, copying distribution or action taken in reliance on the contents of these documents is strictly prohibited. If you RECEIVED 02-07-20 12:17 FROM- TO-

ORANGE REGIONAL MEDICAL Physician Order Form		1	VAVI KAS				
DO NOT USE ABBREVIATIONS:			NAVIKAS, MKAZ				
U MS MSO4 MgSO4 QC			MRN 214800				
OUTPATIENT DARBEPOETIN (Aranesp®) Erythropoetic Stimulating Agent (ESA) Order Form							
Initial Hemoglobin (Hgb)/Hematocrit (Hct) Date/level: Wt:kg Current Hgb: Hct:							
DIAGNOSIS (Check one):							
□ Chemotherapy induced anemia Hemoglobin (Hgb) / Hematocrit (Hct) must be < 10 g/dl and < 30% Darbepoetin initial dose ≤ 2.25 mcg/kg/week On week 4 doses may be increased by 25% if Hgb/Hct rise is < 1/3. By week 8 discontinue treatment unless a rise of 1/3 is documented							
□ End Stage Renal Disease (ESRD) on dialysis Chronic Kidney Disease (CRD) not on dialysis For chronic kidney disease creatinine ≥ 3 and CrCl < 60 mL/min Monitor to ensure transferring saturation > 20% and/or serum ferritin > 100 ng/mL							
□ HIV/AIDS anemia induced by zidovudine or other nucleoside reverse transcriptase inhibitors							
Anemia related to myelodysplastic syndrome (MDS) excluding AML and CML Bone marrow biopsy < 5% blasts Erythropoietin level 100 or less After 2 months of treatment consider discontinuing if there is no significant increase in Hgb/Hcr and/or decreased transitisions							
☐ Treatment of anemia of selected chronic diseases Rheumatoid arthritis, systemic lupus erythematosus, inflammatory bowel diseases, and hepatitis C undergoing treatment							
Initial Hgb/Hct must be < 10 g/dl and < 30%. For all uses other than chemotherapy induced anemia titrate dose of darbepoetin to maintain a target Hgb of 10-12 g/dl and Hct of 30-36% by appropriate timed dose adjustment.							
Date of last ESA agent: Do not give Darbepoetin more frequently than once weekly							
Drug	Dose (mcg)		Route	Frequency	RN/Time		
Darbepoetin (Aranesp [©])	60 mcg		⊠ SubQ □ IV	ween			
Print Last Name RAMA SW WMY Print Last Name:							
Physician Signature / a. /		Nurse Signature:Date:					
DateTiov2.Time:		RBV INITTime: Faxed □ Time Faxed					