	From:	84	52946486			То	:3331148		Page
					,				3 -
	— .								
			•		•				
•								inal	
			•	Ш			1 4	JUEN	
				Ш	•			800	selsky
	IPI ORAL	NGE		IL.	1			104	•
	REGIO		Ch	1EN	OTHERA	ŀΡΥ	.]		
ſ	TO SE COMO	ETED BY PHYSI		<u>ЦО</u>	RDERS		-	D-	
17	Dato Wrilton:	ELED BA BHASI	CIAN:	Ц_		Patient Name: I	~-L		lient Lebal
			<u> </u>	Щ	Date o	Patient Name: PAdministration	<u>~ </u>	S R KADO	<u>#:6-13-5</u>
J.	Totocol / Ragin	ICerativ	15 COI	14	TNM 2		····	'-	
	77410				, — —				NKOA
٦	.,,,,,,,	r		.	Cycle	र्वा			· L_MKUA
	···				Pay			1	
	nous Access:		CARtra	į.	·			Ţ "	
	right	Weight Q	Actual	 - -	Body Surf	eca le-]	
- •	6 tt O in	148165	ideal Adjusted		Area (m)	- C M	c Level . Inimal	1	
<u> </u>		0	Dosing		1	1,103 M	Oderala		
	Orderar	CBCDIFF	BMP	T:	Мадлевіцт		gh		
	d Parameteros		<u></u>	 	1 Wagnesium	UA			
	C lose them:	WEC IRES (fr:	<u>an:</u> Pt	100	er than,	<i>U</i>	_		
200	ehemotherae					Hgb/Hct less t		SCr great	er than:
	V Fluide≀ N	e	/hr)	-			 -∤	<u>RPh Initials</u>	/ Nurse intital
/ ' -		ntivi		ሐፈ	og ir		- 1		
-		_ 					<u>c</u> [
			<u> </u>	 	<u>30m</u>	<u> </u>			 -
- 1				+					
<u> </u>				. [7		
		 -		+				,	
		<u> </u>		_			-+	-	╁ ── <u></u>
									
Solution)n, voluma sna					<u>></u>			
Golution agants	on, volume and Will be munder e nearest viol	Musion rate ara	Per manufak	Ouren Vithin	hospital guide 5% of calcula	ilines unless oth	ciwise ape	edifled. Chen	Dipers comp
Solution against the	in, voluma and Will be munda e nagrest vigi p	Musion rate ara	Per manufak	Wren Vithin dose	ncepital guide 5% of celeute s. Plasse adm	lines unless oth ted dose. Slolog litister chemothe	crwise spe losi agont	ediled. Chen s will be rour	neiherapeutic
1		Musion rate are d down to newest to if within 10% o	Per manufadi (viet stre (f viet site) of colculated	,	A PART ACIT	Deserolation	crwise spe lost agent rady in se	edilea. Civen s will be rour suenço lista	tolherapeutic 1984 down to 1 below.
1	on, voluma snd swil be munda e negreal vigi p emotherapy,	Musion rate are of down to neures ize if within 10% of Dose per that	Per manufadi I viai aize if w of colculaiad Dosa Reducti (maim	Фл°	Calculated	Dose Oispensed (Rounding to	rady in se	QUENÇO Rela	below.
· Che	emotherspy,	Musion rate are; of down to neures ize if within 10%; Dose per Unit (m². kg, AUC)	Per manufadi I viai aize if w of colculated Dosa Reducti	Фл°	A PART ACIT	Dose Dispensed (Rounding to	crwise spe loat agent rapy in se Route	ecified. Chen s will be rour guence lieta infuelon Reta	RPh /
· Che		Musion rate are; of down to neures ize if within 10%; Dose per Unit (m². kg, AUC)	Per manufadi I viai aize if w of colculaiad Dosa Reducti (maim	Фл°	Calculated	Dose Oispensed (Rounding to	rady in se	Infusion	below.
· Che	emotherspy,	Musion rate are; of down to neures ize if within 10%; Dose per Unit (m². kg, AUC)	Per manufadi I viai aize if w of colculaiad Dosa Reducti (maim	Фл°	Calculated	Dose Dispensed (Rounding to	rady in se	Infusion	RPh /
· Che	emotherspy,	Musion rate are; of down to neures ize if within 10%; Dose per Unit (m². kg, AUC)	Per manufadi I viai aize if w of colculaiad Dosa Reducti (maim	Фл°	Calculated	Dose Dispensed (Rounding to	rady in se	Infusion	RPh /
· Che	emotherspy,	Musion rate are; of down to neures ize if within 10%; Dose per Unit (m². kg, AUC)	Per manufadi I viai aize if w of colculaiad Dosa Reducti (maim	Фл°	Calculated	Dose Dispensed (Rounding to	rady in se	Infusion	RPh /
· Che	emotherspy,	Musion rate are; of down to neures ize if within 10%; Dose per Unit (m². kg, AUC)	Per manufadi I viai aize if w of colculaiad Dosa Reducti (maim	Фл°	Calculated	Dose Dispensed (Rounding to	rady in se	Infusion	RPh /
enty	omotherapy, ✓iO 300x	Musion rate are of down to neures ize if within 10% of Dose par Unit (m². kg. AUC)	Per manufaa i vial atze if w of celculated Doss Reducti (mg/m mg/ka, Al	on'	Calculated	Dose Dispensed (Rounding to	rady in se	Infusion	RPh /
Enty	omotherapy,	Infusion rate are of down to nearest ize if within 10% of Dose per Unit (m². kg. AUC)	per manufaa i vial alze (f w of colculated Doss Reducti (mg/m mg/ka, Al	on'	Calculated	Dose Dispensed (Rounding to	rady in se	Infusion	RPh /
Enty	omotherapy,	Infusion rate are of down to nearest ize if within 10% of Dose per Unit (m². kg. AUC)	per manufaa i vial alze (f w of colculated Doss Reducti (mg/m mg/ka, Al	φη* ΨC)	Calculated Doso	plater chemothe Dose Olspensed (Rounding to the completed by RPh)	Route	Intusion Rela	RPh / Nurse Initials
"If using a MD Name	a dose reduction (Print)	Infusion rate are of down to neures its infusion to neures its infusion 10% of the control of th	per manufaa i vial atze if w of celculated Dose Reducti (mg/m mg/ka, At	on'	Calculated Doso	Dose Olspensed (Rounding to be comploied by RPh)	Route Date/Figure	intuelon Rela	RPh / Nurse Initials
"If using a MD Name	a dose reduction (Print)	Infusion rate are of down to neures its infusion to neures its infusion 10% of the control of th	per manufaa i vial atze if w of celculated Dose Reducti (mg/m mg/ka, At	on'	Calculated Doso	inister chemothe Dose Otspensed (Rounding to be completed by RPh)	Route Route Date/Time	infusion Reta	RPh / Nurse Initials
"If using a MD Name RPh Name	a dose reduction (Print)	Infusion rate are of down to nearest ize if within 10% of Dose per Unit (m². kg. AUC)	per manufaa k val alke if w of celculated Does Reducti (mg/m mg/kg, Al MD Signe RN Signe RN Signe	on" C) clure clure	Calculated Doso	inister chemothe Dose Otspensed (Rounding to be completed by RPh)	Route Route Date/Time	intuelon Rela	RPh / Nurse Initials