
a
Separate LTR representation from its divergence profile
Left fraction × middle fraction = right fraction; each panel has its own percentage axis
LTR / all hits
Low-divergence / LTR
Low-divergence LTR / all hits
D. wrighti
D. organi
D. amphileucus
D. gvnigeusgwotli
D. aureatus
D. aeneus
D. mavrokoilius
D. kanawha
D. marmoratus
D. intermedius
D. orestes
D. abditus
D. welteri
D. auriculatus
D. lycos
D. bairdi
D. valtos
D. fuscus
D. valentinei
D. pascagoula
D. perlapsus
D. ocoee
D. apalachicolae
D. monticola
D. cheaha
D. balsameus
D. adatsihi
D. campi
D. santeetlah
D. tilleyi
D. carolinensis
D. anicetus
D. catahoula
D. conanti
0 10 20 30 40 50
0.0 2.5 5.0 7.5 10.0 12.5
0 2 4 6
Aligned hit bp (%)
Aquatic eggs
Terrestrial eggs + aquatic larvae
Direct development
b
Within-LTR signature, body size and strategy
34 species; strategy + SVL; omnibus P = 5.7e-04
Low-divergence LTR hit bp (% of LTR hit bp)
12.5
10.0
7.5
5.0
2.5
30
50
80
120
Maximum SVL (mm; log scale)
c
Phylogenetic strategy contrasts
Reference: aquatic eggs (n = 7); other groups: n = 24 and 3
Terrestrial eggs + aquatic larvae
Direct development
-1.0
-0.5
0.0
0.5
Difference in logit within-LTR fraction (95% CI)
Unadjusted
SVL adjusted