

a
Animal support
Numbers inside points: animals
42 morphology animals; 50 IOD animals
D. wrighti
D. amphileucus
D. gvnigeusgwotli
D. folkertsi
D. aeneus
D. mavrokoilius
D. kanawha
D. marmoratus
D. intermedius
D. orestes
D. ochrophaeus
D. welteri
D. auriculatus
D. bairdi
D. valtos
D. fuscus
D. brimleyorum
D. perlapsus
D. ocoee
D. apalachicolae
D. monticola
D. campi
D. tilleyi
D. anicetus
25%
50%
75%
100%
Largest animal share of objects
Nuclear IOD
Paired morphology
b
Equal-animal summaries
Change from released summaries
All observed animals retained
-5
0
Change in species summary (%)
Genome size
Cell area
Nucleus area
c
Conditional relationships under both summaries
Same full24 log scaling; 95% t intervals
All species (24)
Genome → cell | SVL
Cell → nucleus | SVL
Genome → nucleus | cell + SVL
Without aeneus + wrighti (22)
-0.5
0.0
0.5
1.0
Standardized coefficient
Released summaries
Equal-animal summaries
d
Model support under measurement and species sensitivity
Four candidates; all specify SVL upstream
250 animal/image/object resamples
Cell bridge
Genome bridge
Nucleus bridge
SVL alone
216/250 first
12/250 first
21/250 first
1/250 first
24 species-removal fits
24/24 first
0/24 first
0/24 first
0/24 first
0.0
0.5
1.0
Relative candidate weight