

Increase in venous [K+] During Hyperbaric Exposure Independent of Changes in pH or O<sub>2</sub> Concentration

Increase in venous [K+] During Hyperbaric Exposure

## Abstract

Plasma potassium regulation is important for function of numerous cells in the body. Changes in potassium levels during exposure to an increased O<sub>2</sub> concentration is thought to be the result of the changes in pH and increasing reactive oxygen species. However, the effects of hyperbaria on plasma potassium concentration are not well understood.

Eight subjects were exposed to 1.3 atmospheres absolute (ATA) of hyperbaric air for 90 minutes, 10-times (M-F) over 12-days. Another eight subjects were exposed to 100% oxygen at 1 ATA over the same interval. Four venous blood draws were taken. On day 1 the first draw was taken immediately preceding treatment and the second was taken immediately following treatment. The third draw was taken prior to the 10th treatment and the 4th draw was taken 72 hours post final treatment. We analyzed samples on a blood gas analyzer and performed statistical analysis using a paired Wilcoxon signed-rank test.

The concentration group saw strong trend towards an increase in the potassium concentration from  $4.09 \pm 0.12$  (mmol/L) to  $4.28 \pm 0.28$  (mmol/L) ( $p = 0.065$ ). In the hyperbaric group we see a significant increase in potassium concentration from  $4.19 \pm 0.26$  (mmol/L) to  $4.55 \pm 0.27$  (mmol/L) ( $p = 0.0068$ ). In the concentration group we also see a significant increase in pH concentration from  $7.37 \pm 0.03$  to  $7.39 \pm 0.01$  ( $p = 0.021$ ). A similar significant increase is not seen in the hyperbaric group.

These findings suggest that changes in potassium concentration in response to hyperbaria are not the result of oxygen concentration nor pH. Possible explanations include increased nitrogen levels due to hyperbaric air, increased CO<sub>2</sub> concentration in hyperbaric chamber or changes in the activity of Na<sup>+</sup>, K<sup>+</sup> ATPase pumps at the cellular level which may be a homeostatic response to combat pulmonary edema.

Keywords: Potassium, Hyperbarics, Hyperoxia

## Introduction

Plasma potassium levels are dependent on a multitude of factors and result from the interplay of intracellular and extracellular changes and intake and excretion rates.<sup>1-3</sup> Plasma potassium levels are heavily regulated to fall within a narrow range of 3.5 to 5.0 mmol per liter and maintenance at these

concentrations are of extreme importance to the function of many organs in the body.<sup>3</sup> This is underscored by the fact that both hypokalemia or hyperkalemia lead to an increased death rate in certain conditions.<sup>5,6</sup> This places importance on the determination of pathways of regulation and disease that lead to changes in plasma potassium levels. Factors investigated in this study that may lead to changes in plasma potassium levels, but remains understudied, are the effect of  $O_2$  concentration and increased pressure.

Modulation of inspired  $O_2$  concentration has been used as an effective treatment for many conditions. The two main methods to administer this concentration increase are by directly varying concentration of inspired gases or by varying the ambient pressure. By combining these to aspects of concentration, three of the main modes of  $O_2$  administration are obtained. These are hyperbaric  $O_2$ , concentrated  $O_2$  and hyperbaric treatment. According to the UHMS hyperbaric oxygen therapy ( $HBO_2$ ) is defined as exposure to near 100%  $O_2$  while inside a pressurized chamber at greater than sea level pressure.<sup>7</sup> As of July 2021 the FDA has cleared  $HBO_2$  for 13 different conditions.<sup>8</sup> In contrast hyperbaric treatment is exposure to an increase in pressure without an increase in administered gas  $O_2$  concentration. Finally concentrated oxygen is the exposure to  $O_2$  gas with a higher concentration of  $O_2$  than normal air. Concentrated oxygen and hyperbaric treatment were the two treatments administered in this study.

Effects of hyperoxia on plasma potassium levels remain ambiguous.<sup>2</sup> A few studies have demonstrated an increase in plasma potassium as the result of hyperoxic exposure while others have found no similar increase.<sup>2,9</sup> However research has shown that hyperoxia can effect voltage-gated  $K^+$  channels in humans<sup>10-12</sup> One group found that long-term exposure to hyperoxia can result in a decrease in activity of voltage-gated  $K^+$  channels in lung tissue.<sup>10,11</sup> Where as another study looking at voltage-gated  $K^+$  channels of the heart and saw a significant reduction of oxygen-sensitive Kv1.5 potassium channel protein expression.<sup>12</sup> Internal potassium homeostasis and the distribution of intracellular and extracellular potassium in the body is primarily the result of ion-exchange pumps and passive efflux.<sup>1,3</sup> Hypoxia has also been shown to cause an increase in arterial potassium levels and cause changes in expression of Voltage gated  $K^+$  channel in pulmonary arterial myocytes through  $HIF1 - \alpha$  indicating some pathway through which potassium levels are modulated via changes in  $O_2$  concentration.<sup>13-16</sup> Therefore changes in activity or expression of these channels offers a potential linkage between effects hyperoxia on plasma potassium.

The main goal of this study was to determine the similarities and differences in effects of concentrated  $O_2$  and hyperbaria on the blood metabolites, specifically plasma potassium levels.

## Methods

Healthy adult subjects were recruited via REDCap using a University of Wisconsin’s email list. A questionnaire was administered to determine characteristics of individuals such as pre-existing conditions and age. Individuals who responded to the survey were selected at random contacted via phone. Participants in the study were separated into two groups. The first group received hyperbaric treatment in altitude sickness bags (n=14) while the second group received concentrated oxygen at room pressure (n=12).

Although treatment varied between groups, treatment schedule remained consistent and was the following. Subjects arrived Monday and height and weight were determined. Blood collected via venipuncture before treatment was used for a baseline metabolic panel and analyzed on arterial blood gas (ABG) machine. Following blood draws subjects underwent 1.5 hours of their respective treatments. After treatment blood was collected to determine acute response in blood metabolites. Following this, treatment was given Tuesday through Friday of the following week, taking a break for weekends where no treatment was administered. A blood draw and ABG analysis was performed before the tenth and final treatment. After a weekend of no treatment following the final treatment a blood draw occurred and subsequent analysis was performed.

A total of 36 metabolic indicators were reported by the ABG machine. Indicators were compared pairwise between the four draws. Statistical analysis and data visualization was performed using R (The R Foundation) and paired student t-tests.

## Results

Tables one and two display values for classical ions of interest in potassium homeostasis for concentrated  $O_2$  and hyperbaric air respectively. The concentrated  $O_2$  group saw a significant increase in pH from baseline levels of  $7.374 \pm 0.029$  to post exposure levels of  $7.393 \pm 0.014$  ( $p=0.023$ ) while the hyperbaric group saw no similar significant increase. Potassium levels were also increased in both the concentrated  $O_2$  group and the hyperbaric oxygen group immediately post exposure. In the concentrated group the potassium levels were increased from  $4.09 \pm 0.12$  pre-exposure to  $4.28 \pm 0.28$  post-exposure ( $p=0.046$ ). Whereas in the hyperbaric group a greater increase was observed with pre-exposure levels at  $4.19 \pm 0.26$  and post-exposure levels at  $4.55 \pm 0.27^*$  ( $p=0.007046$ ). Calcium levels were also increased from baseline levels of  $1.19 \pm 0.04$  for both immediately post-exposure at  $1.21 \pm 0.03$  ( $p=0.043$ ) and 24 hours post-exposure at  $1.22 \pm 0.02^*$  ( $p=0.047$ ) however no similar change was observed in the hyperbaric group.

Table 1: Metabolite concentrations for concentrated oxygen group from pre, post, day 10 and final blood draws. Significance denoted by (\* < 0.05), (\*\* < 0.001)

Metabolite	Pre (n=13)	Post (n=13)	Day 10 (n=10)	Final (n=9)
<i>pH</i>	$7.374 \pm 0.029$	$7.393 \pm 0.014^*$	$7.368 \pm 0.033$	$7.384 \pm 0.027$
<i>pCO<sub>2</sub></i> ( <i>mmHg</i> )	$48.5 \pm 5.6$	$47.1 \pm 4.3$	$44.7 \pm 6.3$	$38.8 \pm 8.7^{**}$
<i>pO<sub>2</sub></i> ( <i>mmHg</i> )	$26.3 \pm 8.2$	$21.8 \pm 5.2$	$30.5 \pm 19.6$	$26.9 \pm 10.2$
<i>Na</i> + ( <i>mmol/L</i> )	$139.6 \pm 0.9$	$139.2 \pm 1.2$	$139.4 \pm 1.7$	$138.6 \pm 3.9$
<i>K</i> + ( <i>mmol/L</i> )	$4.09 \pm 0.12$	$4.28 \pm 0.28^*$	$4.28 \pm 0.44$	$4.29 \pm 0.27$
<i>Cl</i> – ( <i>mmol/L</i> )	$106.9 \pm 1.5$	$107.4 \pm 1.3$	$105.1 \pm 1.7^*$	$106.4 \pm 3.4$
<i>Ca</i> + ( <i>mmol/L</i> )	$1.19 \pm 0.04$	$1.21 \pm 0.03^*$	$1.22 \pm 0.02^*$	$1.21 \pm 0.04$

Table 2: Metabolite concentrations for hyperbaric group from pre, post, day 10 and final blood draws. Significance denoted by (\* < 0.05), (\*\* < 0.001)

Metabolite	Pre (n=11)	Post (n=11)	Day 10 (n=11)	Final (n=11)
<i>pH</i>	$7.384 \pm 0.033$	$7.408 \pm 0.029$	$7.387 \pm 0.049$ (n=8)	$7.387 \pm 0.037$
<i>pCO<sub>2</sub></i> ( <i>mmHg</i> )	$35.8 \pm 6.8$	$34.5 \pm 4.5$	$36.6 \pm 10.2$ (n=8)	$39.3 \pm 10.0$
<i>pO<sub>2</sub></i> ( <i>mmHg</i> )	$42.9 \pm 22.9$ (n=8)	$25.7 \pm 6.6$ (n=7)	$41.6 \pm 37.4$ (n=10)	$43.8 \pm 31.8$
<i>Na</i> ( <i>mmol/L</i> )	$134.0 \pm 1.2$	$139.7 \pm 1.3$	$139.2 \pm 1.2$	$139.4 \pm 1.3$
<i>K</i> + ( <i>mmol/L</i> )	$4.19 \pm 0.26$	$4.55 \pm 0.27^*$	$4.15 \pm 0.24$	$4.23 \pm 0.25$
<i>Cl</i> – ( <i>mmol/L</i> )	$106.5 \pm 1.6$	$106.4 \pm 1.7$	$106.2 \pm 2.0$	$106.2 \pm 2.0$
<i>Ca</i> + ( <i>mmol/L</i> )	$1.24 \pm 0.03$	$1.25 \pm 0.03$	$1.22 \pm 0.04$	$1.22 \pm 0.05$

## Discussion

Based on the fact that the classical metabolites implicated in plasma potassium levels experienced no significant fluctuations this hints at a possible novel

relationship between plasma potassium and hyperbaric treatment. However, limitations of this study mean that an exact explication for the observed plasma potassium levels in the hyperbaric group was not be determined. In spite of this fact several possible explanations based on findings in the literature and conditions present in this study arose and were as follows: an increase in hyperbaric chamber  $CO_2$  resulting in mild respiratory acidosis, increase in  $N_2$  and creation of reactive nitrogen species (RNS) in tissue, potassium mediated fluctuations in vasoconstriction/dilation in response to changes in pressure and changes in potassium channel expression via  $HIF - 1\alpha$ .

Correlations between respiratory acidosis and an increase in plasma potassium have been reported with few exceptions.<sup>2</sup> During hyperbaric exposure subjects experienced an increase of levels of  $CO_2$  by =FIXME:INSERT  $CO_2$  LEVELS IN CHAMBER==. Subject expired gases and a lack of outflow of those respired gases in the altitude sickness bags is most likely responsible for the increase in  $CO_2$  observed in chamber.

Further limitations of this study include lack of dietary restrictions on subjects, single Pre vs Post draw time point and variations between treatment cohorts.

## References

- (1) Aronson, P. S.; Giebisch, G. Effects of pH on Potassium: New Explanations for Old Observations. *Journal of the American Society of Nephrology* **2011**, *22*, 1981–1989.
- (2) Adrogué, H. J.; Madias, N. E. Changes in Plasma Potassium Concentration During Acute Acid-Base Disturbances. *The American Journal of Medicine* **1981**, *71* (3), 456–467. [https://doi.org/10.1016/0002-9343\(81\)90182-0](https://doi.org/10.1016/0002-9343(81)90182-0).
- (3) Gumz, M. L.; Rabinowitz, L.; Wingo, C. S. An Integrated View of Potassium Homeostasis. *New England Journal of Medicine* **2015**, *373* (1), 60–72. <https://doi.org/10.1056/NEJMr1313341>.
- (4) Youn, J. H.; McDonough, A. A. Recent Advances in Understanding Integrative Control of Potassium Homeostasis. *Annual Review of Physiology* **2009**, *71* (1), 381–401. <https://doi.org/10.1146/annurev.physiol.010908.163241>.
- (5) Torlén, K.; Kalantar-Zadeh, K.; Molnar, M. Z.; Vashistha, T.; Mehrotra, R. Serum Potassium and Cause-Specific Mortality in a Large Peritoneal Dialysis Cohort. *Clinical Journal of the American Society of Nephrology* **2012**, *7* (8), 1272–1284. <https://doi.org/10.2215/CJN.00960112>.
- (6) Goyal, A.; Spertus, J. A.; Gosch, K.; Venkitachalam, L.; Jones, P. G.; Van den Berghe, G.; Kosiborod, M. Serum Potassium Levels and Mortality in Acute Myocardial Infarction. *JAMA* **2012**, *307* (2), 157–164. <https://doi.org/10.1001/jama.2011.1967>.

- (7) *Hyperbaric Oxygen Therapy Indications*; Moon, R. E., Ed.; Undersea; Hyperbaric Medical Society, 2019; Vol. 14th.
- (8) Hyperbaric Oxygen Therapy: Get the Facts. *Consumer Updates*, 2021.
- (9) Fraley, D. S.; Adler, S. Isohydric Regulation of Plasma Potassium by Bicarbonate in the Rat. *Kidney International* **1976**, *9* (4), 333–343. <https://doi.org/10.1038/ki.1976.39>.
- (10) Zyrianova, T.; Lopez, B.; Olcese, R.; Belperio, J.; Waters, C. M.; Wong, L.; Nguyen, V.; Talapaneni, S.; Schwingshackl, A. K2p2.1 (TREK-1) Potassium Channel Activation Protects Against Hyperoxia-Induced Lung Injury. *Scientific Reports* **2020**, *10* (1), 22011. <https://doi.org/10.1038/s41598-020-78886-y>.
- (11) Schwingshackl, A.; Teng, B.; Ghosh, M.; West, A. N.; Makena, P.; Gorantla, V.; Sinclair, S. E.; Waters, C. M. Regulation and Function of the Two-Pore-Domain (K2p) Potassium Channel Trek-1 in Alveolar Epithelial Cells. *American Journal of Physiology-Lung Cellular and Molecular Physiology* **2012**, *302* (1), L93–L102. <https://doi.org/10.1152/ajplung.00078.2011>.
- (12) Chapalamadugu, K. C.; Panguluri, S. K.; Bennett, E. S.; Koliputi, N.; Tipparaju, S. M. High Level of Oxygen Treatment Causes Cardiotoxicity with Arrhythmias and Redox Modulation. *Toxicology and Applied Pharmacology* **2015**, *282* (1), 100–107. <https://doi.org/10.1016/j.taap.2014.10.019>.
- (13) Barlow, C. W.; Qayyum, M. S.; Davey, P. P.; Paterson, D. J.; Robbins, P. A. Effect of Hypoxia on Arterial Potassium Concentration at Rest and During Exercise in Man. *Exp Physiol* **1994**, *79* (2), 257–260. <https://doi.org/10.1113/expphysiol.1994.sp003759>.
- (14) Whitman, E. M.; Pisarcik, S.; Luke, T.; Fallon, M.; Wang, J.; Sylvester, J. T.; Semenza, G. L.; Shimoda, L. A. Endothelin-1 Mediates Hypoxia-Induced Inhibition of Voltage-Gated K<sup>+</sup> Channel Expression in Pulmonary Arterial Myocytes. *American Journal of Physiology-Lung Cellular and Molecular Physiology* **2008**, *294* (2), L309–L318. <https://doi.org/10.1152/ajplung.00091.2007>.
- (15) Dong, Q.; Zhao, N.; Xia, C. K.; Du, L. L.; Fu, X. X.; Du, Y. M. Hypoxia Induces Voltage-Gated K<sup>+</sup> (Kv) Channel Expression in Pulmonary Arterial Smooth Muscle Cells Through Hypoxia-Inducible Factor-1 (HIF-1). *Bosn J Basic Med Sci* **2012**, *12* (3), 158–163. <https://doi.org/10.17305/bjbms.2012.2463>.
- (16) Semenza, G. L. Regulation of Oxygen Homeostasis by Hypoxia-Inducible Factor 1. *Physiology* **2009**. <https://doi.org/10.1152/physiol.00045.2008>.