

Rapid #: -19163701

CROSS REF ID: **3667265**

LENDER: **NTE :: Main Library**

BORROWER: **GZM :: Memorial Library**

TYPE: Article CC:CCG

JOURNAL TITLE: Undersea & hyperbaric medicine

USER JOURNAL TITLE: Undersea & hyperbaric medicine

ARTICLE TITLE: Decrease in dysbaric osteonecrosis severity as a result of 45-minute oxygen pre-breathe

ARTICLE AUTHOR: Lamers, Jacob

VOLUME: 48

ISSUE: 3

MONTH:

YEAR: 2021

PAGES: 279-286

ISSN: 1066-2936

OCLC #:

Processed by RapidX: 6/10/2022 1:50:40 PM

This material may be protected by copyright law (Title 17 U.S. Code)

RESEARCH ARTICLE

Decrease in dysbaric osteonecrosis severity as a result of 45-minute oxygen pre-breathe

Jacob Lamers^{1,2}; Kent MacLaughlin^{1,2}; Dave Pegelow^{1,2}; Marlowe Eldridge, MD^{1,2}; Aleksey Sobakin DVM, PhD^{1,2}

¹ Department of Pediatrics, Division of Critical Care, University of Wisconsin School of Medicine and Public Health Madison, Wisconsin U.S.

² John Rankin Laboratory of Pulmonary Medicine, University of Wisconsin, Madison, Wisconsin U.S.

CORRESPONDING AUTHOR: Aleksey Sobakin – sobakin@wisc.edu

ABSTRACT

Sudden decompression can result in bubble formation as the result of nitrogen gas (N₂) dissolved in tissue during disabled submarine escape (DISSUB). This may cause dysbaric osteonecrosis (DON), a condition in long bones where bubbles in fatty marrow result in ischemia and necrosis. Previous research has shown that oxygen (O₂) pre-breathe of two hours resulted in a reduction of DON; however, effects of shorter O₂ pre-breathe remain uncertain.

This study's aim was to understand the effect of shorter lengths of O₂ pre-breathe. Eight adult Suffolk ewes (89.5 ± 11.5 kg) were exposed to 33 feet of seawater (fsw) for 24 hours. They were placed randomly into four groups and exposed to either 45, 30 or 15 minutes of O₂ (91-88%) pre-breathe; the controls received none.

They were then rapidly decompressed. Alizarin complexone was later injected intravenously to visualize the extent of DON in the right and left long bones (radii, tibiae, femur and humeri). The 30- and 15-minute pre-breathe groups saw the greatest deposition. There was significant decrease of variance in the 45-minute group when compared with all other treatments, suggesting that 45 minutes of O₂ pre-breathe is required to effectively increase confidence in the reduction of DON. Similar confidence was not reflected in the 30-minute and 15-minute groups: 45 minutes of pre-breathe was the minimum amount needed to effectively prevent against DON in DISSUB escape at 33 fsw. However, future research is needed to determine how to calculate effective dosages of O₂ pre-breathe to prevent DON in any given scenario. ■

INTRODUCTION

Dysbaric osteonecrosis (DON) is the formation of necrotic lesions in the long bone fatty marrow as a result of a sudden drop in barometric or hydrostatic pressure, causing bubble formation in the intramedullary space. DON can arise after a single exposure to a sudden decrease in environmental pressure [1] but it is more common in individuals who experience multiple exposures, including compressed-air workers [2,3] diving fishermen [4,5] and divers [6].

Rapid evacuation during a DISSUB scenario has also been shown to lead to DON [7]. Clinically, DON may manifest as chronic pain in the hips, knees and shoulders, and in extreme cases, total joint failure [8].

Among groups regularly exposed to pressures required for the formation of DON lesions, military divers experience the lowest rates of DON [9-11]. The highest rates of DON were documented in Turkish sponge

divers and Hawaiian fishermen, with prevalence rates of 71% and 65% respectively [4,5]. These values mirror other diving fisherman communities that are at increased risk because of a lack of education about, and resources for, the prevention and treatment of DON [4,5,9]. Compressed-air workers are also at risk of developing DON, with prevalence rates of around 17% [12,13]. The true prevalence of DON may be inaccurate for some populations because detection and reporting are often expensive and challenging [10,13].

The main risk factors that predispose a person to the development of DON are multiple exposures to radical drop in pressure and/or greater maximal diving depth [13,14]. Other potential risk factors for DON are higher levels of plasminogen activator inhibitor-1, fat embolisms and hyperlipidemia [14,15]. Like many other forms of osteonecrosis the underlying mechanism for DON is not entirely understood. This provides certain

KEYWORDS: dysbaric; osteonecrosis; decompression; DISSUB

challenges when trying to determine what pre-existing habits and conditions can increase the chances of developing these lesions [8].

The University of Wisconsin-Madison Sheep Model of decompression injury developed by Lehner, et al. (1997) is a reliable surrogate for DON in humans [16]. The ability to produce accelerated decompression conditions in adult sheep provides an effective human surrogate, as these animals have similar bone size, structure, and fatty marrow distribution to humans [16]. The sheep have similar body weights as well as metabolic and tissue perfusion rates which are factors that control the amount of dissolved gas. This means the likelihood of bubble formation and radiographic and histological findings of DON are remarkably similar to those of humans [16].

When exposed to high ambient pressure, nitrogen (N_2) dissolves in body tissue relative to the partial pressure [17]. Factors such as time of exposure, tissue perfusion and solubility of gas in tissue play vital roles in determining the N_2 load [17]. When studying DON it becomes clear that tissue perfusion rates and solubility of N_2 in fatty tissue play an important role in developing lesions. Fatty marrow in long bones has a low perfusion rate, and N_2 is five times more soluble in fat than in aqueous tissue. This dissolved N_2 can form bubbles on active hydrophobic surfaces when ambient pressure is reduced [18]. One widely accepted theory for the formation of these lesions is that extravascular bubbles lead to an increase in intramedullary pressure [13,17,19]. Blood vessels then experience vascular stasis as a result of the Starling resistor mechanism [19]. If the intramedullary pressure is not reduced, and the bubbles do not disappear, fatty marrow and bone begin to experience ischemia, which can lead to bone necrosis [19].

It has also been proposed that intravascular bubbles may block the sinusoids of the fatty bone marrow, leading to ischemia and necrosis [13]. Another hypothesis is that DON occurs because N_2 bubbles can lead to the release of liquid fat and thromboplastin [15]. This can cause an increase in blood coagulation in intravascular regions, causing vessel blockage [15]. Although the exact mechanism for necrosis that produces DON lesions is not completely understood, it is known that increased environmental pressure causes an increase in N_2 saturation [17,19]. This leads to increased saturation of N_2 in fatty marrow tissue, which results in the formation of gas bubbles when pressure is rapidly decreased [15,17,18]. These bubbles are then responsible for the formation of DON lesions [15, 17-19].

We hypothesize that O_2 washout treatment in the form of O_2 pre-breathe is an effective form of prevention for DON. The aim of this study is to determine the effect of shorter durations of O_2 pre-breathe on DON at 33 fsw.

METHODS

The Institutional Animal Care and Use Committee of the University of Wisconsin reviewed and approved this protocol to perform DISSUB sheep O_2 pre-breathe trials for Naval Sea System Command that followed guidelines for medical research in relation to animals set forth by the Helsinki Animal Committee.

Subjects

Eight adult female Suffolk ewes (89.5 ± 11.5 kg) were exposed to 33 feet of seawater (fsw) for 24 hours in a high-pressure chamber at the University of Wisconsin Biotron Laboratory. In order to avoid acclimatization sheep selected for this study had no prior exposure to a hyperbaric environment. Each animal was randomized to one of four groups ($n = 2$):

- Group 1 received 45 minutes of O_2 (91-88%) pre-breathe;
- Group 2 received 30 minutes of O_2 (91-88%) pre-breathe;
- Group 3 received 15 minutes of O_2 (91-88%) pre-breathe; and
- Group 4 controls received only compressed air before dropout decompression.

Diving procedures

Sheep enter the hyperbaric chamber while experiencing normal surface pressure. The chamber is then compressed at a rate of 30 feet per minute to a final pressure of 33 fsw. The sheep then remain in the chamber for 24 hours. After 24 hours, each group is given its respective treatment. The sheep then undergo decompression at the same rate that they were compressed, reaching surface pressure in 60-90 seconds.

Analysis

Alizarin complexone, a fluorochemical used to visualize areas of bone remodeling [20], was injected into sheep one week prior to euthanasia. MRI images of long bones prior to exposure and then five weeks post-treatment were used to confirm that areas of alizarin complexone deposition were indeed DON lesions. The tibia, humerus, femur, and radius were removed during autopsy and segmented longitudinally to observe areas of DON

lesions. The long bones were preserved through fixation with a 10% solution of neutral buffered formalin to preserve the bone and alizarin color. Digital images were taken to analyze the gross pathology of long bones. Raw images were processed with ImageJ Scion Image software (Scion Corporation, Frederick, Maryland) and the software extension WEKA segmentation (University of Waikato, Verena Kaynig, Johannes Schindelin). The images were then used to calculate the area of alizarin complexone deposition and the total bone area as shown in Figure 1. These values were used to calculate total bone percent alizarin complexone deposition, as well as percent deposition values for the proximal, distal, and middle thirds of each bone.

Statistical analysis

All statistical work was done in Prism 9 (GraphPad Software) and R (The R Foundation). Percent alizarin deposition was compared using a T-test with Wilcoxon correction to check for statistically differences between groups for each of the bones studied. F-test was used to determine significance of variation between groups in each of the four bones. Descriptive statistics were used to determine mean, standard deviations and variation values.

RESULTS

When looking across the total bone at alizarin complexone deposition a trend was evident in the 15-minute and 30-minute oxygen pre-breathe groups: Both saw an increase in the percent deposition when compared to the 45-minute group (Figure 2). This, however, is not a statistically significant increase, with no bone showing significant increases between groups when compared by t-test. But when looking at Figure 3, the radius and humerus show a significant decrease in variance of DON occurrence when comparing the 45-minute group to the 30-minute, 15-minute and controls ($p < 0.0001$, $p < 0.05$). Both the femur and tibia saw significant decreases in variance when comparing the 45-minute group to the 30-minute and 15-minute groups, ($p < 0.0005$), but no significant differences were observed between the 45-minute group and the controls. In all bones the 15-minute and 30-minute groups showed no significant differences in variance of deposition. The radius and tibia saw the highest rates of lesions. The radius mean depositions were 0.05 ± 0.07 in the 45-minute group, 4.86 ± 5.00 in the 30-minute group, 13.15 ± 14.79 in the 15-minute group and 6.377 ± 12.36 for the controls.



The tibia mean depositions were 0.13 ± 0.18 in the 45-minute group, 4.75 ± 5.99 in the 30-minute group, 6.78 ± 11.23 in the 15-minute group and 0.61 ± 0.4 for the controls. There was, however, no significant increase in deposition between any of the bones.

When evaluating the alizarin complexone deposition in the proximal third, a similar trend was observed. Figure 4 shows what looks like a trend toward an increase in deposition in the 30- and 15-minute groups, which is not significant. When comparing the 45-minute to the 15-minute, 30-minute and the controls the only bone that showed significant decreases in the variance in this third was the humerus ($p < 0.05$). In all other bones both the 45-minute group and the controls saw a significant decrease in variance of deposition when compared to the 15-minute and 30-minute groups. The bones that saw




Figure 2: Percent deposition across the entire extent of the femur, humerus, radius and tibia. Treatment was 33 fsw for every sheep with one of four durations of pre-breathe. The groups were as follows: 45 minutes (n=2); 30 minutes (n=2); 15 minutes (n=2); and the controls (n=2).




Figure 3: Variance of percent deposition across the entire extent of the femur, humerus, radius and tibia. Treatment was 33 fsw for every sheep with one of four durations of pre-breathe.

the most DON in the proximal third was the humerus and tibia. The humerus saw mean percent deposition rates of 0.322 ± 0.631 in the 45-minute group, 8.91 ± 7.69 in the 30-minute group, 4.06 ± 4.78 in the 15-minute group and 2.13 ± 2.87 in the control group. In the tibia the mean percent deposition rates were 0.0441 ± 0.0874 in the 45-minute group, 5.55 ± 6.63 in the 30-minute group, 7.82 ± 12.5 in the 15-minute group and 0.708 ± 0.932 in the control group. The humerus saw a significant increase in deposition in this third when compared to the femur ($p < 0.05$). However, all other differences in deposition between bones in this third were not significant.

Figure 5 shows that when compared to the distal and proximal thirds, the middle saw the least deposition in all four treatment groups. The humerus and tibia remained almost completely unaffected in this area, with the highest percent depositions occurring in the tibia of the 30-minute and 15-minute groups at rates of 1.71 ± 2.58 and 1.6 ± 1.99 , respectively. The femur saw deposition at rates of 4.5 ± 8.98 in the 15-minute group, but all other groups remained unaffected. Of the four bones, the radius was the only bone to show significant signs of deposition, but these differences were not statistically significant. The 45-minute group showed significant decrease in variance of deposition when compared with the 15-minute, 30-minute and control groups ($p < 0.0005$). The 30-minute group also saw a significant decrease in variance when compared with the 15-minute and the control groups ($p < 0.05$). The 15-minute and the




Figure 4: Percent deposition for the proximal third of the femur, humerus, radius and tibia. Treatment was 33 fsw for every sheep with one of four durations of pre-breathe. The groups were as follows: 45 minutes (n=2); 30 minutes (n=2); 15 minutes (n=2); and the controls (n=2).





Figure 5: Percent deposition for the middle third of the femur, humerus, radius and tibia. Treatment was 33 fsw for every sheep with one of four durations of pre-breathe. The groups were as follows: 45 minutes (n=2); 30 minutes (n=2); 15 minutes (n=2); and the controls (n=2).

controls did not show significant differences in variances. In the radius the mean percent deposition rates were 0.0166 ± 0.0164 in the 45-minute group, 1.39 ± 1.26 in the 30-minute group, 7.12 ± 7.93 in the 15-minute group and 3.22 ± 6.4 in the control group. However, most of the deposition in radius in this third occurred near the distal region.

The trend observed in the proximal third and when looking at the total bone was also seen in the distal region as well (Figure 6). The 45-minute group saw the lowest amounts of deposition when compared to the other treatment groups. These differences were not significant. The 45-minute group saw significantly lower variance when compared to the 15-minute, 30-minute and control groups in the radius and humerus ($p < 0.005$). In the femur and tibia, the 45-minute group saw significantly lower variance when compared to the 15-minute and the 30-minute groups ($p < 0.005$). However, in these bones the 45-minute variance was not significantly lower than in the control groups. The two most affected bones in this third were the radius and tibia. In the radius mean percent deposition rates were 0.11 ± 0.169 in the 45-minute group, 8.61 ± 14.6 in the 30-minute group, 25.4 ± 30.2 in the 15-minute group and 12.6 ± 24.3 in the control group. As for the tibia the mean percent deposition rates were 0.405 ± 0.729 in the 45-minute group, 6.8 ± 11.2 in the 30-minute group, 9.57 ± 17.3 in the 15-minute group and 0.696 ± 0.75 in the control group. These differences between bones were not significant.



DISCUSSION

The 30-minute, 15-minute and control groups all showed large amounts of deposition in at least one of the bones studied. However, because of the small n value for these groups and the sporadic nature of DON lesion formation these differences did not appear to be significant. In spite of this fact, significantly high variability in DON lesions in at least one of the bones studied for all groups other than the 45-minute suggests there is confidence in the preventative capabilities of 45-minutes of O₂ pre-breathe under this study's conditions. We hypothesize that the variance observed in the 30-minute, 15-minute and control groups is indicative of an ineffective duration of O₂ pre-breathe which allowed for varying rates of DON based on a multitude of factors; these include body weight, bone composition, duration of dive and time at depth. The 45-minute pre-breathe was, however, enough at this study's depth and time to overcome these factors and increase the confidence in the reduction of DON. The fact that the 15-minute, 30-minute, and controls did not see a similar reduction in variance means that confidence in their preventative capabilities is not justified.

The increased N₂ partial pressure, high solubility and low perfusion rates in fatty marrow means that N₂ does not rapidly diffuse out of marrow. Because bubble formation is the main factor determining the development of DON, removing this dissolved N₂ may be an effective form of prevention. According to Dalton's law, raising the partial pressure of O₂ in blood through O₂ pre-breathe will lower the partial pressure of N₂. This lowering of the

partial pressure of N₂ means it can diffuse more rapidly from highly saturated fatty marrow. The lower concentration of N₂ should therefore decrease the probability of bubble formation. A previous study done in the UW Sheep Model showed that two hours of O₂ pre-breathe greatly reduced the formation of DON lesions.

In that study, using similar methods, sheep were subjected to 15 minutes, one hour and two hours of O₂ pre-breathe at a depth of 60 fsw, while the sheep in our study were subjected to 33 fsw. (The 15-minute and one-hour groups were ineffective at preventing the formation of DON lesions.) However two hours saw a significant reduction in DON lesions. The discrepancy in the time of pre-breathe needed is most likely the result of the difference in pressure. As pressure increases during a dive the amount of dissolved gas in the tissue and blood also increases. The increased pressure in the previous study would have resulted in a greater amount of dissolved N₂. The exact amount of time required for N₂ to be flushed out completely remains uncertain and most likely varies between individuals because of factors that control the amount of dissolved N₂ and the rate at which N₂ can be washed out. In spite of this our results suggest that greater dive depth and therefore greater dissolved N₂ mean a greater length of pre-breathe is required. However, without further research into the exact relationship between these two factors this remains uncertain.

Confounding variables appear to have an impact on our results, most notably in the control group. Both the tibia and femur showed

no significant difference in variance between the 45-minute group and the controls. The fact that the control group saw a lower extent of DON lesions compared to the 15-minute and 30-minute groups appears to contradict the claim that 45-minute pre-breathe is causing the reduction of variance. However, DON does not uniformly affect those who experience rapid drops in ambient pressure. For example, when comparing DON to decompression sickness – another disease caused by bubble formation as the result of a rapid reduction in pressure – it is generally accepted that not all those who experience decompression sickness also experience DON [24]. In spite of this the UW Sheep model has been shown to accurately recreate DON lesions like those seen in humans [16]. This leads us to believe that the control results are a product of our small sample size and not indicative of larger relationship where both the no oxygen pre-breathe and 45-minute pre-breathe see similar preventative qualities in our study environment. On the contrary, the lower percent deposition in the controls is most likely the cause of their lower weights compared to the other treatment groups. As previously mentioned, the mechanism for DON is not entirely understood, and a complete list of risk factors remains uncertain. Variables such as higher body mass index (BMI), fat mass, concentration of lipids in blood and plasminogen activator inhibitor-1 have all been linked to increases in dysbaric disease [14,25-27]. However

without further study the effect of these factors remains uncertain. More research is needed to better understand the effects of BMI and bone composition along with the previously mentioned effects of pressure and depth.

CONCLUSIONS

This study's finding of low rates of DON lesion formation in the 45-minute pre-breathe group as well as a significant reduction in variance of DON lesions compared to controls and other pre-breathe groups suggests the possible preventative capability of O₂ pre-breathe. However, a larger sample size is required to determine the efficacy of O₂ pre-breathe treatment. Moreover, in order to effectively determine the amount of O₂ pre-breathe needed to prevent DON lesions for a specific individual, a better understanding of the relationship between factors including but not limited to BMI, bone composition, dive depth and time at depth is needed. ■

Acknowledgments

Thank you to NAVSEA for their continued support of our work and to everyone at the Eldridge lab for all of their help during this project. Special thanks to Greg Barton for helping us find a story.

Funding details

This work was supported by NAVSEA under grant N0463A-12-C-0004.

REFERENCES

1. Nellen JR, Kindwall EP. Aseptic necrosis of bone secondary to occupational exposure to compressed air: roentgenologic findings in 59 cases. *Am J Roentgenol Radium Ther Nucl Med.* 1972;115(3):512-524.
2. Kindwall EP, Nellen JR, Spiegelhoff DR. Aseptic necrosis in compressed air tunnel workers using current OSHA decompression schedules. *J Occup Med.* 1982;24(10):741-745.
3. Gregg PJ, Walder DN. Caisson disease of bone. *Clin Orthop Relat Res.* 1986(210):43-54.
4. Wade CE, Hayashi EM, Cashman TM, Jr., Beckman EL. Incidence of dysbaric osteonecrosis in Hawaii's diving fishermen. *Undersea Biomed Res.* 1978;5(2):137-147.
5. Toklu AS, Cimsit M. Dysbaric osteonecrosis in Turkish sponge divers. *Undersea Hyperb Med.* 2001;28(2):83-88.
6. Elliott DH, Harrison JA. Bone necrosis--an occupational hazard of diving. *J R Nav Med Serv.* 1970;56(1):140-161.
7. James M. Late bone lesions in caisson disease: three cases in submarine personnel. *The Lancet (British edition): Elsevier Ltd;* 1945. 6-8.
8. Mankin HJ. Nontraumatic necrosis of bone (osteonecrosis). *N Engl J Med.* 1992;326(22):1473-1479.
9. Uguen M, Pougnet R, Uguen A, Lodde B, Dewitte JD. Dysbaric osteonecrosis among professional divers: a literature review. *Undersea Hyperb Med.* 2014;41(6):579-587.
10. Bolte H, Koch A, Tetzlaff K, Bettinghausen E, Heller M, Reuter M. Detection of dysbaric osteonecrosis in military divers using magnetic resonance imaging. *Eur Radiol.* 2005;15(2):368-375.
11. Uzun G, Toklu AS, Yildiz S, Sonmez G, et al. Dysbaric osteonecrosis screening in Turkish Navy divers. *Aviat Space Environ Med.* 2008;79(1):44-46.

12. Davidson JK. Dysbaric disorders: aseptic bone necrosis in tunnel workers and divers. *Baillieres Clin Rheumatol*. 1989; 3(1):1-23.
13. McCallum RI, Walder DN. Bone lesions in compressed air workers, with special reference to men who worked on the Clyde Tunnels 1958 to 1963. Report of Decompression Sickness Panel Medical Research Council. *J Bone Joint Surg Br*. 1966;48(2): 207-235.
14. Miyanishi K, Kamo Y, Ihara H, Naka T, et al. Risk factors for dysbaric osteonecrosis. *Rheumatology (Oxford)*. 2006;45(7): 855-858.
15. Jones JP, Jr., Ramirez S, Doty SB. The pathophysiologic role of fat in dysbaric osteonecrosis. *Clin Orthop Relat Res*. 1993(296): 256-264.
16. Lehner CE, Adams WM, Dubielzig RR, Palta M, Lanphier EH. Dysbaric osteonecrosis in divers and caisson workers. An animal model. *Clin Orthop Relat Res*. 1997(344):320-332.
17. Hutter CD. Dysbaric osteonecrosis: a reassessment and hypothesis. *Med Hypotheses*. 2000;54(4):585-590.
18. Arieli R. Nanobubbles form at active hydrophobic spots on the luminal aspect of blood vessels: consequences for decompression illness in diving and possible implications for autoimmune disease-an overview. *Front Physiol*. 2017;8:591.
19. Lanphier EH, Lehner CE, Lemos SE. The role of intramedullary pressure in hyperbaric osteonecrosis engineering and health in compressed air work: Proceedings of the International Conference, Oxford, September 1992. 1994:148-155.
20. Adkins KF. Alizarin red S as an intravital fluorochrome in mineralizing tissues. *Stain Technol*. 1965;40:69-70.
21. Sobakin AS, Wilson MA, Lehner CE, Dueland RT, Gendron-Fitzpatrick AP. Oxygen pre-breathing decreases dysbaric diseases in UW sheep undergoing hyperbaric exposure. *Undersea Hyperb Med*. 2008;35(1):61-67.
22. Bove AA. Diving medicine. *Am J Respir Crit Care Med*: American Thoracic Society; 2014. 1479-1486.
23. Walker III JR, Murphy-Lavoie HM. Diving In water recompression. *StatPearls [Internet]*. 2020.
24. Walder DN. Bone necrosis. In: Jardine FM MR, editor. *Engineering and Health in Compressed Air Work: Proceedings of the International Conference, Oxford, September 1992*: CRC Press, Spon Press; 1994.
25. Gempp E, Louge P, de Maistre S. Predictive factors of dysbaric osteonecrosis following musculoskeletal decompression sickness in recreational scuba divers. *Joint Bone Spine*. 2016; 83(3):357-358.
26. Cialoni D, Pieri M, Balestra C, Marroni A. Dive risk factors, gas bubble formation, and decompression illness in recreational scuba diving: analysis of DAN Europe DSL data base. *Front Psychol*. 2017;8:1587.
27. Arieli R. Fatty diet, active hydrophobic spots, and decompression sickness. *Diving Hyperb Med*. 2018;48(3):197.



Copyright of Undersea & Hyperbaric Medicine is the property of Undersea & Hyperbaric Medical Society and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.