Scalar on Function Regression

Jonathan Willnow, Jakob Juergens, Jonghun Baek

Presentation Day

Introduction

Jona Introductory Example \rightarrow Octane/NIR-spectrum

Jona

Motivation from multivariate regression (multivariate dgp).

Jonghun

- Random Functions (name square integrable functions)
- Motivate continuous stochastic processes (growth curves/electricity consumption/yield curves/stonks)
- Use curves to predict a scalar response (show typical dgp)

Jonghun

- Basis expansions (b-splines and fourier)
- Talk about purposes
- Plots and show bias variance tradeoff

Jakob

- Random function represented as linear combination of basis functions
- Just transform to multiple linear regression setting
- You already know that from the beginning

Estimation via Basis Representation

Assume the following data generating process

$$Y(\omega) = \alpha + \int_0^1 \beta(s) F(\omega)(s) ds + \epsilon(\omega)$$

lacksquare Y and ϵ realize in $\mathbb R$ and F realizes in $\mathbb L^2[0,1]$

Assume that we have a data set containing observations each of which is made up of:

- y_i : a scalar realizations of Y
- $f_i(t)$: a realization of F



Estimation via Basis Representation

Let $\{b_i(t) \mid i=1,\ldots,\infty\}$ be a basis of $\mathbb{L}^2[0,1]$

Then we have the following representation of $\beta(t)$

$$eta(t) = \sum_{j=1}^{\infty} \psi_j b_j(t) = \sum_{j=1}^L \psi_j b_j(t) + \delta(t) pprox \sum_{j=1}^L \psi_j b_j(t)$$

and we can transform the data generating process into:

$$Y(\omega) = \alpha + \int_0^1 \left[\left(\sum_{j=1}^\infty \psi_j b_j(s) \right) F(\omega)(s) \right] ds + \epsilon(\omega)$$
$$= \alpha + \sum_{j=1}^\infty \left[\psi_j \int_0^1 F(\omega)(s) b_j(s) ds \right] + \epsilon(\omega)$$

$$\int_0^1 F(\omega)(s)b_j(s)\mathrm{d}s$$

This is a scalar we can calculate for each observation and for each deterministic basis function $b_i(t)$.

Theory - FPCA

Jakob

- Let's assume you know the theory of PCA (pc from varcov matrix) √
- Introduce mean and covariance functions of random functions ✓
- There is another cool basis \rightarrow Eigenbasis (Karhunen-Loeve Expansion) \checkmark
- Sample Analog! (create a basis from observations and use for basis regression) ✓
- Plot fpcs and approximation of function realization



Spectral Representation of Random Vectors

Let $X(\omega)$ be a random vector realizing in \mathbb{R}^p .

- Let $\mu_X = \mathbb{E}(X)$ and $\Sigma_X = Cov(X)$
- Let $\{\gamma_i \mid i=1,\ldots,p\}$ be the orthonormal **Eigenvectors** of Σ_X
- Let $\{\lambda_i \mid i=1,\ldots,p\}$ be the corresponding **Eigenvalues** of Σ_X

Then X can also be represented as

$$X(\omega) = \mu_{x} + \sum_{i=1}^{p} \xi_{i}(\omega)\gamma_{i}$$

where the $\xi_i(\omega)$ have the following properties

$$\mathbb{E}[\xi_i(\omega)] = 0$$

2
$$Var(\xi_i(\omega)) = \lambda_i$$

3
$$Cov(\xi_i(\omega), \xi_j(\omega)) = 0$$
 for $i \neq i$



Karhunen-Loéve Expansion

Mean Function:

$$\mu(t) = \mathbb{E}\left[F(\omega)(t)\right]$$

Autocovariance Function:

$$c(t,s) = \mathbb{E}\big[\left(F(\omega)(t) - \mu(t)\right)\left(F(\omega)(s) - \mu(s)\right)\big]$$

The **Eigenvalues** and **Eigenfunctions**: $\{(\lambda_i, \nu_i) \mid i \in \mathcal{I}\}$ are solutions of the following equation:

$$\int_0^1 c(t,s)\nu(s)\mathrm{d}s = \lambda\nu(t)$$



Karhunen-Loéve Expansion

A random function F can be expressed in terms of its mean function and its Eigenfunctions:

$$F(\omega)(t) = \mu(t) + \sum_{j=1}^{\infty} \xi_j(\omega)\nu_j(t)$$

Where the ξ_j are scalar-valued random variables with the following properties.

$$2 Var(\xi_i(\omega)) = \lambda_i$$

3
$$Cov(\xi_i(\omega), \xi_j(\omega)) = 0$$
 for $i \neq j$

This representation is called the **Karhunen-Loéve Expansion** of the random function F and the Eigenfunctions can serve as a basis to represent the function.

Principal Component Analysis

A related concept is **Principal Component Analysis** (PCA).

Σ_X unknown o sample analogues

- Let $\mathbf{X} \in \mathbb{R}^{n \times p}$ contain the standardized regressors
- Let $\hat{\Sigma}_X = \frac{\mathbf{X}'\mathbf{X}}{n}$
- Let $\{\hat{\gamma}_i \mid i=1,\ldots,p\}$ be the orthonormal **Eigenvectors** of $\hat{\Sigma}_X$
- Let $\{\hat{\lambda}_i | i=1,\ldots,p\}$ be the corresponding **Eigenvalues** of $\hat{\Sigma}_X$

Then $Z_i(\omega) = \hat{\gamma}_i' X(\omega)$ is called the i'th principal component and

3
$$Cov(Z_i(\omega), Z_j(\omega)) = 0$$
 for $i \neq j$



Functional Principal Component Analysis

This idea can be extended to functional regressors in the form of **Functional Principal Component Analysis** (FPCA).

Empirical Mean Function:

$$\hat{\mu}(t) = \frac{1}{n} \sum_{j=1}^{n} f_j(t)$$

Empirical Autocovariance Function:

$$\hat{c}(t,s) = rac{1}{n} \sum_{j=1}^{n} \left(f_j(t) - \hat{\mu}(t) \right) \left(f_j(s) - \hat{\mu}(s) \right)$$

Functional Principal Component Analysis

The **Eigenvalues** and **Eigenfunctions**: $\{(\hat{\lambda}_i, \hat{\nu}_i) \mid i \in \mathcal{I}\}$ are solutions of the following equation:

$$\int_0^1 \hat{c}(t,s)\hat{\nu}(s)\mathrm{d}s = \hat{\lambda}\hat{\nu}(t)$$

The $\{\hat{\nu}_i(s) \mid i \in \mathcal{I}\}$ are called **Functional Principal Components** and can serve as a basis for representing the original curves.

The corresponding scores $\hat{\xi}_i$ can be derived as

$$\hat{\xi}_j(\omega) = \int_0^1 (F(\omega)(s) - \hat{\mu}(s))\hat{\nu}_j(s)\mathrm{d}s$$



Simulation Setup & Application

Jona

- Compare b-spline / fourier regression chosen via criterion (cv/aic/...)
- Similar for fpca
- generate new curves from observed curves motivated by Karhunen-Loeve expansion
- Compare optimal variants with test and training sets
- Connect to Application

Summary

Jona

Just summarize what we have done...

further reading

Put footnotes here!

