

Causality

Causal BNs and Structural Causal Models

Maciej Liśkiewicz

University of Lübeck

December, 2022

Causal Models

Agenda

- *Causal* Bayesian Networks
- Modelling of controlled change in the model / controlled experiments
- Intervention
- *do*-operator
- Intervention joint distribution
- Causal effects
- Structural Causal Models / Functional Causal Models

Causal Bayesian Networks

- So far, we have considered a BN as a carrier of conditional independence relationships
- However, such an “associational knowledge representation” is not sufficient when we want to make decisions under uncertainty
- Particularly, the interpretation of DAGs as carriers of CI assumptions does not necessarily imply causation

Causal Bayesian Networks

- So far, we have considered a BN as a carrier of conditional independence relationships
- However, such an “associational knowledge representation” is not sufficient when we want to make decisions under uncertainty
- Particularly, the interpretation of DAGs as carriers of CI assumptions does not necessarily imply causation
- For example, in our “recommendation letter” BN, the (in)dependence between I – student’s intelligence and S – the student’s SAT score, can be encoded as

$$I \rightarrow S \quad \text{and} \quad I \leftarrow S$$

Causal Bayesian Networks

- So far, we have considered a BN as a carrier of conditional independence relationships
- However, such an “associational knowledge representation” is not sufficient when we want to make decisions under uncertainty
- Particularly, the interpretation of DAGs as carriers of CI assumptions does not necessarily imply causation
- For example, in our “recommendation letter” BN, the (in)dependence between I – student’s intelligence and S – the student’s SAT score, can be encoded as

$$I \rightarrow S \quad \text{and} \quad I \leftarrow S$$

- Our goal: causal interpretation of DAGs
 - ▶ the ubiquity of graphical models in statistical and AI applications stems primarily from their causal interpretation

Causal Bayesian Networks


- The advantages of building models around **causal** rather than **associational** information
 1. The judgments required in the construction of a DAG are more *meaningful* and therefore more *reliable*
 2. The ability to represent and respond to external or spontaneous *changes*
- **Example: “slippery pavement” (Pearl (2009, Ch. 1))**
- All these imply that causal models (assuming they are valid) are much more informative than probability models

Causal Networks and Interventions

Primacy of causal over associational knowledge...

Example: “Slippery Pavement” (Pearl (2009, ch. 1))

- X_1 : the season of the year (spring, summer, fall, winter)
- X_2 : whether rain falls
- X_3 : whether the sprinkler is on
- X_4 : whether the pavement would get wet
- X_5 : whether the pavement would be slippery




- ... meaningful judgments in the construction of a DAG

Causal Networks and Interventions

Primacy of causal over associational knowledge...

Example: "Slippery Pavement" (Pearl (2009, ch. 1))

- X_1 : the season of the year (spring, summer, fall, winter)
- X_2 : whether rain falls
- X_3 : whether the sprinkler is on
- X_4 : whether the pavement would get wet
- X_5 : whether the pavement would be slippery




- ... meaningful judgments in the construction of a DAG
- **Exercise:** attempt to construct a DAG representation for the associations in DAG above along the ordering $(X_5, X_1, X_3, X_2, X_4)$

Causal Networks and Interventions

Primacy of causal over associational knowledge...


Example “Slippery Pavement” (Pearl (2009, ch. 1))



- ... the ability to represent and respond to external or spontaneous changes
- Each parent-child relationship represents a *stable* and *autonomous* physical mechanism
- It is conceivable to change one parent-child relationship without changing the others
- This property allows the following
 - ▶ A joint distribution tells us how probabilities would change with subsequent observations (as in case of BNs)
 - ▶ A causal model **also** tells us how these probabilities would change as a **result of external interventions**


Causal Networks and Interventions

- Direct experimentation



Causal Networks and Interventions

- A causal model tells how the probabilities would change as a **result of external interventions**



- $P(x_1, x_2, x_3, x_4, x_5) = P(x_1) P(x_2 | x_1) P(x_3 | x_1) P(x_4 | x_2, x_3) P(x_5 | x_4)$
- In a causal model there is a deep connection between **modularity** and **interventions**
- Instead of specifying a **new probability function** P for each of the many possible interventions, we specify merely the immediate change implied by the intervention
- As a result we get that the **effect of an intervention** can be predicted by modifying the corresponding factors in the decomposition $P(x_1, x_2, x_3, x_4, \dots)$

Causal Networks and Interventions

- A causal model tells how the probabilities would change as a **result of external interventions**



- Our “pre-intervention” probability function

$$P(x_1, x_2, x_3, x_4, x_5) = P(x_1) P(x_2 | x_1) P(x_3 | x_1) P(x_4 | x_2, x_3) P(x_5 | x_4)$$

- Example: to represent the (external) action “turn the sprinkler On” we

- ▶ remove $X_1 \rightarrow X_3$ from the model and
- ▶ assign X_3 to the value “ON”

- We get the post-intervention probability function

$$P_{X_3=ON}(x_1, x_2, x_4, x_5) = P(x_1) P(x_2 | x_1) P(x_4 | x_2, X_3 = ON) P(x_5 | x_4)$$

Causal Networks and Interventions

- A causal model tells how the probabilities would change as a **result of external interventions**



- The post-intervention probability function

$$P_{X_3=ON}(x_1, x_2, x_4, x_5) = P(x_1) P(x_2 | x_1) P(x_4 | x_2, X_3 = ON) P(x_5 | x_4)$$

- The **deletion** of $P(x_3 | x_1)$ from the factorization means, whatever relationship existed between X_1 and X_3 **prior to the action**, that relationship is no longer valid while we perform the action
- We denote the action using the **do-operator** introduced by Pearl:

$$do(X_3 = ON)$$

Causal Networks and Interventions

- A causal model tells how the probabilities would change as a **result of external interventions**



- We will use both notations (meaning the same)

$$P_{X_3=\text{ON}}(x_1, x_2, x_4, x_5) \quad \text{and} \quad P(x_1, x_2, x_4, x_5 \mid \text{do}(X_3 = \text{ON}))$$

but we will prefer the last one


- According to the previously accepted notation this leads also to the following

$$P_{x_3}(x_1, x_2, x_4, x_5) \quad \text{and} \quad P(x_1, x_2, x_4, x_5 \mid \text{do}(x_3))$$

where by x_3 we mean a specific value of X_3 (in our case $x_3 = \text{ON}$)

Causal Networks and Interventions

- A causal model tells how the probabilities would change as a **result of external interventions**



- Note the difference between $P(x_1, x_2, x_4, x_5 \mid X_3 = \text{ON})$ and $P(x_1, x_2, x_4, x_5 \mid do(X_3 = \text{ON}))$
- $P(x_1, x_2, x_4, x_5 \mid X_3 = \text{ON})$ is obtained by conditioning in the (left) model
- As a consequence: after **observing** that $X_3 = \text{ON}$ (evidence), we wish to infer
 - that the season is dry
 - that it probably did not rain
 - etc.
- After the **action** “turning the sprinkler On” (right) we infer quite different knowledge

Causal Networks and Interventions

- Let \mathbf{v} be a sequence of values of variables V_1, \dots, V_n
- Let $\mathbf{X} \subseteq \mathbf{V} = \{V_1, \dots, V_n\}$ and let \mathbf{x} be a sequence of values of variables \mathbf{X}
- We say that $\mathbf{v} = (v_1, \dots, v_n)$ is consistent with $\mathbf{x} = (v'_{i_1}, \dots, v'_{i_k})$ for all i_j if $v_{i_j} = v'_{i_j}$

Causal Networks and Interventions

- Let \mathbf{v} be a sequence of values of variables V_1, \dots, V_n
- Let $\mathbf{X} \subseteq \mathbf{V} = \{V_1, \dots, V_n\}$ and let \mathbf{x} be a sequence of values of variables \mathbf{X}
- We say that $\mathbf{v} = (v_1, \dots, v_n)$ is consistent with $\mathbf{x} = (v'_{i_1}, \dots, v'_{i_k})$ for all i_j if $v_{i_j} = v'_{i_j}$

Interventional Distribution, Causal Effect

- Assume $G = (\mathbf{V} = \{V_1, \dots, V_n\}, \mathbf{E})$ is a causal Bayesian network
- The **interventional distribution** $P(\mathbf{v} \mid do(\mathbf{x}))$ resulting from any intervention $do(\mathbf{X} = \mathbf{x})$ is defined as a truncated factorization:

$$P(\mathbf{v} \mid do(\mathbf{x})) = \prod_{\{i: V_i \notin \mathbf{X}\}} P(v_i \mid pa_i) \quad \text{for all } \mathbf{v} \text{ consistent with } \mathbf{x}$$

Causal Networks and Interventions

- Let \mathbf{v} be a sequence of values of variables V_1, \dots, V_n
- Let $\mathbf{X} \subseteq \mathbf{V} = \{V_1, \dots, V_n\}$ and let \mathbf{x} be a sequence of values of variables \mathbf{X}
- We say that $\mathbf{v} = (v_1, \dots, v_n)$ is consistent with $\mathbf{x} = (v'_{i_1}, \dots, v'_{i_k})$ for all i_j if $v_{i_j} = v'_{i_j}$

Interventional Distribution, Causal Effect

- Assume $G = (\mathbf{V} = \{V_1, \dots, V_n\}, \mathbf{E})$ is a causal Bayesian network
- The **interventional distribution** $P(\mathbf{v} \mid do(\mathbf{x}))$ resulting from any intervention $do(\mathbf{X} = \mathbf{x})$ is defined as a truncated factorization:

$$P(\mathbf{v} \mid do(\mathbf{x})) = \prod_{\{i: V_i \notin \mathbf{X}\}} P(v_i \mid pa_i) \quad \text{for all } \mathbf{v} \text{ consistent with } \mathbf{x}$$

- The **causal effect** of \mathbf{X} on outcome variables \mathbf{Y} , denoted as

$$P(\mathbf{y} \mid do(\mathbf{x}))$$

is defined as the probability distribution of variables \mathbf{Y} after the intervention

Causal Networks and Interventions

- Let \mathbf{v} be a sequence of values of variables V_1, \dots, V_n
- Let $\mathbf{X} \subseteq \mathbf{V} = \{V_1, \dots, V_n\}$ and let \mathbf{x} be a sequence of values of variables \mathbf{X}
- We say that $\mathbf{v} = (v_1, \dots, v_n)$ is consistent with $\mathbf{x} = (v'_{i_1}, \dots, v'_{i_k})$ for all i_j if $v_{i_j} = v'_{i_j}$

Interventional Distribution, Causal Effect

- Assume $G = (\mathbf{V} = \{V_1, \dots, V_n\}, \mathbf{E})$ is a causal Bayesian network
- The **interventional distribution** $P(\mathbf{v} \mid do(\mathbf{x}))$ resulting from any intervention $do(\mathbf{X} = \mathbf{x})$ is defined as a truncated factorization:

$$P(\mathbf{v} \mid do(\mathbf{x})) = \prod_{\{i: V_i \notin \mathbf{X}\}} P(v_i \mid pa_i) \quad \text{for all } \mathbf{v} \text{ consistent with } \mathbf{x}$$

- The **causal effect** of \mathbf{X} on outcome variables \mathbf{Y} , denoted as

$$P(\mathbf{y} \mid do(\mathbf{x}))$$

is defined as the probability distribution of variables \mathbf{Y} after the intervention

- Notice that if \mathbf{v} is not consistent with \mathbf{x} , we let $P(\mathbf{v} \mid do(\mathbf{x})) = 0$

Causal Networks and Interventions

Example: Observation vs. Intervention

- Assume $\mathbf{V} = \{R, W\}$
- R : it is raining ($R = 1$) or not ($R = 0$);
- W : the street is wet ($W = 1$) or not ($W = 0$)

Causal Networks and Interventions

Example: Observation vs. Intervention


- Assume $\mathbf{V} = \{R, W\}$
- R : it is raining ($R = 1$) or not ($R = 0$);
- W : the street is wet ($W = 1$) or not ($W = 0$)
- **Observation vs. Intervention** R : it is raining ($R = 1$) or not ($R = 0$);
 W : the street is wet ($W = 1$) or not ($W = 0$)

$$R \longrightarrow W$$

Causal Networks and Interventions

Example: Observation vs. Intervention

- Assume $\mathbf{V} = \{R, W\}$
- R : it is raining ($R = 1$) or not ($R = 0$);
- W : the street is wet ($W = 1$) or not ($W = 0$)



- ▶ Suppose: $P(R = 1) = 0.01$, and $P(W = 1 | R = 1) = 1$, $P(W = 1 | R = 0) = 0.001$.
- ▶ Let's suppose we **observe** that the street is wet. Using Bayes Theorem, we get:

$$P(R = 1 | W = 1) = 0.91$$

Causal Networks and Interventions

Example: Observation vs. Intervention

- Assume $\mathbf{V} = \{R, W\}$
- R : it is raining ($R = 1$) or not ($R = 0$);
- W : the street is wet ($W = 1$) or not ($W = 0$)

$$R \rightarrow W$$



- ▶ Suppose: $P(R = 1) = 0.01$, and $P(W = 1 | R = 1) = 1$, $P(W = 1 | R = 0) = 0.001$.
- ▶ Let's suppose we **observe** that the street is wet. Using Bayes Theorem, we get:

$$P(R = 1 | W = 1) = 0.91$$

- ▶ If we **intervene** to make the street wet, we get:

$$P(R = 1 | do(W = 1)) = 0.01$$

Causal Networks and Interventions

Example: Observation vs. Intervention

- Assume $\mathbf{V} = \{R, W\}$
- R : it is raining ($R = 1$) or not ($R = 0$);
- W : the street is wet ($W = 1$) or not ($W = 0$)

$$R \rightarrow W$$



- ▶ Suppose: $P(R = 1) = 0.01$, and $P(W = 1 | R = 1) = 1$, $P(W = 1 | R = 0) = 0.001$.
- ▶ Let's suppose we **observe** that the street is wet. Using Bayes Theorem, we get:

$$P(R = 1 | W = 1) = 0.91$$

- ▶ If we **intervene** to make the street wet, we get:

$$P(R = 1 | do(W = 1)) = 0.01$$

- ▶ We get, also that $P(R = 1 | do(W = 0)) = 0.01$
- ▶ Conclusion: W has no causal effect on $R = 1$


The Identifiability Problem

Example: Smoking and the Genotype Theory (Pearl (2009))

- Does smoking (S) cause lung cancer (C)?
- Assume that the variables are binary, taking on true (1) or false (0) values
- No relevant factors are assumed
- Moreover, assume the following (hypothetical) data set from a study on the relations among cancer and cigarette smoking

	Group Type	% of Population	% of Cancer cases
$S = 0$	Nonsmokers	50	9.75
$S = 1$	Smokers	50	85.25

- Task: compute the causal effects $P(C = 1 \mid do(s))$ from data in this model



$$P(c \mid do(s)) = P(c \mid s)$$

- $P(C = 1 \mid do(S = 0)) = 0.0975$
- $P(C = 1 \mid do(S = 1)) = 0.8525$

Causal Networks and Interventions

The Identifiability Problem

Identifiability of Causal Effects

The identification of the total causal effect (in general, non-parametric models), called the **identification problem**, is defined as follows:

- for a given DAG $G = (\mathbf{V}, \mathbf{E})$, a set $\mathbf{R} \subseteq \mathbf{V}$ representing observed variables, and two disjoint subsets $\mathbf{X}, \mathbf{Y} \subseteq \mathbf{V}$,
- express the causal total effect $P(\mathbf{y} \mid do(\mathbf{x}))$ using only pre-intervention (i.e. do-operator free) probabilities involving variables in \mathbf{R} , or output that this is not possible.

When such a formula exists, we say that the causal effect of \mathbf{X} on \mathbf{Y} in G is **identifiable**.


The Identifiability Problem

Example: Smoking and the Genotype Theory (Pearl (2009))

- Does smoking (S) cause lung cancer (C)?
- Assume, to forestall antismoking legislation, the tobacco industry has argued that the observed correlation between smoking and lung cancer could be explained by some sort of **carcinogenic genotype** that involves inborn craving for nicotine
- Thus, consider in our model the relevant factor: Genotype (G)
- Unfortunately, the feature is **not measurable** (called also **unobserved**)
- Can the causal effects $P(C = 1 \mid do(s))$ be estimated from data in this model?

	Group Type	% of Population	% of Cancer cases
$S = 0$	Nonsmokers	50	9.75
$S = 1$	Smokers	50	85.25

Genotype (unobserved)



$$P(c \mid do(s)) = \text{not identifiable !}$$

The Identifiability Problem

Example: Smoking and the Genotype Theory (Pearl (2009))

- Does smoking (S) cause lung cancer (C)?
- Consider now the amount of tar deposited in a person's lungs
- The relevant factors in our model are now: Genotype (G , unobserved), Tar in the lungs (T binary, taking on true (1) or false (0))
- How to compute the causal effects $P(C = 1 \mid do(s))$ from data:

		Group Type	$P(s, t)$ % of Population	$P(C = 1 \mid s, t)$ % of Cancer cases
$S = 0$	$T = 0$	Nonsmokers, No tar	47.5	10
$S = 1$	$T = 0$	Smokers, No tar	2.5	90
$S = 0$	$T = 1$	Nonsmokers, Tar	2.5	5
$S = 1$	$T = 1$	Smokers, Tar	47.5	85


The Identifiability Problem

Example: Smoking and the Genotype Theory (Pearl (2009))

- Task: compute the causal effects $P(C = 1 \mid do(s))$ from data:

		Group Type	$P(s, t)$ % of Population	$P(C = 1 \mid s, t)$ % of Cancer cases
$S = 0$	$T = 0$	Nonsmokers, No tar	47.5	10
$S = 1$	$T = 0$	Smokers, No tar	2.5	90
$S = 0$	$T = 1$	Nonsmokers, Tar	2.5	5
$S = 1$	$T = 1$	Smokers, Tar	47.5	85

Genotype (unobserved)




The Identifiability Problem

Example: Smoking and the Genotype Theory (Pearl (2009))

- Task: compute the causal effects $P(C = 1 \mid do(s))$ from data:

		Group Type	$P(s, t)$ % of Population	$P(C = 1 \mid s, t)$ % of Cancer cases
$S = 0$	$T = 0$	Nonsmokers, No tar	47.5	10
$S = 1$	$T = 0$	Smokers, No tar	2.5	90
$S = 0$	$T = 1$	Nonsmokers, Tar	2.5	5
$S = 1$	$T = 1$	Smokers, Tar	47.5	85

Genotype (unobserved)



Smoking Tar Cancer

In our course we will show that the following formula can be used:


$$P(c \mid do(s)) = \sum_t P(t \mid s) \sum_{s'} P(c \mid s', t) P(s')$$

- $P(C = 1 \mid do(S = 0)) = .95(.10 \times .50 + .90 \times .50) + .05(.05 \times .50 + .85 \times .50)$
 $= .95 \times .50 + .05 \times .45 = .4975$
- $P(C = 1 \mid do(S = 1)) = .05(.10 \times .50 + .90 \times .50) + .95(.05 \times .50 + .85 \times .50)$
 $= .05 \times .50 + .95 \times .45 = .4525$

Structural Causal Models

- Next we define more general causal models:
- Structural Causal Models (SCM) called also Functional Causal Models
- The first component of an SCM is a collection of assignments, e.g.
 - ▶ $x = f_X(u_X)$
 - ▶ $y = f_Y(x, u_Y)$

that induces a DAG



- The second component is the probability function $P(U_X, U_Y)$

Structural Causal Models

Definition

Structural Causal Model

- A structural causal model (SCM) $M = (S, P)$ consists of two sets of variables $\mathbf{V} = \{X_1, \dots, X_n\}$ and $\mathbf{U} = \{U_1, \dots, U_n\}$ and

- ▶ a collection S of n (structural) assignments

$$X_i = f_i(Pa_i, U_i)$$

where $Pa_i \subseteq \mathbf{V}$ are called parents of X_i and

- ▶ a joint distribution P over \mathbf{U}
- The variables in \mathbf{U} are called **exogenous** variables, meaning that they are external to the model
- The variables in \mathbf{V} are **endogenous**
- The graph G associated with an SCM is obtained by creating one vertex for each X_i and drawing directed edges from each parent in Pa_i to X_i
- We assume the resulting G is a DAG

Structural Causal Models

Basic Properties

- We sometimes call the exogenous variables in \mathbf{U} the **noise** or **error** variables
- The elements of Pa_i are called not only parents but also
 - ▶ **direct causes** of X_i , and
 - ▶ we call X_i a **direct effect** of each of its direct causes

Structural Causal Models

Basic Properties

- We sometimes call the exogenous variables in \mathbf{U} the **noise** or **error** variables
- The elements of Pa_i are called not only parents but also
 - ▶ **direct causes** of X_i , and
 - ▶ we call X_i a **direct effect** of each of its direct causes
- P specifies the distribution of all exogenous variables
- In the case when the causal diagram is **acyclic** (as assumed in our definition), the corresponding model is called **semi-Markovian**

Structural Causal Models

Basic Properties

- We sometimes call the exogenous variables in \mathbf{U} the **noise** or **error** variables
- The elements of Pa_i are called not only parents but also
 - ▶ **direct causes** of X_i , and
 - ▶ we call X_i a **direct effect** of each of its direct causes
- P specifies the distribution of all exogenous variables
- In the case when the causal diagram is **acyclic** (as assumed in our definition), the corresponding model is called **semi-Markovian**
- In such models the values of the X_i variables will be uniquely determined by those of the \mathbf{U} variables
- Under such conditions, the joint distribution $P(x_1, \dots, x_n)$ is determined uniquely by the distribution $P(\mathbf{u})$ of the error variables

Structural Causal Models

- If, in addition to acyclicity, the error terms are jointly independent, the model is called **Markovian**
- Example of an Markovian and non-Markovian SCM



- Sometimes, in DAGs associated with Markovian SCMs we will not show explicitly the exogene variables U_1, U_2, \dots
- By convention, this implies that they are assumed to be mutually independent.
- Additionally, we have the following properties
 - ▶ Every endogenous variable in a model is a descendant of at least one exogenous variable
 - ▶ Exogenous variables cannot be descendants of any other variables, and in particular, cannot be a descendant of an endogenous variable

Structural Causal Models

Basic Properties

Theorem (Causal Markov Condition)

Assume $M = (S, P)$ is a Markovian SCM. Then

- M defines a unique distribution over the variables $\mathbf{V} = \{X_1, \dots, X_n\}$, such that $X_i = f_i(Pa_i, U_i)$ in distribution for $i = 1, \dots, n$
- We refer to it as the associated with M distribution and write $P_{\mathbf{V}}$
- The distribution $P_{\mathbf{V}}(x_1, \dots, x_n)$ satisfies the parental Markov (local) condition relative the causal diagram G ; that is, for each variable X_i :

$$(X_i \perp\!\!\!\perp \mathbf{V} \setminus (De(X_i) \cup Pa(X_i)) \mid Pa(X_i))_{P_{\mathbf{V}}}$$

Structural Causal Models

Basic Properties

Theorem (Causal Markov Condition)

Assume $M = (S, P)$ is a Markovian SCM. Then

- M defines a unique distribution over the variables $\mathbf{V} = \{X_1, \dots, X_n\}$, such that $X_i = f_i(Pa_i, U_i)$ in distribution for $i = 1, \dots, n$
- We refer to it as the associated with M distribution and write $P_{\mathbf{V}}$
- The distribution $P_{\mathbf{V}}(x_1, \dots, x_n)$ satisfies the parental Markov (local) condition relative the causal diagram G ; that is, for each variable X_i :

$$(X_i \perp\!\!\!\perp \mathbf{V} \setminus (De(X_i) \cup Pa(X_i)) \mid Pa(X_i))_{P_{\mathbf{V}}}$$



Exercise Compute $P(x, y)$ in M , with $x = u_X$, $y = x \cdot u_Y + (1 - x)(1 - u_Y)$ where U_X and U_Y are two independent binary variables with $P(u_X = 1) = P(u_Y = 1) = 1/2$ (e.g., random coins)

Structural Causal Models

Interventions and Causal Effects in SCM

Definition (Intervention in SCMs, Causal Effect)

- Consider an SCM $M = (S, P)$ over $\mathbf{V} = \{X_1, \dots, X_n\}$ and \mathbf{U} , with S defined for $i = 1, \dots, n$ as

$$X_i = f_i(Pa_i, U_i)$$

- To model the **intervention**

$$do(X_i = x_i), \quad \text{or in general} \quad do(X_{i_1} = x_{i_1}, \dots, X_{i_k} = x_{i_k})$$

we replace one (or several) of the structural assignments f for the variables to intervene, to obtain a new SCM $M' = (S', P)$, with

$$f'_i(Pa_i, U_i) := x_i$$

and in general case

$$f'_{i_1}(Pa_{i_1}, U_{i_1}) := x_{i_1}, \dots, f'_{i_k}(Pa_{i_k}, U_{i_k}) := x_{i_k}$$

- We define the **causal effect** of $\mathbf{X} \subseteq \mathbf{V}$ on outcome variables $\mathbf{Y} \subseteq \mathbf{V}$, denoted as

$$P_{\mathbf{V}}(\mathbf{y} \mid do(\mathbf{x}))$$


as the probability distribution of variables \mathbf{Y} after the intervention

Structural Causal Models

Interventions and Causal Effects in SCM

Example (Pattern Recognition (Peters, Janzing, Schölkopf (2017), Sec. 1.4.1)

Structural causal models of handwritten digit data sets




- Model M_1 : a human is provided with **class labels y** produces a corresponding **handwritten digit image x** ; We model the process as a suitable function (or mechanism) f_X of the class label Y and some independent exogenous (noise) variable U_X
- We can then compute $P_{X,Y}$ from P_{U_X, U_Y} , and f_X
- There are two possible interventions in M_1 , which lead to intervention distributions

Structural Causal Models

Interventions and Causal Effects in SCM

Example (Pattern Recognition (Peters, Janzing, Schölkopf (2017), Sec. 1.4.1)

Structural causal models of handwritten digit data sets




- Model M_2 : the human decides which class to write (Z) and produces both **images (X)** and **class labels (Y)**; Both the image X and the recorded class label Y are functions of the writer's intention (Z)
- We assume Z , U_X , and U_Y are independent

Structural Causal Models

Interventions and Causal Effects in SCM

Example (Pattern Recognition (Peters, Janzing, Schölkopf (2017), Sec. 1.4.1)

Structural causal models of handwritten digit data sets



- If the functions f_X in M_1 and f_X, f_Y in M_2 and noise terms are chosen suitably, we can ensure that M_1 and M_2 entail the same observational distributions $P_{X,Y}$
- However they are interventionally different

Structural Causal Models vs Causal BN

Linear SCMs (SEMs)

- In its general form, an SCM $M = (S, P)$ consists of a set S of equations of the form

$$X_j = f_j(Pa_j, U_j) \quad j = 1, \dots, n$$

- The equations are nonlinear, nonparametric generalization of the (recursive) **linear structural equation models** (SEMs)

$$X_j = \sum_{i < j} c_{ji} X_i + U_j, \quad j = 1, \dots, n.$$

- Parameters c_{ji} are called a **path coefficients** and they describe **direct causal effects** of X_i on X_j
- Values u_j represent **error terms** and there is assumed they have **normal distribution**

Structural Causal Models vs Causal BN

Linear SCMs (SEMs)

- In its general form, an SCM $M = (S, P)$ consists of a set S of equations of the form

$$X_j = f_j(Pa_j, U_j) \quad j = 1, \dots, n$$

- The equations are nonlinear, nonparametric generalization of the (recursive) **linear structural equation models** (SEMs)

$$X_j = \sum_{i < j} c_{ji} X_i + U_j, \quad j = 1, \dots, n.$$

- Parameters c_{ji} are called a **path coefficients** and they describe **direct causal effects** of X_i on X_j
- Values u_j represent **error terms** and there is assumed they have **normal distribution**
- Example**

$$Z_1 = U_1$$

$$Z_2 = U_2$$


$$X_1 = a_1 Z_1 + a_2 Z_2 + U_3$$

$$X_2 = b_1 Z_1 + b_2 Z_2 + U_4$$

$$Y = c_1 X_1 + c_2 X_2 + U_5$$

$$\text{Cov}(U_3, U_5) = \beta_1 \neq 0$$

$$\text{Cov}(U_4, U_5) = \beta_2 \neq 0$$



Structural Causal Models vs Causal BN

Linear SCMs (SEMs)

- In its general form, an SCM $M = (S, P)$ consists of a set S of equations of the form

$$X_j = f_j(Pa_j, U_j) \quad j = 1, \dots, n$$

- The equations are nonlinear, nonparametric generalization of the (recursive) **linear structural equation models** (SEMs)

$$X_j = \sum_{i < j} c_{ji} X_i + U_j, \quad j = 1, \dots, n.$$

- Parameters c_{ji} are called a **path coefficients** and they describe **direct causal effects** of X_i on X_j
- Values u_j represent **error terms** and there is assumed they have **normal distribution**
- Example**

$$Z_1 = U_1$$

$$Z_2 = U_2$$


$$X_1 = a_1 Z_1 + a_2 Z_2 + U_3$$

$$X_2 = b_1 Z_1 + b_2 Z_2 + U_4$$

$$Y = c_1 X_1 + c_2 X_2 + U_5$$

$$\text{Cov}(U_3, U_5) = \beta_1 \neq 0$$

$$\text{Cov}(U_4, U_5) = \beta_2 \neq 0$$



- SEM is a powerful multivariate analysis technique that is widely used by many applied researchers in the social and behavioral sciences

Literature

- J. Pearl (2009), Ch.1
- J. Pearl, M. Glymour, and N.P. Jewell (2016), Ch. 1,2