FACULTY OF FUNDAMENTAL PROBLEMS OF TECHNOLOGY WROCŁAW UNIVERSITY OF SCIENCE AND TECHNOLOGY

EIGENMODES IN NEARLY INTEGRABLE QUANTUM CHAINS

Jakub Pawłowski

Index number: 250193

Master thesis under supervision of prof. dr hab. Marcin Mierzejewski



This page is intentionally left blank. $\,$

I would like to express my sincere gratitude to prof. dr hab. Marcin Mierzejewski for being my

This page is intentionally left blank.

Abstract

Apparent incompatibility of classical irreversible thermodynamics with

 ${\bf Keywords:}\ integrals\ of\ motion,\ ETH,\ integrability\ breaking,\ XXZ\ model$

This page is intentionally left blank.

Contents

1	Introduction	1
2	Lanczos algorithm	3
	2.1 Lanczos method for ground state calculation	3
3	Summary	5
Bi	ibliography	7
A	Appendix	7



1 Introduction

AAA



Lanczos algorithm

The main purpose of this thesis is to develop and test a set of numerical tools based on the Lanczos algorithm, which is an iterative method for finding extremal eigenvalues of a large matrix. In its simplest form, it is useful for efficient calculation of a ground state energy and eigenvector, and subsequently for determining the ground state properties of a system of choice. However, in this work we are mainly interested in infinite temperature properties, which requires sampling of the whole spectrum of Hamiltonian in question.

2.1 Lanczos method for ground state calculation



3 Summary

AAA



Appendix

