

SC223 - Linear Algebra

Aditya Tatu

Lecture 6



August 23, 2022

Gaussian Elimination/Row Reduction

- Let us use the following example:

$$2x_2 + 5x_3 + 4x_4 + 2x_5 = 2$$

$$x_1 - x_2 + 2x_3 + 3x_4 - x_5 = 1$$

$$2x_1 + x_2 + 4x_4 + 2x_5 = -1$$

$$3x_1 + x_2 + 3x_3 + 2x_4 - 2x_5 = 3$$

Gaussian Elimination/Row Reduction

- Let us use the following example:

$$2x_2 + 5x_3 + 4x_4 + 2x_5 = 2$$

$$x_1 - x_2 + 2x_3 + 3x_4 - x_5 = 1$$

$$2x_1 + x_2 + 4x_4 + 2x_5 = -1$$

$$3x_1 + x_2 + 3x_3 + 2x_4 - 2x_5 = 3$$

- When are linear equations easy to solve?

Gaussian Elimination/Row Reduction

- Let us use the following example:

$$2x_2 + 5x_3 + 4x_4 + 2x_5 = 2$$

$$x_1 - x_2 + 2x_3 + 3x_4 - x_5 = 1$$

$$2x_1 + x_2 + 4x_4 + 2x_5 = -1$$

$$3x_1 + x_2 + 3x_3 + 2x_4 - 2x_5 = 3$$

- When are linear equations easy to solve? As few variables (ideally 1) as possible.

Gaussian Elimination/Row Reduction

- Let us use the following example:

$$2x_2 + 5x_3 + 4x_4 + 2x_5 = 2$$

$$x_1 - x_2 + 2x_3 + 3x_4 - x_5 = 1$$

$$2x_1 + x_2 + 4x_4 + 2x_5 = -1$$

$$3x_1 + x_2 + 3x_3 + 2x_4 - 2x_5 = 3$$

- When are linear equations easy to solve? As few variables (ideally 1) as possible.
- Representation as a matrix

$$\left[\begin{array}{ccccc|c} 0 & 2 & 5 & 4 & 2 & 2 \\ 1 & -1 & 2 & 3 & -1 & 1 \\ 2 & 1 & 0 & 4 & 2 & -1 \\ 3 & 1 & 3 & 2 & -2 & 3 \end{array} \right]$$

- This is called the **Augmented matrix**.

- What can we do to reduce the number of variables, while still preserving the solutions?

- What can we do to reduce the number of variables, while still preserving the solutions?
- Combining two (or more) equations to get a simpler equation is equivalent to replacing a row with a linear combination of two (or more) rows of the AM!

- What can we do to reduce the number of variables, while still preserving the solutions?
- Combining two (or more) equations to get a simpler equation is equivalent to replacing a row with a linear combination of two (or more) rows of the AM!
- **Elementary row operations:** Exchanging two rows, adding a multiple of one row to another, and multiplying a constant to all entries of a row. All of them preserve solutions to linear equations.

- What can we do to reduce the number of variables, while still preserving the solutions?
- Combining two (or more) equations to get a simpler equation is equivalent to replacing a row with a linear combination of two (or more) rows of the AM!
- **Elementary row operations:** Exchanging two rows, adding a multiple of one row to another, and multiplying a constant to all entries of a row. All of them preserve solutions to linear equations.
- Idea is to use row operations to get an **upper triangular** AM:

$$\left[\begin{array}{ccccc|c} * & - & - & - & - & - \\ 0 & * & - & - & - & - \\ 0 & 0 & * & - & - & - \\ 0 & 0 & 0 & * & - & - \end{array} \right]$$

- The * positions are called **leading entries**, and are the left-most non-zero entry of each row.

● **Definition:**(*Echelon form*) A matrix is said to be in Echelon form if

- ▶ All non-zero rows are above any zero rows (if any).
- ▶ Each leading entry (leftmost non-zero entry) in a row is in a column to the right of the leading entry of the row above it.

● Examples of Echelon forms:

$$\begin{bmatrix} * & - & - & - & - & - \\ 0 & 0 & * & - & - & - \\ 0 & 0 & 0 & * & - & - \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} * & - & - & - & - \\ 0 & 0 & * & - & - \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} * & - & - \\ 0 & 0 & * \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

● **Definition:**(*Echelon form*) A matrix is said to be in Echelon form if

- ▶ All non-zero rows are above any zero rows (if any).
- ▶ Each leading entry (leftmost non-zero entry) in a row is in a column to the right of the leading entry of the row above it.

● Examples of Echelon forms:

$$\begin{bmatrix} * & - & - & - & - & - \\ 0 & 0 & * & - & - & - \\ 0 & 0 & 0 & * & - & - \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} * & - & - & - & - \\ 0 & 0 & * & - & - \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} * & - & - \\ 0 & 0 & * \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

● Examples of non-Echelon forms:

$$\begin{bmatrix} * & - & - & - & - & - \\ 0 & 0 & * & - & - & - \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & * & - & - \end{bmatrix} \begin{bmatrix} * & - & - & - & - & - \\ 0 & 0 & 0 & * & - & - \\ 0 & 0 & 0 & * & - & - \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} * & - & - \\ 0 & 0 & * \\ 0 & * & - \\ 0 & 0 & 0 \end{bmatrix}$$

- **Definition:** (*Row-reduced Echelon form (RREF)*) A matrix in Echelon form is said to be in a Row-Reduced Echelon Form if
- ▶ If the leading entry in each row is 1
 - ▶ If the leading term is the only non-zero term in that column.

- **Definition:** (*Row-reduced Echelon form (RREF)*) A matrix in Echelon form is said to be in a Row-Reduced Echelon Form if
 - ▶ If the leading entry in each row is 1
 - ▶ If the leading term is the only non-zero term in that column.
- Examples of RREF:

$$\begin{bmatrix} * & - & 0 & 0 & - & - \\ 0 & 0 & * & 0 & - & - \\ 0 & 0 & 0 & * & - & - \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad \begin{bmatrix} * & - & 0 & 0 & - & - \\ 0 & 0 & * & 0 & - & - \\ 0 & 0 & 0 & * & - & - \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad \begin{bmatrix} * & - & 0 \\ 0 & 0 & * \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

- Let us use the following example:

$$2x_2 + 5x_3 + 4x_4 + 2x_5 = 2$$

$$x_1 - x_2 + 2x_3 + 3x_4 - x_5 = 1$$

$$2x_1 + x_2 + 4x_4 + 2x_5 = -1$$

$$3x_1 + x_2 + 3x_3 + 2x_4 - 2x_5 = 3$$

- Representation as a matrix

$$\left[\begin{array}{ccccc|c} 0 & 2 & 5 & 4 & 2 & 2 \\ 1 & -1 & 2 & 3 & -1 & 1 \\ 2 & 1 & 0 & 4 & 2 & -1 \\ 3 & 1 & 3 & 2 & -2 & 3 \end{array} \right]$$

Solving Linear Equations

$$\left[\begin{array}{ccccc|c} 0 & 2 & 5 & 4 & 2 & 2 \\ 1 & -1 & 2 & 3 & -1 & 1 \\ 2 & 1 & 0 & 4 & 2 & -1 \\ 3 & 1 & 3 & 2 & -2 & 3 \end{array} \right]$$

Solving Linear Equations

$$\left[\begin{array}{ccccc|c} 0 & 2 & 5 & 4 & 2 & 2 \\ 1 & -1 & 2 & 3 & -1 & 1 \\ 2 & 1 & 0 & 4 & 2 & -1 \\ 3 & 1 & 3 & 2 & -2 & 3 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_2} \left[\begin{array}{ccccc|c} \mathbf{1} & -1 & 2 & 3 & -1 & 1 \\ 0 & 2 & 5 & 4 & 2 & 2 \\ 2 & 1 & 0 & 4 & 2 & -1 \\ 3 & 1 & 3 & 2 & -2 & 3 \end{array} \right]$$

Solving Linear Equations

$$\left[\begin{array}{ccccc|c} 0 & 2 & 5 & 4 & 2 & 2 \\ 1 & -1 & 2 & 3 & -1 & 1 \\ 2 & 1 & 0 & 4 & 2 & -1 \\ 3 & 1 & 3 & 2 & -2 & 3 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_2} \left[\begin{array}{ccccc|c} \mathbf{1} & -1 & 2 & 3 & -1 & 1 \\ 0 & 2 & 5 & 4 & 2 & 2 \\ 2 & 1 & 0 & 4 & 2 & -1 \\ 3 & 1 & 3 & 2 & -2 & 3 \end{array} \right] \xrightarrow[\begin{array}{l} R_4 \leftarrow R_4 - 3R_1 \\ R_3 \leftarrow R_3 - 2R_1 \end{array}]{}$$
$$\left[\begin{array}{ccccc|c} \mathbf{1} & -1 & 2 & 3 & -1 & 1 \\ 0 & \mathbf{2} & 5 & 4 & 2 & 2 \\ 0 & 3 & -4 & -2 & 4 & -3 \\ 0 & 4 & -3 & -7 & 1 & 0 \end{array} \right]$$

Solving Linear Equations

$$\begin{aligned} & \left[\begin{array}{ccccc|c} 0 & 2 & 5 & 4 & 2 & 2 \\ 1 & -1 & 2 & 3 & -1 & 1 \\ 2 & 1 & 0 & 4 & 2 & -1 \\ 3 & 1 & 3 & 2 & -2 & 3 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_2} \left[\begin{array}{ccccc|c} 1 & -1 & 2 & 3 & -1 & 1 \\ 0 & 2 & 5 & 4 & 2 & 2 \\ 2 & 1 & 0 & 4 & 2 & -1 \\ 3 & 1 & 3 & 2 & -2 & 3 \end{array} \right] \xrightarrow{\begin{array}{l} R_4 \leftarrow R_4 - 3R_1 \\ R_3 \leftarrow R_3 - 2R_1 \end{array}} \\ & \left[\begin{array}{ccccc|c} 1 & -1 & 2 & 3 & -1 & 1 \\ 0 & 2 & 5 & 4 & 2 & 2 \\ 0 & 3 & -4 & -2 & 4 & -3 \\ 0 & 4 & -3 & -7 & 1 & 0 \end{array} \right] \xrightarrow{\begin{array}{l} R_4 \leftarrow R_4 - 2R_2 \\ R_3 \leftarrow 2R_3 - 3R_2 \end{array}} \left[\begin{array}{ccccc|c} 1 & -1 & 2 & 3 & -1 & 1 \\ 0 & 2 & 5 & 4 & 2 & 2 \\ 0 & 0 & -23 & -16 & 2 & -12 \\ 0 & 0 & -13 & -15 & -3 & -4 \end{array} \right] \end{aligned}$$

Solving Linear Equations

$$\begin{aligned} & \left[\begin{array}{ccccc|c} 0 & 2 & 5 & 4 & 2 & 2 \\ 1 & -1 & 2 & 3 & -1 & 1 \\ 2 & 1 & 0 & 4 & 2 & -1 \\ 3 & 1 & 3 & 2 & -2 & 3 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_2} \left[\begin{array}{ccccc|c} 1 & -1 & 2 & 3 & -1 & 1 \\ 0 & 2 & 5 & 4 & 2 & 2 \\ 2 & 1 & 0 & 4 & 2 & -1 \\ 3 & 1 & 3 & 2 & -2 & 3 \end{array} \right] \xrightarrow{\begin{array}{l} R_4 \leftarrow R_4 - 3R_1 \\ R_3 \leftarrow R_3 - 2R_1 \end{array}} \\ & \left[\begin{array}{ccccc|c} 1 & -1 & 2 & 3 & -1 & 1 \\ 0 & 2 & 5 & 4 & 2 & 2 \\ 0 & 3 & -4 & -2 & 4 & -3 \\ 0 & 4 & -3 & -7 & 1 & 0 \end{array} \right] \xrightarrow{\begin{array}{l} R_4 \leftarrow R_4 - 2R_2 \\ R_3 \leftarrow 2R_3 - 3R_2 \end{array}} \left[\begin{array}{ccccc|c} 1 & -1 & 2 & 3 & -1 & 1 \\ 0 & 2 & 5 & 4 & 2 & 2 \\ 0 & 0 & -23 & -16 & 2 & -12 \\ 0 & 0 & -13 & -15 & -3 & -4 \end{array} \right] \\ & \xrightarrow{R_4 \leftarrow -23R_4 - 13R_3} \underbrace{\left[\begin{array}{ccccc|c} 1 & -1 & 2 & 3 & -1 & 1 \\ 0 & 2 & 5 & 4 & 2 & 2 \\ 0 & 0 & -23 & -16 & 2 & -12 \\ 0 & 0 & 0 & -137 & -95 & 64 \end{array} \right]}_U \end{aligned}$$

- Any Elementary row operation can be represented as a matrix.

- ▶ Any Elementary row operation can be represented as a matrix.
- ▶ These matrices will be called *Elementary Row Transformations*.

- ▶ Any Elementary row operation can be represented as a matrix.
- ▶ These matrices will be called *Elementary Row Transformations*.

$$(R_1 \leftrightarrow R_2) \rightarrow$$

- ▶ Any Elementary row operation can be represented as a matrix.
- ▶ These matrices will be called *Elementary Row Transformations*.

$$(R_1 \leftrightarrow R_2) \rightarrow P_{12} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$(R_3 \leftarrow 2R_3 - 3R_2) \rightarrow$$

- Any Elementary row operation can be represented as a matrix.
- These matrices will be called *Elementary Row Transformations*.

$$(R_1 \leftrightarrow R_2) \rightarrow P_{12} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$(R_3 \leftarrow 2R_3 - 3R_2) \rightarrow L_{32} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & -3 & 2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

- Any Elementary row operation can be represented as a matrix.
- These matrices will be called *Elementary Row Transformations*.

$$(R_1 \leftrightarrow R_2) \rightarrow P_{12} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$(R_3 \leftarrow 2R_3 - 3R_2) \rightarrow L_{32} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & -3 & 2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

- Notice that each elementary row operation except permutation can be written using a *lower triangular matrix*.

- Any Elementary row operation can be represented as a matrix.
- These matrices will be called *Elementary Row Transformations*.

$$(R_1 \leftrightarrow R_2) \rightarrow P_{12} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$(R_3 \leftarrow 2R_3 - 3R_2) \rightarrow L_{32} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & -3 & 2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

- Notice that each elementary row operation except permutation can be written using a *lower triangular matrix*.
- $U = L_{43}L_{32}L_{42}L_{31}L_{41}P_{12}A$

- Any Elementary row operation can be represented as a matrix.
- These matrices will be called *Elementary Row Transformations*.

$$(R_1 \leftrightarrow R_2) \rightarrow P_{12} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$(R_3 \leftarrow 2R_3 - 3R_2) \rightarrow L_{32} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & -3 & 2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

- Notice that each elementary row operation except permutation can be written using a *lower triangular matrix*.
- $U = L_{43}L_{32}L_{42}L_{31}L_{41}P_{12}A$

Properties of ERT

- Are all elementary row transformations invertible?

Properties of ERT

- Are all elementary row transformations invertible?
- L_{32}^{-1}

Properties of ERT

- Are all elementary row transformations invertible?

- $L_{32}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 3/2 & 1/2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$

Properties of ERT

- Are all elementary row transformations invertible?

- $L_{32}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 3/2 & 1/2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$

- Moreover, the inverse is also lower triangular!

Properties of ERT

- Are all elementary row transformations invertible?

- $L_{32}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 3/2 & 1/2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$

- Moreover, the inverse is also lower triangular!
- What matrix will you get if you multiply two lower triangular matrices?

Properties of ERT

- Are all elementary row transformations invertible?

- $L_{32}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 3/2 & 1/2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$

- Moreover, the inverse is also lower triangular!
- What matrix will you get if you multiply two lower triangular matrices? → Lower Triangular!

Properties of ERT

- Are all elementary row transformations invertible?

- $L_{32}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 3/2 & 1/2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$

- Moreover, the inverse is also lower triangular!
- What matrix will you get if you multiply two lower triangular matrices? → Lower Triangular!
- Thus, $P_{12}A = L_{43}^{-1}L_{32}^{-1}L_{42}^{-1}L_{31}^{-1}L_{41}^{-1}U = LU.$

LU Decomposition

- In general, any matrix $A \in \mathbb{R}^{m \times n}$ can be decomposed into a product of lower and upper triangular matrices, with appropriate permutations:

$$PA = LU,$$

where $P \in \mathbb{R}^{m \times m}$, $L \in \mathbb{R}^{m \times m}$, $U \in \mathbb{R}^{m \times n}$.

LU Decomposition

- In general, any matrix $A \in \mathbb{R}^{m \times n}$ can be decomposed into a product of lower and upper triangular matrices, with appropriate permutations:

$$PA = LU,$$

where $P \in \mathbb{R}^{m \times m}$, $L \in \mathbb{R}^{m \times m}$, $U \in \mathbb{R}^{m \times n}$.

- Solving linear equations with LU decomposition:
 $Ax = b \rightarrow LUx = b$.

LU Decomposition

- In general, any matrix $A \in \mathbb{R}^{m \times n}$ can be decomposed into a product of lower and upper triangular matrices, with appropriate permutations:

$$PA = LU,$$

where $P \in \mathbb{R}^{m \times m}$, $L \in \mathbb{R}^{m \times m}$, $U \in \mathbb{R}^{m \times n}$.

- Solving linear equations with LU decomposition:

$$Ax = b \rightarrow LUx = b.$$

- First let $Ux = y$ and solve $Ly = b$, and next solve for x in $Ux = y$.