

SC223 - Linear Algebra

Aditya Tatu

Lecture 7



August 25, 2022

Solving Linear Equations

$$\begin{aligned}
 & \left[\begin{array}{ccccc|c} 0 & 2 & 5 & 4 & 2 & 2 \\ 1 & -1 & 2 & 3 & -1 & 1 \\ 2 & 1 & 0 & 4 & 2 & -1 \\ 3 & 1 & 3 & 2 & -2 & 3 \end{array} \right] \xrightarrow{R_1 \leftrightarrow R_2} \left[\begin{array}{ccccc|c} 1 & -1 & 2 & 3 & -1 & 1 \\ 0 & 2 & 5 & 4 & 2 & 2 \\ 2 & 1 & 0 & 4 & 2 & -1 \\ 3 & 1 & 3 & 2 & -2 & 3 \end{array} \right] \xrightarrow{\begin{array}{l} R_4 \leftarrow R_4 - 3R_1 \\ R_3 \leftarrow R_3 - 2R_1 \end{array}} \\
 & \left[\begin{array}{ccccc|c} 1 & -1 & 2 & 3 & -1 & 1 \\ 0 & 2 & 5 & 4 & 2 & 2 \\ 0 & 3 & -4 & -2 & 4 & -3 \\ 0 & 4 & -3 & -7 & 1 & 0 \end{array} \right] \xrightarrow{\begin{array}{l} R_4 \leftarrow R_4 - 2R_2 \\ R_3 \leftarrow 2R_3 - 3R_2 \end{array}} \left[\begin{array}{ccccc|c} 1 & -1 & 2 & 3 & -1 & 1 \\ 0 & 2 & 5 & 4 & 2 & 2 \\ 0 & 0 & -23 & -16 & 2 & -12 \\ 0 & 0 & -13 & -15 & -3 & -4 \end{array} \right] \\
 & \xrightarrow{R_4 \leftarrow 23R_4 - 13R_3} \underbrace{\left[\begin{array}{ccccc|c} 1 & -1 & 2 & 3 & -1 & 1 \\ 0 & 2 & 5 & 4 & 2 & 2 \\ 0 & 0 & -23 & -16 & 2 & -12 \\ 0 & 0 & 0 & -137 & -95 & 64 \end{array} \right]}_{U \text{ (except the last column)}}
 \end{aligned}$$

- ▶ Any Elementary row operation can be represented as a matrix.
- ▶ These matrices will be called *Elementary Row Transformations*.

$$(R_1 \leftrightarrow R_2) \rightarrow P_{12} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$(R_3 \leftarrow 2R_3 - 3R_2) \rightarrow$$

- Any Elementary row operation can be represented as a matrix.
- These matrices will be called *Elementary Row Transformations*.

$$(R_1 \leftrightarrow R_2) \rightarrow P_{12} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$(R_3 \leftarrow 2R_3 - 3R_2) \rightarrow L_{32} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & -3 & 2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

- Any Elementary row operation can be represented as a matrix.
- These matrices will be called *Elementary Row Transformations*.

$$(R_1 \leftrightarrow R_2) \rightarrow P_{12} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$
$$(R_3 \leftarrow 2R_3 - 3R_2) \rightarrow L_{32} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & -3 & 2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

- Notice that each elementary row operation except permutation can be written using a *lower triangular matrix*.

- Any Elementary row operation can be represented as a matrix.
- These matrices will be called *Elementary Row Transformations*.

$$(R_1 \leftrightarrow R_2) \rightarrow P_{12} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$(R_3 \leftarrow 2R_3 - 3R_2) \rightarrow L_{32} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & -3 & 2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

- Notice that each elementary row operation except permutation can be written using a *lower triangular matrix*.
- $U = L_{43}L_{32}L_{42}L_{31}L_{41}P_{12}A$.

Properties of ERT

- $L_{32} \rightarrow (R_3 \leftarrow 2R_3 - 3R_2)$. Is this invertible?

Properties of ERT

- $L_{32} \rightarrow (R_3 \leftarrow 2R_3 - 3R_2)$. Is this invertible? Can I get back R_3 from $2R_3 - 3R_2$?

Properties of ERT

- $L_{32} \rightarrow (R_3 \leftarrow 2R_3 - 3R_2)$. Is this invertible? Can I get back R_3 from $2R_3 - 3R_2$?
- L_{32}^{-1}

Properties of ERT

- $L_{32} \rightarrow (R_3 \leftarrow 2R_3 - 3R_2)$. Is this invertible? Can I get back R_3 from $2R_3 - 3R_2$?

- $L_{32}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 3/2 & 1/2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$

Properties of ERT

- $L_{32} \rightarrow (R_3 \leftarrow 2R_3 - 3R_2)$. Is this invertible? Can I get back R_3 from $2R_3 - 3R_2$?

- $L_{32}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 3/2 & 1/2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$

- Are all elementary row transformations invertible?

Properties of ERT

- $L_{32} \rightarrow (R_3 \leftarrow 2R_3 - 3R_2)$. Is this invertible? Can I get back R_3 from $2R_3 - 3R_2$?

- $L_{32}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 3/2 & 1/2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$

- Are all elementary row transformations invertible?
- Moreover, the inverse is also lower triangular! (except for permutation RT)

Properties of ERT

- $L_{32} \rightarrow (R_3 \leftarrow 2R_3 - 3R_2)$. Is this invertible? Can I get back R_3 from $2R_3 - 3R_2$?

- $L_{32}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 3/2 & 1/2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$

- Are all elementary row transformations invertible?
- Moreover, the inverse is also lower triangular! (except for permutation RT)
- What matrix will you get if you multiply two lower triangular matrices?

Properties of ERT

- $L_{32} \rightarrow (R_3 \leftarrow 2R_3 - 3R_2)$. Is this invertible? Can I get back R_3 from $2R_3 - 3R_2$?

- $L_{32}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 3/2 & 1/2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$

- Are all elementary row transformations invertible?
- Moreover, the inverse is also lower triangular! (except for permutation RT)
- What matrix will you get if you multiply two lower triangular matrices? → Lower Triangular!

Properties of ERT

- $L_{32} \rightarrow (R_3 \leftarrow 2R_3 - 3R_2)$. Is this invertible? Can I get back R_3 from $2R_3 - 3R_2$?

- $L_{32}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 3/2 & 1/2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}.$

- Are all elementary row transformations invertible?
- Moreover, the inverse is also lower triangular! (except for permutation RT)
- What matrix will you get if you multiply two lower triangular matrices? \rightarrow Lower Triangular!
- Thus, $P_{12}A = L_{43}^{-1}L_{32}^{-1}L_{42}^{-1}L_{31}^{-1}L_{41}^{-1}U = LU.$

LU Decomposition

- Any matrix $A \in \mathbb{R}^{m \times n}$ can be decomposed into a product of lower and upper triangular matrices, with appropriate permutations:

$$PA = LU,$$

where $P \in \mathbb{R}^{m \times m}$, $L \in \mathbb{R}^{m \times m}$, $U \in \mathbb{R}^{m \times n}$.

LU Decomposition

- Any matrix $A \in \mathbb{R}^{m \times n}$ can be decomposed into a product of lower and upper triangular matrices, with appropriate permutations:

$$PA = LU,$$

where $P \in \mathbb{R}^{m \times m}$, $L \in \mathbb{R}^{m \times m}$, $U \in \mathbb{R}^{m \times n}$.

- Solving linear equations with LU decomposition:
 $Ax = b \rightarrow LUx = b$.

LU Decomposition

- Any matrix $A \in \mathbb{R}^{m \times n}$ can be decomposed into a product of lower and upper triangular matrices, with appropriate permutations:

$$PA = LU,$$

where $P \in \mathbb{R}^{m \times m}$, $L \in \mathbb{R}^{m \times m}$, $U \in \mathbb{R}^{m \times n}$.

- Solving linear equations with LU decomposition:

$$Ax = b \rightarrow LUx = b.$$

- First let $Ux = y$ and solve $Ly = b$, and next solve for x in $Ux = y$.

Example

$$\bullet \quad U = \begin{bmatrix} \mathbf{1} & -1 & 2 & 3 & -1 \\ 0 & \mathbf{2} & 5 & 4 & 2 \\ 0 & 0 & -\mathbf{23} & -16 & 2 \\ 0 & 0 & 0 & -\mathbf{137} & -95 \end{bmatrix}, y = \begin{bmatrix} 1 \\ 2 \\ -12 \\ 64 \end{bmatrix}.$$

Example

$$\bullet U = \begin{bmatrix} \mathbf{1} & -1 & 2 & 3 & -1 \\ 0 & \mathbf{2} & 5 & 4 & 2 \\ 0 & 0 & -\mathbf{23} & -16 & 2 \\ 0 & 0 & 0 & -\mathbf{137} & -95 \end{bmatrix}, y = \begin{bmatrix} 1 \\ 2 \\ -12 \\ 64 \end{bmatrix}.$$

● Solve for x in

$$\begin{bmatrix} \mathbf{1} & -1 & 2 & 3 & -1 \\ 0 & \mathbf{2} & 5 & 4 & 2 \\ 0 & 0 & -\mathbf{23} & -16 & 2 \\ 0 & 0 & 0 & -\mathbf{137} & -95 \end{bmatrix} x = \begin{bmatrix} 1 \\ 2 \\ -12 \\ 64 \end{bmatrix}$$

Computing Matrix Inverses

- A^{-1} is the unique matrix such that $A^{-1}A = AA^{-1} = I$.

Computing Matrix Inverses

- A^{-1} is the unique matrix such that $A^{-1}A = AA^{-1} = I$.
- Simultaneously solve n linear equations:

$$\underbrace{\begin{bmatrix} a_{*1} & \dots & a_{*n} \\ a_{*1} & \dots & a_{*n} \end{bmatrix}}_{AM} \begin{bmatrix} x_1 & \dots & x_n \end{bmatrix} = I, x_i \in \mathbb{R}^n, i = 1, \dots, n$$
$$\begin{bmatrix} LU & | & I \end{bmatrix} \Rightarrow LUx_1 = I_{*1}, \dots, LUx_n = I_{*n}.$$

Computing Matrix Inverses

- A^{-1} is the unique matrix such that $A^{-1}A = AA^{-1} = I$.
- Simultaneously solve n linear equations:

$$\underbrace{\begin{bmatrix} a_{*1} & \dots & a_{*n} \end{bmatrix}}_{AM} \begin{bmatrix} x_1 & \dots & x_n \end{bmatrix} = I, x_i \in \mathbb{R}^n, i = 1, \dots, n$$
$$\begin{bmatrix} LU & | & I \end{bmatrix} \Rightarrow LUx_1 = I_{*1}, \dots, LUx_n = I_{*n}.$$

- Gauss-Jordan Method: Let R_1, \dots, R_k represent row transformation matrices, not necessarily lower triangular, such that $R_k \cdot R_{k-1} \cdot \dots \cdot R_1 A = I$, then $A^{-1} =$

Computing Matrix Inverses

- A^{-1} is the unique matrix such that $A^{-1}A = AA^{-1} = I$.
- Simultaneously solve n linear equations:

$$\underbrace{\begin{bmatrix} a_{*1} & \dots & a_{*n} \end{bmatrix}}_{AM} \begin{bmatrix} x_1 & \dots & x_n \end{bmatrix} = I, x_i \in \mathbb{R}^n, i = 1, \dots, n$$
$$\begin{bmatrix} LU & | & I \end{bmatrix} \Rightarrow LUx_1 = I_{*1}, \dots, LUx_n = I_{*n}.$$

- Gauss-Jordan Method: Let R_1, \dots, R_k represent row transformation matrices, not necessarily lower triangular, such that $R_k \cdot R_{k-1} \cdot \dots \cdot R_1 A = I$, then $A^{-1} = R_k \cdot R_{k-1} \cdot \dots \cdot R_1$.

Computing Matrix Inverses

- A^{-1} is the unique matrix such that $A^{-1}A = AA^{-1} = I$.
- Simultaneously solve n linear equations:

$$\underbrace{\begin{bmatrix} a_{*1} & \dots & a_{*n} \\ a_{*1} & \dots & a_{*n} \end{bmatrix}}_{AM} \begin{bmatrix} x_1 & \dots & x_n \end{bmatrix} = I, x_i \in \mathbb{R}^n, i = 1, \dots, n$$
$$\begin{bmatrix} LU & | & I \end{bmatrix} \Rightarrow LUx_1 = I_{*1}, \dots, LUx_n = I_{*n}.$$

- Gauss-Jordan Method: Let R_1, \dots, R_k represent row transformation matrices, not necessarily lower triangular, such that $R_k \cdot R_{k-1} \cdot \dots \cdot R_1 A = I$, then $A^{-1} = R_k \cdot R_{k-1} \cdot \dots \cdot R_1$.
- Thus,

$$R_k \cdot R_{k-1} \cdot \dots \cdot R_1 \begin{bmatrix} A & | & I \end{bmatrix} =$$

Computing Matrix Inverses

- A^{-1} is the unique matrix such that $A^{-1}A = AA^{-1} = I$.
- Simultaneously solve n linear equations:

$$\underbrace{\begin{bmatrix} a_{*1} & \dots & a_{*n} \end{bmatrix}}_{AM} \begin{bmatrix} x_1 & \dots & x_n \end{bmatrix} = I, x_i \in \mathbb{R}^n, i = 1, \dots, n$$
$$\begin{bmatrix} LU & | & I \end{bmatrix} \Rightarrow LUx_1 = I_{*1}, \dots, LUx_n = I_{*n}.$$

- Gauss-Jordan Method: Let R_1, \dots, R_k represent row transformation matrices, not necessarily lower triangular, such that $R_k \cdot R_{k-1} \cdot \dots \cdot R_1 A = I$, then $A^{-1} = R_k \cdot R_{k-1} \cdot \dots \cdot R_1$.
- Thus,

$$R_k \cdot R_{k-1} \cdot \dots \cdot R_1 \begin{bmatrix} A & | & I \end{bmatrix} = \begin{bmatrix} I & | & A^{-1} \end{bmatrix}$$

Computational Cost

- Additions and multiplications in worst case in computing the LU decomposition?

Computational Cost

- Additions and multiplications in worst case in computing the LU decomposition?
- $\mathcal{O}(n^3)$

Computational Cost

- Additions and multiplications in worst case in computing the LU decomposition?
- $\mathcal{O}(n^3)$
- Cost involved in Back-substitution?

Computational Cost

- Additions and multiplications in worst case in computing the LU decomposition?
- $\mathcal{O}(n^3)$
- Cost involved in Back-substitution?
- $\mathcal{O}(n^2)$

Computational Cost

- Additions and multiplications in worst case in computing the LU decomposition?
- $\mathcal{O}(n^3)$
- Cost involved in Back-substitution?
- $\mathcal{O}(n^2)$
- Cost for computing A^{-1} using Gauss-Jordan method?

Computational Cost

- Additions and multiplications in worst case in computing the LU decomposition?
- $\mathcal{O}(n^3)$
- Cost involved in Back-substitution?
- $\mathcal{O}(n^2)$
- Cost for computing A^{-1} using Gauss-Jordan method?
- $\mathcal{O}(n^3)$

Computational Cost

- Additions and multiplications in worst case in computing the LU decomposition?
- $\mathcal{O}(n^3)$
- Cost involved in Back-substitution?
- $\mathcal{O}(n^2)$
- Cost for computing A^{-1} using Gauss-Jordan method?
- $\mathcal{O}(n^3)$
- Why should one use LU decomposition?