

SC223 - Linear Algebra

Aditya Tatu

Lecture 1



August 2, 2022

What is Linear algebra?

What is Linear algebra?

- It is the study of *structures* that behave like a *line*.

What is a Structure?

What is a Structure?



What is a Structure?



Figure: Even though all of these are different, but we recognize them as having the *structure* of a chair. Image source: freepik.com

Example

- Let $A = \{a, b, c\}$. Equip the set $\mathcal{P}(A)$ with the binary relation:
 $xRy \Leftrightarrow x \subseteq y$.

Example

- Let $A = \{a, b, c\}$. Equip the set $\mathcal{P}(A)$ with the binary relation: $xRy \Leftrightarrow x \subseteq y$.
- Let $B = \{1, 2, 3, 5, 6, 10, 15, 30\}$ consist of all factors of 30. Define the binary relation: $xRy \Leftrightarrow x|y$.

Example

- Let $A = \{a, b, c\}$. Equip the set $\mathcal{P}(A)$ with the binary relation: $xRy \Leftrightarrow x \subseteq y$.
- Let $B = \{1, 2, 3, 5, 6, 10, 15, 30\}$ consist of all factors of 30. Define the binary relation: $xRy \Leftrightarrow x|y$.
- What are $(\mathcal{P}(A), \subseteq)$, $(B, |)$? Is there any similarity between the two?

Example

- Let $A = \{a, b, c\}$. Equip the set $\mathcal{P}(A)$ with the binary relation: $xRy \Leftrightarrow x \subseteq y$.
- Let $B = \{1, 2, 3, 5, 6, 10, 15, 30\}$ consist of all factors of 30. Define the binary relation: $xRy \Leftrightarrow x|y$.
- What are $(\mathcal{P}(A), \subseteq)$, $(B, |)$? Is there any similarity between the two?





Figure: Hasse Diagram for $(\mathcal{P}(A), \subseteq)$ and $(B, |)$





Figure: Hasse Diagram for $(\mathcal{P}(A), \subseteq)$ and $(B, |)$

- Although the elements are different, the relationship between them is same!





Figure: Hasse Diagram for $(\mathcal{P}(A), \subseteq)$ and $(B, |)$

- Although the elements are different, the relationship between them is same!
- They have the structure of a *Poset*.

Linear Algebra

- System of Linear Equations.
- Vector Spaces.
- Linear Transformations and Matrices.
- Eigenvalues and Eigenvectors.
- Inner Products and Norms.
- Complex Vector Spaces.

Linear Equations

- Linear systems in 2 variables: Solve for x and y

$$2x + 3y = 5$$

$$x - 5y = 10$$

Linear Equations

- Linear systems in 2 variables: Solve for x and y

$$2x + 3y = 5$$

$$x - 5y = 10$$

- Linear systems in 3 variables: Solve for x, y and z

$$2x + 3y - z = 5$$

$$x - 5y + 2z = 10$$

$$3x + 2y + z = 1$$

Linear Equations

- We will now study them in the form:

$$\begin{bmatrix} 2 & 3 \\ 1 & -5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 5 \\ 10 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 3 & -1 \\ 1 & -5 & 2 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 5 \\ 10 \\ 1 \end{bmatrix}$$

Linear Equations

- We will now study them in the form:

$$\begin{bmatrix} 2 & 3 \\ 1 & -5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 5 \\ 10 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 3 & -1 \\ 1 & -5 & 2 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 5 \\ 10 \\ 1 \end{bmatrix}$$

- In general, solve for x in

$$\underbrace{\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}}_A \underbrace{\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}}_x = \underbrace{\begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}}_b,$$

where $A \in \mathbb{R}^{m \times n}$, $x \in \mathbb{R}^n$ and $b \in \mathbb{R}^m$.

What are we interested in?

- How to solve $Ax = b$?

What are we interested in?

- How to solve $Ax = b$?
- Under what conditions is it possible to solve a given $Ax = b$?

What are we interested in?

- How to solve $Ax = b$?
- Under what conditions is it possible to solve a given $Ax = b$?
- Under what conditions is it not possible to solve a given $Ax = b$?

What are we interested in?

- How to solve $Ax = b$?
- Under what conditions is it possible to solve a given $Ax = b$?
- Under what conditions is it not possible to solve a given $Ax = b$?
- Under what conditions can there be multiple solutions to $Ax = b$?

What are we interested in?

- How to solve $Ax = b$?
- Under what conditions is it possible to solve a given $Ax = b$?
- Under what conditions is it not possible to solve a given $Ax = b$?
- Under what conditions can there be multiple solutions to $Ax = b$?
- What can be done if no solution exists?

What are we interested in?

- How to solve $Ax = b$?
- Under what conditions is it possible to solve a given $Ax = b$?
- Under what conditions is it not possible to solve a given $Ax = b$?
- Under what conditions can there be multiple solutions to $Ax = b$?
- What can be done if no solution exists?
- What is the Computational cost of the algorithm to solve $Ax = b$.

Row Picture

- Let us look at each row of the system:

$$\begin{bmatrix} 2 & 3 \\ 1 & -5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 5 \\ 10 \end{bmatrix}.$$

Possibilities for a 2×2 system

Possibilities for a 3×3 system

Column Picture

$$\begin{bmatrix} 2 & 3 \\ 1 & -5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 5 \\ 10 \end{bmatrix}$$

can be re-written as

$$x \cdot \begin{bmatrix} 2 \\ 1 \end{bmatrix} + y \cdot \begin{bmatrix} 3 \\ -5 \end{bmatrix} = \begin{bmatrix} 5 \\ 10 \end{bmatrix},$$

where $x \cdot \begin{bmatrix} a \\ b \end{bmatrix} := \begin{bmatrix} ax \\ bx \end{bmatrix}$, and $\begin{bmatrix} a \\ b \end{bmatrix} + \begin{bmatrix} c \\ d \end{bmatrix} := \begin{bmatrix} a + c \\ b + d \end{bmatrix}$.

- In general, for an $m \times n$ system of linear equations $Ax = b$,

$$x_1 \cdot \underbrace{\begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix}}_{a_{*1} \in \mathbb{R}^n} + x_2 \cdot \underbrace{\begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix}}_{a_{*2} \in \mathbb{R}^n} + \dots + x_n \cdot \underbrace{\begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix}}_{a_{*n} \in \mathbb{R}^n} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

- **Linear combination** of a_{*i} and a_{*j} with real numbers x_i and x_j is defined as

$$x_i \cdot \begin{bmatrix} a_{1i} \\ a_{2i} \\ \vdots \\ a_{mi} \end{bmatrix} + x_j \cdot \begin{bmatrix} a_{1j} \\ a_{2j} \\ \vdots \\ a_{mj} \end{bmatrix} = \begin{bmatrix} x_i a_{1i} + x_j a_{1j} \\ x_i a_{2i} + x_j a_{2j} \\ \vdots \\ x_i a_{mi} + x_j a_{mj} \end{bmatrix}$$

When does a solution exist?

When does a solution exist?

- Solution to $Ax = b$ exists if and only if b belongs to the set of *all possible linear combinations* of columns of A .

When does a solution exist?

- ▶ Solution to $Ax = b$ exists if and only if b belongs to the set of *all possible linear combinations* of columns of A .
- ▶ **Column Space:** The set of all possible linear combinations of columns of A is called the Column space of matrix A , and is denoted by $C(A)$.

$$C(A) = \{Ax \mid \forall x \in \mathbb{R}^n\}, C(A) \subseteq \mathbb{R}^m$$

When does a solution exist?

- ▶ Solution to $Ax = b$ exists if and only if b belongs to the set of *all possible linear combinations* of columns of A .
- ▶ **Column Space:** The set of all possible linear combinations of columns of A is called the Column space of matrix A , and is denoted by $C(A)$.

$$C(A) = \{Ax \mid \forall x \in \mathbb{R}^n\}, C(A) \subseteq \mathbb{R}^m$$

- ▶ Properties:

When does a solution exist?

- ▶ Solution to $Ax = b$ exists if and only if b belongs to the set of *all possible linear combinations* of columns of A .
- ▶ **Column Space:** The set of all possible linear combinations of columns of A is called the Column space of matrix A , and is denoted by $C(A)$.

$$C(A) = \{Ax \mid \forall x \in \mathbb{R}^n\}, C(A) \subseteq \mathbb{R}^m$$

- ▶ Properties:

- ▶ Let $\mathbf{0}_m = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}_{m \times 1}$.

When does a solution exist?

- ▶ Solution to $Ax = b$ exists if and only if b belongs to the set of *all possible linear combinations* of columns of A .
- ▶ **Column Space:** The set of all possible linear combinations of columns of A is called the Column space of matrix A , and is denoted by $C(A)$.

$$C(A) = \{Ax \mid \forall x \in \mathbb{R}^n\}, C(A) \subseteq \mathbb{R}^m$$

- ▶ Properties:

- ▶ Let $\mathbf{0}_m = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}_{m \times 1}$. $\mathbf{0}_m \in C(A)$ for any matrix A

- ▶ If $b_1, b_2 \in C(A)$, $\forall p, q \in \mathbb{R}$, $p \cdot b_1 + q \cdot b_2$

When does a solution exist?

- ▶ Solution to $Ax = b$ exists if and only if b belongs to the set of *all possible linear combinations* of columns of A .
- ▶ **Column Space:** The set of all possible linear combinations of columns of A is called the Column space of matrix A , and is denoted by $C(A)$.

$$C(A) = \{Ax \mid \forall x \in \mathbb{R}^n\}, C(A) \subseteq \mathbb{R}^m$$

- ▶ Properties:

- ▶ Let $\mathbf{0}_m = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}_{m \times 1}$. $\mathbf{0}_m \in C(A)$ for any matrix A

- ▶ If $b_1, b_2 \in C(A)$, $\forall p, q \in \mathbb{R}$, $p \cdot b_1 + q \cdot b_2 \in C(A)$.

What about multiple solutions?

- Let $Ax = b$ and $Ay = b$, with $x \neq y$.

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) =$

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) =$

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!
- ▶ **Nullspace:** For a matrix $A \in \mathbb{R}^{m \times n}$, the *Nullspace* is the set of vectors that get mapped to $\mathbf{0}_m$, and is denoted by $N(A)$.

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!
- ▶ **Nullspace:** For a matrix $A \in \mathbb{R}^{m \times n}$, the *Nullspace* is the set of vectors that get mapped to $\mathbf{0}_m$, and is denoted by $N(A)$.

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶ $Ax = \mathbf{0}_m$ are also called **Homogeneous equations**.

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!
- ▶ **Nullspace:** For a matrix $A \in \mathbb{R}^{m \times n}$, the *Nullspace* is the set of vectors that get mapped to $\mathbf{0}_m$, and is denoted by $N(A)$.

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶ $Ax = \mathbf{0}_m$ are also called **Homogeneous equations**.
- ▶ Properties:

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!
- ▶ **Nullspace:** For a matrix $A \in \mathbb{R}^{m \times n}$, the *Nullspace* is the set of vectors that get mapped to $\mathbf{0}_m$, and is denoted by $N(A)$.

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶ $Ax = \mathbf{0}_m$ are also called **Homogeneous equations**.
- ▶ Properties:
- ▶ $\mathbf{0}_n$

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!
- ▶ **Nullspace:** For a matrix $A \in \mathbb{R}^{m \times n}$, the *Nullspace* is the set of vectors that get mapped to $\mathbf{0}_m$, and is denoted by $N(A)$.

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶ $Ax = \mathbf{0}_m$ are also called **Homogeneous equations**.
- ▶ Properties:
- ▶ $\mathbf{0}_n \in N(A)$.

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!
- ▶ **Nullspace:** For a matrix $A \in \mathbb{R}^{m \times n}$, the *Nullspace* is the set of vectors that get mapped to $\mathbf{0}_m$, and is denoted by $N(A)$.

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶ $Ax = \mathbf{0}_m$ are also called **Homogeneous equations**.
- ▶ Properties:
- ▶ $\mathbf{0}_n \in N(A)$.
- ▶ If $x, y \in N(A)$, $\forall p, q \in \mathbb{R}, p \cdot x + q \cdot y$

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!
- ▶ **Nullspace:** For a matrix $A \in \mathbb{R}^{m \times n}$, the *Nullspace* is the set of vectors that get mapped to $\mathbf{0}_m$, and is denoted by $N(A)$.

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶ $Ax = \mathbf{0}_m$ are also called **Homogeneous equations**.
- ▶ Properties:
- ▶ $\mathbf{0}_n \in N(A)$.
- ▶ If $x, y \in N(A)$, $\forall p, q \in \mathbb{R}, p \cdot x + q \cdot y \in N(A)$.

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!
- ▶ **Nullspace:** For a matrix $A \in \mathbb{R}^{m \times n}$, the *Nullspace* is the set of vectors that get mapped to $\mathbf{0}_m$, and is denoted by $N(A)$.

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶ $Ax = \mathbf{0}_m$ are also called **Homogeneous equations**.
- ▶ Properties:
- ▶ $\mathbf{0}_n \in N(A)$.
- ▶ If $x, y \in N(A)$, $\forall p, q \in \mathbb{R}$, $p \cdot x + q \cdot y \in N(A)$.
- ▶ If $\exists z \in N(A)$, $z \neq \mathbf{0}_n$, then $Ax = b$ will have

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!
- ▶ **Nullspace:** For a matrix $A \in \mathbb{R}^{m \times n}$, the *Nullspace* is the set of vectors that get mapped to $\mathbf{0}_m$, and is denoted by $N(A)$.

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶ $Ax = \mathbf{0}_m$ are also called **Homogeneous equations**.
- ▶ Properties:
- ▶ $\mathbf{0}_n \in N(A)$.
- ▶ If $x, y \in N(A)$, $\forall p, q \in \mathbb{R}$, $p \cdot x + q \cdot y \in N(A)$.
- ▶ If $\exists z \in N(A)$, $z \neq \mathbf{0}_n$, then $Ax = b$ will have infinitely many solutions, if one exists!