

SC223 - Linear Algebra

Aditya Tatu

Lecture 2



August 4, 2022

Row Picture

- Let us look at each row of the system:

$$\begin{bmatrix} 2 & 3 \\ 1 & -5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 5 \\ 10 \end{bmatrix}.$$

Possibilities for a 2×2 system

Possibilities for a 3×3 system

Column Picture

$$\begin{bmatrix} 2 & 3 \\ 1 & -5 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 5 \\ 10 \end{bmatrix}$$

can be re-written as

$$x \cdot \begin{bmatrix} 2 \\ 1 \end{bmatrix} + y \cdot \begin{bmatrix} 3 \\ -5 \end{bmatrix} = \begin{bmatrix} 5 \\ 10 \end{bmatrix},$$

where $x \cdot \begin{bmatrix} a \\ b \end{bmatrix} := \begin{bmatrix} ax \\ bx \end{bmatrix}$, and $\begin{bmatrix} a \\ b \end{bmatrix} + \begin{bmatrix} c \\ d \end{bmatrix} := \begin{bmatrix} a + c \\ b + d \end{bmatrix}$.

- In general, for an $m \times n$ system of linear equations $Ax = b$,

$$x_1 \cdot \underbrace{\begin{bmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{bmatrix}}_{a_{*1} \in \mathbb{R}^n} + x_2 \cdot \underbrace{\begin{bmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{bmatrix}}_{a_{*2} \in \mathbb{R}^n} + \dots + x_n \cdot \underbrace{\begin{bmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{bmatrix}}_{a_{*n} \in \mathbb{R}^n} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

- **Linear combination** of a_{*i} and a_{*j} with real numbers x_i and x_j is defined as

$$x_i \cdot \begin{bmatrix} a_{1i} \\ a_{2i} \\ \vdots \\ a_{mi} \end{bmatrix} + x_j \cdot \begin{bmatrix} a_{1j} \\ a_{2j} \\ \vdots \\ a_{mj} \end{bmatrix} = \begin{bmatrix} x_i a_{1i} + x_j a_{1j} \\ x_i a_{2i} + x_j a_{2j} \\ \vdots \\ x_i a_{mi} + x_j a_{mj} \end{bmatrix}$$

When does a solution exist?

When does a solution exist?

- Solution to $Ax = b$ exists if and only if b belongs to the set of *all possible linear combinations* of columns of A .

When does a solution exist?

- ▶ Solution to $Ax = b$ exists if and only if b belongs to the set of *all possible linear combinations* of columns of A .
- ▶ **Column Space:** The set of all possible linear combinations of columns of A is called the Column space of matrix A , and is denoted by $C(A)$.

$$C(A) = \{Ax \mid \forall x \in \mathbb{R}^n\}, C(A) \subseteq \mathbb{R}^m$$

When does a solution exist?

- ▶ Solution to $Ax = b$ exists if and only if b belongs to the set of *all possible linear combinations* of columns of A .
- ▶ **Column Space:** The set of all possible linear combinations of columns of A is called the Column space of matrix A , and is denoted by $C(A)$.

$$C(A) = \{Ax \mid \forall x \in \mathbb{R}^n\}, C(A) \subseteq \mathbb{R}^m$$

- ▶ Properties:

When does a solution exist?

- ▶ Solution to $Ax = b$ exists if and only if b belongs to the set of *all possible linear combinations* of columns of A .
- ▶ **Column Space:** The set of all possible linear combinations of columns of A is called the Column space of matrix A , and is denoted by $C(A)$.

$$C(A) = \{Ax \mid \forall x \in \mathbb{R}^n\}, C(A) \subseteq \mathbb{R}^m$$

- ▶ Properties:

- ▶ Let $\mathbf{0}_m = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}_{m \times 1}$.

When does a solution exist?

- ▶ Solution to $Ax = b$ exists if and only if b belongs to the set of *all possible linear combinations* of columns of A .
- ▶ **Column Space:** The set of all possible linear combinations of columns of A is called the Column space of matrix A , and is denoted by $C(A)$.

$$C(A) = \{Ax \mid \forall x \in \mathbb{R}^n\}, C(A) \subseteq \mathbb{R}^m$$

- ▶ Properties:

- ▶ Let $\mathbf{0}_m = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}_{m \times 1}$. $\mathbf{0}_m \in C(A)$ for any matrix A

- ▶ If $b_1, b_2 \in C(A)$, $\forall p, q \in \mathbb{R}$, $p \cdot b_1 + q \cdot b_2$

When does a solution exist?

- ▶ Solution to $Ax = b$ exists if and only if b belongs to the set of *all possible linear combinations* of columns of A .
- ▶ **Column Space:** The set of all possible linear combinations of columns of A is called the Column space of matrix A , and is denoted by $C(A)$.

$$C(A) = \{Ax \mid \forall x \in \mathbb{R}^n\}, C(A) \subseteq \mathbb{R}^m$$

- ▶ Properties:

- ▶ Let $\mathbf{0}_m = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}_{m \times 1}$. $\mathbf{0}_m \in C(A)$ for any matrix A

- ▶ If $b_1, b_2 \in C(A)$, $\forall p, q \in \mathbb{R}$, $p \cdot b_1 + q \cdot b_2 \in C(A)$.

What about multiple solutions?

- Let $Ax = b$ and $Ay = b$, with $x \neq y$.

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) =$

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) =$

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!
- ▶ **Nullspace:** For a matrix $A \in \mathbb{R}^{m \times n}$, the *Nullspace* is the set of vectors that get mapped to $\mathbf{0}_m$, and is denoted by $N(A)$.

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!
- ▶ **Nullspace:** For a matrix $A \in \mathbb{R}^{m \times n}$, the *Nullspace* is the set of vectors that get mapped to $\mathbf{0}_m$, and is denoted by $N(A)$.

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶ $Ax = \mathbf{0}_m$ are also called **Homogeneous equations**.

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!
- ▶ **Nullspace:** For a matrix $A \in \mathbb{R}^{m \times n}$, the *Nullspace* is the set of vectors that get mapped to $\mathbf{0}_m$, and is denoted by $N(A)$.

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶ $Ax = \mathbf{0}_m$ are also called **Homogeneous equations**.
- ▶ Properties:

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!
- ▶ **Nullspace:** For a matrix $A \in \mathbb{R}^{m \times n}$, the *Nullspace* is the set of vectors that get mapped to $\mathbf{0}_m$, and is denoted by $N(A)$.

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶ $Ax = \mathbf{0}_m$ are also called **Homogeneous equations**.
- ▶ Properties:
- ▶ $\mathbf{0}_n$

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!
- ▶ **Nullspace:** For a matrix $A \in \mathbb{R}^{m \times n}$, the *Nullspace* is the set of vectors that get mapped to $\mathbf{0}_m$, and is denoted by $N(A)$.

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶ $Ax = \mathbf{0}_m$ are also called **Homogeneous equations**.
- ▶ Properties:
- ▶ $\mathbf{0}_n \in N(A)$.

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!
- ▶ **Nullspace:** For a matrix $A \in \mathbb{R}^{m \times n}$, the *Nullspace* is the set of vectors that get mapped to $\mathbf{0}_m$, and is denoted by $N(A)$.

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶ $Ax = \mathbf{0}_m$ are also called **Homogeneous equations**.
- ▶ Properties:
- ▶ $\mathbf{0}_n \in N(A)$.
- ▶ If $x, y \in N(A)$, $\forall p, q \in \mathbb{R}, p \cdot x + q \cdot y$

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!
- ▶ **Nullspace:** For a matrix $A \in \mathbb{R}^{m \times n}$, the *Nullspace* is the set of vectors that get mapped to $\mathbf{0}_m$, and is denoted by $N(A)$.

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶ $Ax = \mathbf{0}_m$ are also called **Homogeneous equations**.
- ▶ Properties:
- ▶ $\mathbf{0}_n \in N(A)$.
- ▶ If $x, y \in N(A)$, $\forall p, q \in \mathbb{R}, p \cdot x + q \cdot y \in N(A)$.

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!
- ▶ **Nullspace:** For a matrix $A \in \mathbb{R}^{m \times n}$, the *Nullspace* is the set of vectors that get mapped to $\mathbf{0}_m$, and is denoted by $N(A)$.

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶ $Ax = \mathbf{0}_m$ are also called **Homogeneous equations**.
- ▶ Properties:
- ▶ $\mathbf{0}_n \in N(A)$.
- ▶ If $x, y \in N(A)$, $\forall p, q \in \mathbb{R}$, $p \cdot x + q \cdot y \in N(A)$.
- ▶ If $\exists z \in N(A)$, $z \neq \mathbf{0}_n$, then $Ax = b$ will have

What about multiple solutions?

- ▶ Let $Ax = b$ and $Ay = b$, with $x \neq y$.
- ▶ Then, $A(x - y) = \mathbf{0}_m$.
- ▶ Similarly, let $z \in \mathbb{R}^n$ be such that $Az = \mathbf{0}_m$. Then, if $Ax = b$, $A(x + z) = b \Rightarrow$ Multiple Solutions!
- ▶ **Nullspace:** For a matrix $A \in \mathbb{R}^{m \times n}$, the *Nullspace* is the set of vectors that get mapped to $\mathbf{0}_m$, and is denoted by $N(A)$.

$$N(A) := \{x \in \mathbb{R}^n \mid Ax = \mathbf{0}_m\}, N(A) \subseteq \mathbb{R}^n$$

- ▶ $Ax = \mathbf{0}_m$ are also called **Homogeneous equations**.
- ▶ Properties:
- ▶ $\mathbf{0}_n \in N(A)$.
- ▶ If $x, y \in N(A)$, $\forall p, q \in \mathbb{R}$, $p \cdot x + q \cdot y \in N(A)$.
- ▶ If $\exists z \in N(A)$, $z \neq \mathbf{0}_n$, then $Ax = b$ will have infinitely many solutions, if one exists!

- If there are multiple solutions to $Ax = b \Rightarrow \exists z \neq \mathbf{0}_n$ such that $Az = \mathbf{0}_m$.

- If there are multiple solutions to $Ax = b \Rightarrow \exists z \neq \mathbf{0}_n$ such that $Az = \mathbf{0}_m$.
- Re-writing, we get $\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m$, with at least one non-zero entry in z , say z_k .

- If there are multiple solutions to $Ax = b \Rightarrow \exists z \neq \mathbf{0}_n$ such that $Az = \mathbf{0}_m$.
- Re-writing, we get $\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m$, with at least one non-zero entry in z , say z_k .
- $a_{*k} = \sum_{i=1, i \neq k}^n \frac{z_i}{z_k} a_{*i}$. Thus we can write column k as a *linear combination* of other columns.

- If there are multiple solutions to $Ax = b \Rightarrow \exists z \neq \mathbf{0}_n$ such that $Az = \mathbf{0}_m$.
- Re-writing, we get $\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m$, with at least one non-zero entry in z , say z_k .
- $a_{*k} = \sum_{i=1, i \neq k}^n \frac{z_i}{z_k} a_{*i}$. Thus we can write column k as a *linear combination* of other columns. We say that the vectors $\{a_{*i}, i = 1, \dots, n\}$ is a **linearly dependent** set of vectors.

- If there are multiple solutions to $Ax = b \Rightarrow \exists z \neq \mathbf{0}_n$ such that $Az = \mathbf{0}_m$.
- Re-writing, we get $\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m$, with at least one non-zero entry in z , say z_k .
- $a_{*k} = \sum_{i=1, i \neq k}^n \frac{z_i}{z_k} a_{*i}$. Thus we can write column k as a *linear combination* of other columns. We say that the vectors $\{a_{*i}, i = 1, \dots, n\}$ is a **linearly dependent** set of vectors.
- **Linear Independent set of vectors:** A set of vectors $\{a_{*i}, i = 1, \dots, n\}$ is said to be linearly independent if none of the vectors can be written as a linear combination of the remaining vectors.

- If there are multiple solutions to $Ax = b \Rightarrow \exists z \neq \mathbf{0}_n$ such that $Az = \mathbf{0}_m$.
- Re-writing, we get $\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m$, with at least one non-zero entry in z , say z_k .
- $a_{*k} = \sum_{i=1, i \neq k}^n \frac{z_i}{z_k} a_{*i}$. Thus we can write column k as a *linear combination* of other columns. We say that the vectors $\{a_{*i}, i = 1, \dots, n\}$ is a **linearly dependent** set of vectors.
- **Linear Independent set of vectors:** A set of vectors $\{a_{*i}, i = 1, \dots, n\}$ is said to be linearly independent if none of the vectors can be written as a linear combination of the remaining vectors.
- A set of vectors $\{a_{*i}, i = 1, \dots, n\}$ is said to be linearly independent if

$$\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m \Rightarrow$$

- If there are multiple solutions to $Ax = b \Rightarrow \exists z \neq \mathbf{0}_n$ such that $Az = \mathbf{0}_m$.
- Re-writing, we get $\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m$, with at least one non-zero entry in z , say z_k .
- $a_{*k} = \sum_{i=1, i \neq k}^n \frac{z_i}{z_k} a_{*i}$. Thus we can write column k as a *linear combination* of other columns. We say that the vectors $\{a_{*i}, i = 1, \dots, n\}$ is a **linearly dependent** set of vectors.
- **Linear Independent set of vectors:** A set of vectors $\{a_{*i}, i = 1, \dots, n\}$ is said to be linearly independent if none of the vectors can be written as a linear combination of the remaining vectors.
- A set of vectors $\{a_{*i}, i = 1, \dots, n\}$ is said to be linearly independent if

$$\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m \Rightarrow z_1 = z_2 = \dots = z_n = 0$$

- If there are multiple solutions to $Ax = b \Rightarrow \exists z \neq \mathbf{0}_n$ such that $Az = \mathbf{0}_m$.
- Re-writing, we get $\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m$, with at least one non-zero entry in z , say z_k .
- $a_{*k} = \sum_{i=1, i \neq k}^n \frac{z_i}{z_k} a_{*i}$. Thus we can write column k as a *linear combination* of other columns. We say that the vectors $\{a_{*i}, i = 1, \dots, n\}$ is a **linearly dependent** set of vectors.
- **Linear Independent set of vectors:** A set of vectors $\{a_{*i}, i = 1, \dots, n\}$ is said to be linearly independent if none of the vectors can be written as a linear combination of the remaining vectors.
- A set of vectors $\{a_{*i}, i = 1, \dots, n\}$ is said to be linearly independent if

$$\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m \Rightarrow z_1 = z_2 = \dots = z_n = 0$$

- **Column Rank of a Matrix:** The number of linearly independent columns of a matrix is called the Column Rank.

- If there are multiple solutions to $Ax = b \Rightarrow \exists z \neq \mathbf{0}_n$ such that $Az = \mathbf{0}_m$.
- Re-writing, we get $\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m$, with at least one non-zero entry in z , say z_k .
- $a_{*k} = \sum_{i=1, i \neq k}^n \frac{z_i}{z_k} a_{*i}$. Thus we can write column k as a *linear combination* of other columns. We say that the vectors $\{a_{*i}, i = 1, \dots, n\}$ is a **linearly dependent** set of vectors.
- **Linear Independent set of vectors:** A set of vectors $\{a_{*i}, i = 1, \dots, n\}$ is said to be linearly independent if none of the vectors can be written as a linear combination of the remaining vectors.
- A set of vectors $\{a_{*i}, i = 1, \dots, n\}$ is said to be linearly independent if

$$\sum_{i=1}^n z_i a_{*i} = \mathbf{0}_m \Rightarrow z_1 = z_2 = \dots = z_n = 0$$

- **Column Rank of a Matrix:** The number of linearly independent columns of a matrix is called the Column Rank.

- **Matrix Transpose:** For $A \in \mathbb{R}^{m \times n}$ given by

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix},$$

the matrix

$$\begin{bmatrix} a_{11} & a_{21} & \dots & a_{m1} \\ a_{12} & a_{22} & \dots & a_{m2} \\ \vdots & \vdots & \vdots & \vdots \\ \underbrace{a_{1n}}_{a_{1*}} & \underbrace{a_{2n}}_{a_{2*}} & \dots & \underbrace{a_{mn}}_{a_{m*}} \end{bmatrix} \in \mathbb{R}^{n \times m},$$

is called the *transpose of A*, and is denoted by A^T .

- **Matrix Transpose:** For $A \in \mathbb{R}^{m \times n}$ given by

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix},$$

the matrix

$$\begin{bmatrix} a_{11} & a_{21} & \dots & a_{m1} \\ a_{12} & a_{22} & \dots & a_{m2} \\ \vdots & \vdots & \vdots & \vdots \\ \underbrace{a_{1n}}_{a_{1*}} & \underbrace{a_{2n}}_{a_{2*}} & \dots & \underbrace{a_{mn}}_{a_{m*}} \end{bmatrix} \in \mathbb{R}^{n \times m},$$

is called the *transpose of A*, and is denoted by A^T .

- **Beware of the notation:** a_{i*} denotes the i^{th} row of A written as a column matrix.

- All linear combination of columns gives the column space, how about linear combination of rows?

- All linear combination of columns gives the column space, how about linear combination of rows?
- **Rowspace:** The set of all linear combinations of rows of the matrix is called the Rowspace.

- All linear combination of columns gives the column space, how about linear combination of rows?
- **Rowspace:** The set of all linear combinations of rows of the matrix is called the Row space.
- Since rows of A are columns of A^T , notation for row space is $C(A^T)$, and can be written as

$$C(A^T) = \{A^T y \mid \forall y \in \mathbb{R}^m\} \subseteq \mathbb{R}^n$$

- All linear combination of columns gives the column space, how about linear combination of rows?
- **Rowspace:** The set of all linear combinations of rows of the matrix is called the Row space.
- Since rows of A are columns of A^T , notation for row space is $C(A^T)$, and can be written as

$$C(A^T) = \{A^T y \mid \forall y \in \mathbb{R}^m\} \subseteq \mathbb{R}^n$$

- Properties:
 - ▶ $\mathbf{0}_n \in C(A^T)$
 - ▶ If $x_1, x_2 \in C(A^T)$, then $\forall p, q \in \mathbb{R}, p \cdot x_1 + q \cdot x_2 \in C(A^T)$.

- All linear combination of columns gives the column space, how about linear combination of rows?
- **Rowspace:** The set of all linear combinations of rows of the matrix is called the Row space.
- Since rows of A are columns of A^T , notation for row space is $C(A^T)$, and can be written as

$$C(A^T) = \{A^T y \mid \forall y \in \mathbb{R}^m\} \subseteq \mathbb{R}^n$$

- Properties:
 - ▶ $\mathbf{0}_n \in C(A^T)$
 - ▶ If $x_1, x_2 \in C(A^T)$, then $\forall p, q \in \mathbb{R}, p \cdot x_1 + q \cdot x_2 \in C(A^T)$.
- **Row Rank of a Matrix:** The number of linearly independent rows of a matrix is called the Column Rank.

- **Left Nullspace:** The set of solutions in \mathbb{R}^m to $A^T y = \mathbf{0}_n$ is called the *Left Nullspace* of matrix A .

- **Left Nullspace:** The set of solutions in \mathbb{R}^m to $A^T y = \mathbf{0}_n$ is called the *Left Nullspace* of matrix A .
- Since this is the same as Nullspace of the matrix A^T , left nullspace is denoted by $N(A^T)$.

$$N(A^T) := \{y \in \mathbb{R}^m \mid A^T y = \mathbf{0}_n\} \subseteq \mathbb{R}^m$$

- Properties:
 - ▶ $\mathbf{0}_m \in N(A^T)$
 - ▶ If $y_1, y_2 \in N(A^T)$, then $\forall p, q \in \mathbb{R}, p \cdot y_1 + q \cdot y_2 \in N(A^T)$.

- What is $C(A^T) \cap N(A)$?

- What is $C(A^T) \cap N(A)$?
- Let $x \in C(A^T) \cap N(A)$. Since $x \in C(A^T)$, $\exists y \in \mathbb{R}^m$ such that $x = A^T y$, and since $x \in N(A)$, $Ax = \mathbf{0}_m$.

- What is $C(A^T) \cap N(A)$?
- Let $x \in C(A^T) \cap N(A)$. Since $x \in C(A^T)$, $\exists y \in \mathbb{R}^m$ such that $x = A^T y$, and since $x \in N(A)$, $Ax = \mathbf{0}_m$.
- Thus,

$$Ax = AA^T y = \mathbf{0}_m$$

- What is $C(A^T) \cap N(A)$?
- Let $x \in C(A^T) \cap N(A)$. Since $x \in C(A^T)$, $\exists y \in \mathbb{R}^m$ such that $x = A^T y$, and since $x \in N(A)$, $Ax = \mathbf{0}_m$.
- Thus,

$$Ax = AA^T y = \mathbf{0}_m$$

$$y^T \underbrace{AA^T}_{v \in \mathbb{R}^n} y = 0$$

- What is $C(A^T) \cap N(A)$?
- Let $x \in C(A^T) \cap N(A)$. Since $x \in C(A^T)$, $\exists y \in \mathbb{R}^m$ such that $x = A^T y$, and since $x \in N(A)$, $Ax = \mathbf{0}_m$.
- Thus,

$$Ax = AA^T y = \mathbf{0}_m$$

$$y^T \underbrace{AA^T}_{v \in \mathbb{R}^n} y = 0$$

$$\Rightarrow v^T v = 0 \Rightarrow v = A^T y = x = \mathbf{0}_n.$$

Thus $C(A^T) \cap N(A) = \mathbf{0}_n$.

- What is $C(A^T) \cap N(A)$?
- Let $x \in C(A^T) \cap N(A)$. Since $x \in C(A^T)$, $\exists y \in \mathbb{R}^m$ such that $x = A^T y$, and since $x \in N(A)$, $Ax = \mathbf{0}_m$.
- Thus,

$$Ax = AA^T y = \mathbf{0}_m$$

$$y^T \underbrace{AA^T}_{v \in \mathbb{R}^n} y = 0$$

$$\Rightarrow v^T v = 0 \Rightarrow v = A^T y = x = \mathbf{0}_n.$$

Thus $C(A^T) \cap N(A) = \mathbf{0}_n$.

- Similarly, $C(A) \cap N(A^T) = \mathbf{0}_m$.

- Are Column and Row rank of a matrix different?

- Are Column and Row rank of a matrix different?
- Let $A \in \mathbb{R}^{m \times n}$ be a matrix with column rank r . wlog, assume first r columns are LI.

- Are Column and Row rank of a matrix different?
- Let $A \in \mathbb{R}^{m \times n}$ be a matrix with column rank r . wlog, assume first r columns are LI.

$$\begin{aligned}
 & \begin{bmatrix} a_{*1} & \dots & a_{*r} & a_{*r+1} & \dots & a_{*n} \end{bmatrix}_{m \times n} \\
 &= \begin{bmatrix} a_{*1} & \dots & a_{*r} \end{bmatrix}_{m \times r} \underbrace{\begin{bmatrix} I_{r \times r} & b_{*r+1} & \dots & b_{*n} \end{bmatrix}_{r \times n}}_C \\
 & \begin{bmatrix} a_{1*}^T \\ a_{2*}^T \\ \vdots \\ a_{m*}^T \end{bmatrix} = \begin{bmatrix} a_{*1} & \dots & a_{*r} \end{bmatrix}_{m \times r} \begin{bmatrix} c_{1*}^T \\ c_{2*}^T \\ \vdots \\ c_{r*}^T \end{bmatrix}_{r \times n}
 \end{aligned}$$