

SC223 - Linear Algebra

Aditya Tatu

Lecture 12



September 6, 2022

Vector Spaces

● **Definition:** A Vector space is a set V with a **field** $(\mathbb{F}, +_F, \times)$, and two binary operations, vector addition $+$ and scalar multiplication \cdot that satisfy the following axioms:

► $(V, +)$ is an **Abelian group**:

► $\forall x, y \in V, x + y \in V$

► $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$

► $\forall x \in V, \exists y \in V, x + y = y + x = \theta$. We will denote y by

$-x$.

► $\forall x, y, z \in V, (x + y) + z = x + (y + z)$.

► $\forall x, y \in V, x + y = y + x$.

Example

- Symmetry group of a rectangle.
 - What all can we do to the rectangle so that we get the same (not necessarily same vertices) rectangle?




Figure: Symmetries of a rectangle

Example

- Symmetry group of a rectangle.
 - What all can we do to the rectangle so that we get the same (not necessarily same vertices) rectangle?




Figure: Symmetries of a rectangle

- Do nothing! - Identity: I

Example

- Symmetry group of a rectangle.
 - ▶ What all can we do to the rectangle so that we get the same (not necessarily same vertices) rectangle?




Figure: Symmetries of a rectangle

- ▶ Do nothing! - Identity: I
- ▶ Rotate by 180 deg: R_{180}

Example

- Symmetry group of a rectangle.
 - ▶ What all can we do to the rectangle so that we get the same (not necessarily same vertices) rectangle?




Figure: Symmetries of a rectangle

- ▶ Do nothing! - Identity: I
- ▶ Rotate by 180 deg: R_{180}
- ▶ Flip along vertical axis: F_v

Example

- Symmetry group of a rectangle.
 - What all can we do to the rectangle so that we get the same (not necessarily same vertices) rectangle?




Figure: Symmetries of a rectangle

- Do nothing! - Identity: I
- Rotate by 180 deg: R_{180}
- Flip along vertical axis: F_v
- Flip along horizontal axis: F_h .

Example

- Observe the set $D_2 = \{I, R_{180}, F_h, F_v\}$ along with composition \circ operation:

Example

- Observe the set $D_2 = \{I, R_{180}, F_h, F_v\}$ along with composition \circ operation:

| \circ | I | R_{180} | F_h | F_v |
|-----------|-----------|-----------|-----------|-----------|
| I | I | R_{180} | F_h | F_v |
| R_{180} | R_{180} | I | F_v | F_h |
| F_h | F_h | F_v | I | R_{180} |
| F_v | F_v | F_h | R_{180} | I |

Example

- Observe the set $D_2 = \{I, R_{180}, F_h, F_v\}$ along with composition \circ operation:

| \circ | I | R_{180} | F_h | F_v |
|-----------|-----------|-----------|-----------|-----------|
| I | I | R_{180} | F_h | F_v |
| R_{180} | R_{180} | I | F_v | F_h |
| F_h | F_h | F_v | I | R_{180} |
| F_v | F_v | F_h | R_{180} | I |

- (S, \circ) is closed.

Example

- Observe the set $D_2 = \{I, R_{180}, F_h, F_v\}$ along with composition \circ operation:

| \circ | I | R_{180} | F_h | F_v |
|-----------|-----------|-----------|-----------|-----------|
| I | I | R_{180} | F_h | F_v |
| R_{180} | R_{180} | I | F_v | F_h |
| F_h | F_h | F_v | I | R_{180} |
| F_v | F_v | F_h | R_{180} | I |

- ▶ (S, \circ) is closed.
- ▶ I is the identity wrt \circ

Example

- Observe the set $D_2 = \{I, R_{180}, F_h, F_v\}$ along with composition \circ operation:

| \circ | I | R_{180} | F_h | F_v |
|-----------|-----------|-----------|-----------|-----------|
| I | I | R_{180} | F_h | F_v |
| R_{180} | R_{180} | I | F_v | F_h |
| F_h | F_h | F_v | I | R_{180} |
| F_v | F_v | F_h | R_{180} | I |

- ▶ (S, \circ) is closed.
- ▶ I is the identity wrt \circ
- ▶ $-I = I, -R_{180} = R_{180}, -F_h = F_h, -F_v = F_v$

Example

- Observe the set $D_2 = \{I, R_{180}, F_h, F_v\}$ along with composition \circ operation:

| \circ | I | R_{180} | F_h | F_v |
|-----------|-----------|-----------|-----------|-----------|
| I | I | R_{180} | F_h | F_v |
| R_{180} | R_{180} | I | F_v | F_h |
| F_h | F_h | F_v | I | R_{180} |
| F_v | F_v | F_h | R_{180} | I |

- ▶ (S, \circ) is closed.
- ▶ I is the identity wrt \circ
- ▶ $-I = I, -R_{180} = R_{180}, -F_h = F_h, -F_v = F_v$
- ▶ \circ is associative.

Example

- Observe the set $D_2 = \{I, R_{180}, F_h, F_v\}$ along with composition \circ operation:

| \circ | I | R_{180} | F_h | F_v |
|-----------|-----------|-----------|-----------|-----------|
| I | I | R_{180} | F_h | F_v |
| R_{180} | R_{180} | I | F_v | F_h |
| F_h | F_h | F_v | I | R_{180} |
| F_v | F_v | F_h | R_{180} | I |

- ▶ (S, \circ) is closed.
- ▶ I is the identity wrt \circ
- ▶ $-I = I, -R_{180} = R_{180}, -F_h = F_h, -F_v = F_v$
- ▶ \circ is associative.
- ▶ \circ is commutative. Thus (S, \circ) forms an Abelian group.

- Consider the set $S' = \{00, 01, 10, 11\}$ with bitwise addition modulo-2 operation: $+_2$.

| $+_2$ | 00 | 01 | 10 | 11 |
|-------|----|----|----|----|
| 00 | 00 | 01 | 10 | 11 |
| 01 | 01 | 00 | 11 | 10 |
| 10 | 10 | 11 | 00 | 01 |
| 11 | 11 | 10 | 01 | 00 |

- Consider the set $S' = \{00, 01, 10, 11\}$ with bitwise addition modulo-2 operation: $+_2$.

| $+_2$ | 00 | 01 | 10 | 11 |
|-------|----|----|----|----|
| 00 | 00 | 01 | 10 | 11 |
| 01 | 01 | 00 | 11 | 10 |
| 10 | 10 | 11 | 00 | 01 |
| 11 | 11 | 10 | 01 | 00 |

- Compare with:

| \circ | I | R_{180} | F_h | F_v |
|-----------|-----------|-----------|-----------|-----------|
| I | I | R_{180} | F_h | F_v |
| R_{180} | R_{180} | I | F_v | F_h |
| F_h | F_h | F_v | I | R_{180} |
| F_v | F_v | F_h | R_{180} | I |

- Consider the set $S' = \{00, 01, 10, 11\}$ with bitwise addition modulo-2 operation: $+_2$.

| $+_2$ | 00 | 01 | 10 | 11 |
|-------|----|----|----|----|
| 00 | 00 | 01 | 10 | 11 |
| 01 | 01 | 00 | 11 | 10 |
| 10 | 10 | 11 | 00 | 01 |
| 11 | 11 | 10 | 01 | 00 |

- Compare with:

| \circ | I | R_{180} | F_h | F_v |
|-----------|-----------|-----------|-----------|-----------|
| I | I | R_{180} | F_h | F_v |
| R_{180} | R_{180} | I | F_v | F_h |
| F_h | F_h | F_v | I | R_{180} |
| F_v | F_v | F_h | R_{180} | I |

- Both are examples of the *Klein-4* group.

Vector Spaces

● **Definition:** A Vector space is a set V with a **field** $(\mathbb{F}, +_F, \times)$, and two binary operations, vector addition $+$ and scalar multiplication \cdot that satisfy the following axioms:

► $(V, +)$ is an **Abelian group**:

► $\forall x, y \in V, x + y \in V$

► $\exists \theta \in V, \forall x \in V, x + \theta = \theta + x = x$

► $\forall x \in V, \exists y \in V, x + y = y + x = \theta$. We will denote y by

$-x$.

► $\forall x, y, z \in V, (x + y) + z = x + (y + z)$.

► $\forall x, y \in V, x + y = y + x$.

- **Closure with respect to Scalar multiplication:** $\cdot \mathbb{F} \times V \rightarrow V$.

- **Closure with respect to Scalar multiplication:** $\cdot: \mathbb{F} \times V \rightarrow V$.
- **Scalar Multiplication identity:** $\exists 1 \in \mathbb{F}$ such that $1 \cdot v = v, \forall v \in V$.

- **Closure with respect to Scalar multiplication:** $\cdot: \mathbb{F} \times V \rightarrow V$.
- **Scalar Multiplication identity:** $\exists 1 \in \mathbb{F}$ such that $1 \cdot v = v, \forall v \in V$.
- **Distributivity:** $\forall a \in \mathbb{F}, \forall u, v \in V, a \cdot (u + v) = a \cdot u + a \cdot v$,
and $\forall a, b \in \mathbb{F}, \forall u \in V, (a +_F b) \cdot u = a \cdot u + b \cdot u$.

- **Closure with respect to Scalar multiplication:** $\cdot \mathbb{F} \times V \rightarrow V$.
- **Scalar Multiplication identity:** $\exists 1 \in \mathbb{F}$ such that $1 \cdot v = v, \forall v \in V$.
- **Distributivity:** $\forall a \in \mathbb{F}, \forall u, v \in V, a \cdot (u + v) = a \cdot u + a \cdot v$,
and $\forall a, b \in \mathbb{F}, \forall u \in V, (a +_F b) \cdot u = a \cdot u + b \cdot u$.
- **Compatibility of field and scalar multiplication:**
 $\forall a, b \in \mathbb{F}, \forall u \in V, (a \times b) \cdot u = a \cdot (b \cdot u)$.

Field

- **Definition:**(Field). A field is a set \mathbb{F} with two binary operations, addition $+_{\mathbb{F}}$ and multiplication \times that satisfy the following axioms:

Field

- **Definition:**(Field). A field is a set \mathbb{F} with two binary operations, addition $+_F$ and multiplication \times that satisfy the following axioms:
 - ▶ $(\mathbb{F}, +_F)$ is an **Abelian group**. The additive identity will be denoted by 0.

Field

- **Definition:**(Field). A field is a set \mathbb{F} with two binary operations, addition $+_F$ and multiplication \times that satisfy the following axioms:
 - ▶ $(\mathbb{F}, +_F)$ is an **Abelian group**. The additive identity will be denoted by 0.
 - ▶ $(\mathbb{F} - \{0\}, \times)$ is an **Abelian group**. The multiplicative identity will be denoted by 1.

Field

● **Definition:**(Field). A field is a set \mathbb{F} with two binary operations, addition $+_F$ and multiplication \times that satisfy the following axioms:

► $(\mathbb{F}, +_F)$ is an **Abelian group**. The additive identity will be denoted by 0.

► $(\mathbb{F} - \{0\}, \times)$ is an **Abelian group**. The multiplicative identity will be denoted by 1.

► **Distributivity:**

$$\forall a, b, c \in \mathbb{F}, (a +_F b) \times c = a \times c +_F b \times c, a \times (b +_F c) = a \times b +_F a \times c$$

Examples of Fields

► $(\mathbb{Z}_2, +_2, \times)$

Examples of Fields

- ▶ $(\mathbb{Z}_2, +_2, \times)$
- ▶ $(\mathbb{R}, +, \times)$

Examples of Fields

- ▶ $(\mathbb{Z}_2, +_2, \times)$
- ▶ $(\mathbb{R}, +, \times)$
- ▶ $(\mathbb{C}, +, \times)$

Examples of Fields

- ▶ $(\mathbb{Z}_2, +_2, \times)$
- ▶ $(\mathbb{R}, +, \times)$
- ▶ $(\mathbb{C}, +, \times)$
- ▶ $(\mathbb{Q}, +, \times)$

Examples of Fields

- ▶ $(\mathbb{Z}_2, +_2, \times)$
- ▶ $(\mathbb{R}, +, \times)$
- ▶ $(\mathbb{C}, +, \times)$
- ▶ $(\mathbb{Q}, +, \times)$
- ▶ $(\mathbb{R}[x], +, \times)$, where $\mathbb{R}[x]$ is the set of all rational polynomials of the form $\frac{p(x)}{q(x)}$, with $q \neq 0$, and p and q are polynomials in one variable with real coefficients.

Examples of Fields

- ▶ $(\mathbb{Z}_2, +_2, \times)$
- ▶ $(\mathbb{R}, +, \times)$
- ▶ $(\mathbb{C}, +, \times)$
- ▶ $(\mathbb{Q}, +, \times)$
- ▶ $(\mathbb{R}[x], +, \times)$, where $\mathbb{R}[x]$ is the set of all rational polynomials of the form $\frac{p(x)}{q(x)}$, with $q \neq 0$, and p and q are polynomials in one variable with real coefficients.
- ▶ If the 3-tuple $(V, +, \cdot)$ with field $(\mathbb{F}, +_F, \times)$ satisfies all vector space axioms, we say that $(V, +, \cdot)$ forms a vector space over \mathbb{F} .

Examples of Fields

- ▶ $(\mathbb{Z}_2, +_2, \times)$
- ▶ $(\mathbb{R}, +, \times)$
- ▶ $(\mathbb{C}, +, \times)$
- ▶ $(\mathbb{Q}, +, \times)$
- ▶ $(\mathbb{R}[x], +, \times)$, where $\mathbb{R}[x]$ is the set of all rational polynomials of the form $\frac{p(x)}{q(x)}$, with $q \neq 0$, and p and q are polynomials in one variable with real coefficients.
- ▶ If the 3-tuple $(V, +, \cdot)$ with field $(\mathbb{F}, +_F, \times)$ satisfies all vector space axioms, we say that $(V, +, \cdot)$ forms a vector space over \mathbb{F} .
- Any element of the vector space $(V, +, \cdot)$ will be referred to as a **vector**, and any element $a \in \mathbb{F}$ will be referred to as a **scalar**.

Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$ over \mathbb{R} .

Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$ over \mathbb{R} .
- $(\mathbb{R}^n, +, \cdot)$ over \mathbb{R} .

Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$ over \mathbb{R} .
- $(\mathbb{R}^n, +, \cdot)$ over \mathbb{R} .
- $(\mathbb{C}^n, +, \cdot)$ over \mathbb{C} .

Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$ over \mathbb{R} .
- $(\mathbb{R}^n, +, \cdot)$ over \mathbb{R} .
- $(\mathbb{C}^n, +, \cdot)$ over \mathbb{C} .
- $(\mathbb{R}^{\mathbb{Z}}, +, \cdot)$ over \mathbb{R} , where $\mathbb{R}^{\mathbb{Z}}$ is the set of all doubly-infinite sequences.

Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$ over \mathbb{R} .
- $(\mathbb{R}^n, +, \cdot)$ over \mathbb{R} .
- $(\mathbb{C}^n, +, \cdot)$ over \mathbb{C} .
- $(\mathbb{R}^{\mathbb{Z}}, +, \cdot)$ over \mathbb{R} , where $\mathbb{R}^{\mathbb{Z}}$ is the set of all doubly-infinite sequences.
- $(\mathcal{P}(\mathbb{R}), +, \cdot)$ over \mathbb{R} , where $\mathcal{P}(\mathbb{R})$ is the set of all polynomials of one variable with real coefficients.

Examples of Vector spaces

- $(\mathbb{R}, +, \cdot)$ over \mathbb{R} .
- $(\mathbb{R}^n, +, \cdot)$ over \mathbb{R} .
- $(\mathbb{C}^n, +, \cdot)$ over \mathbb{C} .
- $(\mathbb{R}^{\mathbb{Z}}, +, \cdot)$ over \mathbb{R} , where $\mathbb{R}^{\mathbb{Z}}$ is the set of all doubly-infinite sequences.
- $(\mathcal{P}(\mathbb{R}), +, \cdot)$ over \mathbb{R} , where $\mathcal{P}(\mathbb{R})$ is the set of all polynomials of one variable with real coefficients.
- $(\mathbb{L}_2(\mathbb{R}), +, \cdot)$ over \mathbb{R} , where $\mathbb{L}_2(\mathbb{R})$ denotes the set of all square-integrable functions $f : \mathbb{R} \rightarrow \mathbb{R}$.