

# SC223 - Linear Algebra

Aditya Tatu

Lecture 8



August 26, 2022

# LU Decomposition

- Any matrix  $A \in \mathbb{R}^{m \times n}$  can be decomposed into a product of lower and upper triangular matrices, with appropriate permutations:

$$PA = LU,$$

where  $P \in \mathbb{R}^{m \times m}$ ,  $L \in \mathbb{R}^{m \times m}$ ,  $U \in \mathbb{R}^{m \times n}$ .

- Solving linear equations with  $LU$  decomposition:

$$Ax = b \rightarrow LUx = b.$$

- First let  $Ux = y$  and solve  $Ly = b$ , and next solve for  $x$  in  $Ux = y$ .

# LU Decomposition

- Any matrix  $A \in \mathbb{R}^{m \times n}$  can be decomposed into a product of lower and upper triangular matrices, with appropriate permutations:

$$PA = LU,$$

where  $P \in \mathbb{R}^{m \times m}$ ,  $L \in \mathbb{R}^{m \times m}$ ,  $U \in \mathbb{R}^{m \times n}$ .

- Solving linear equations with  $LU$  decomposition:

$$Ax = b \rightarrow LUx = b.$$

- First let  $Ux = y$  and solve  $Ly = b$ , and next solve for  $x$  in  $Ux = y$ .

- In the example, where is  $y$ ?

$$\left[ \begin{array}{ccccc|c} \mathbf{1} & -1 & 2 & 3 & -1 & 1 \\ 0 & \mathbf{2} & 5 & 4 & 2 & 2 \\ 0 & 0 & -\mathbf{23} & -16 & 2 & -12 \\ 0 & 0 & 0 & -\mathbf{137} & -95 & 64 \end{array} \right]$$

## Example

$$\bullet \quad U = \begin{bmatrix} \mathbf{1} & -1 & 2 & 3 & -1 \\ 0 & \mathbf{2} & 5 & 4 & 2 \\ 0 & 0 & -\mathbf{23} & -16 & 2 \\ 0 & 0 & 0 & -\mathbf{137} & -95 \end{bmatrix}, y = \begin{bmatrix} 1 \\ 2 \\ -12 \\ 64 \end{bmatrix}.$$

## Example

$$\bullet U = \begin{bmatrix} \mathbf{1} & -1 & 2 & 3 & -1 \\ 0 & \mathbf{2} & 5 & 4 & 2 \\ 0 & 0 & -\mathbf{23} & -16 & 2 \\ 0 & 0 & 0 & -\mathbf{137} & -95 \end{bmatrix}, y = \begin{bmatrix} 1 \\ 2 \\ -12 \\ 64 \end{bmatrix}.$$

● Solve for  $x$  in

$$\begin{bmatrix} \mathbf{1} & -1 & 2 & 3 & -1 \\ 0 & \mathbf{2} & 5 & 4 & 2 \\ 0 & 0 & -\mathbf{23} & -16 & 2 \\ 0 & 0 & 0 & -\mathbf{137} & -95 \end{bmatrix} x = \begin{bmatrix} 1 \\ 2 \\ -12 \\ 64 \end{bmatrix}$$

# LU Decomposition Algorithm

● Let  $A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \dots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$

► How to eliminate first variable from Eqns 2 to  $n$ ?

# LU Decomposition Algorithm

● Let  $A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \dots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$

► How to eliminate first variable from Eqns 2 to  $n$ ?

►  $L_1 = \begin{bmatrix} 1 & 0 & \dots & 0 \\ -\frac{a_{21}}{a_{11}} & 1 & \dots & 0 \\ -\frac{a_{31}}{a_{11}} & 0 & \dots & 0 \\ \vdots & \vdots & \dots & \vdots \\ -\frac{a_{n1}}{a_{11}} & 0 & \dots & 1 \end{bmatrix}$

# LU Decomposition Algorithm

● Let  $A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \dots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$

► How to eliminate first variable from Eqns 2 to  $n$ ?

►  $L_1 = \begin{bmatrix} 1 & 0 & \dots & 0 \\ -\frac{a_{21}}{a_{11}} & 1 & \dots & 0 \\ -\frac{a_{31}}{a_{11}} & 0 & \dots & 0 \\ \vdots & \vdots & \dots & \vdots \\ -\frac{a_{n1}}{a_{11}} & 0 & \dots & 1 \end{bmatrix}$

► Let  $L_1 A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ 0 & a_{22}^1 & \dots & a_{2n}^1 \\ 0 & a_{32}^1 & \dots & a_{3n}^1 \\ \vdots & \vdots & \dots & \vdots \\ 0 & a_{n2}^1 & \dots & a_{nn}^1 \end{bmatrix}$



- How to eliminate variable 2 from Eqns 3 to  $n$ ?

► How to eliminate variable 2 from Eqns 3 to  $n$ ?

$$\text{► } L_2 = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & -\frac{a_{32}^1}{a_{22}^1} & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & -\frac{a_{n2}^1}{a_{22}^1} & 0 & \dots & 1 \end{bmatrix}$$

► How to eliminate variable 2 from Eqns 3 to  $n$ ?

►  $L_2 = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & -\frac{a_{32}^1}{a_{22}^1} & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & -\frac{a_{n2}^1}{a_{22}^1} & 0 & \dots & 1 \end{bmatrix}, L_2 L_1 A =$

► How to eliminate variable 2 from Eqns 3 to  $n$ ?

$$\text{► } L_2 = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & -\frac{a_{32}^1}{a_{22}^1} & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & -\frac{a_{n2}^1}{a_{22}^1} & 0 & \dots & 1 \end{bmatrix}, L_2 L_1 A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ 0 & a_{22}^1 & a_{23}^1 & \dots & a_{2n}^1 \\ 0 & 0 & a_{33}^2 & \dots & a_{3n}^2 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & a_{n3}^2 & \dots & a_{nn}^2 \end{bmatrix}$$

- How to eliminate variable 2 from Eqns 3 to  $n$ ?

$$\text{► } L_2 = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & -\frac{a_{32}^1}{a_{22}^1} & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & -\frac{a_{n2}^1}{a_{22}^1} & 0 & \dots & 1 \end{bmatrix}, L_2 L_1 A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ 0 & a_{22}^1 & a_{23}^1 & \dots & a_{2n}^1 \\ 0 & 0 & a_{33}^2 & \dots & a_{3n}^2 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & a_{n3}^2 & \dots & a_{nn}^2 \end{bmatrix}$$

- Final Step:  $L_{n-1} \cdot \dots \cdot L_1 A = U$ .

- How to eliminate variable 2 from Eqns 3 to  $n$ ?

$$\text{► } L_2 = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & -\frac{a_{32}^1}{a_{22}^1} & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & -\frac{a_{n2}^1}{a_{22}^1} & 0 & \dots & 1 \end{bmatrix}, L_2 L_1 A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ 0 & a_{22}^1 & a_{23}^1 & \dots & a_{2n}^1 \\ 0 & 0 & a_{33}^2 & \dots & a_{3n}^2 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & a_{n3}^2 & \dots & a_{nn}^2 \end{bmatrix}$$

- Final Step:  $L_{n-1} \cdot \dots \cdot L_1 A = U$ .

- $L_{n-1} \cdot \dots \cdot L_1 \cdot L = I_n$ .

- How to eliminate variable 2 from Eqns 3 to  $n$ ?

$$\text{► } L_2 = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & -\frac{a_{32}^1}{a_{22}^1} & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & -\frac{a_{n2}^1}{a_{22}^1} & 0 & \dots & 1 \end{bmatrix}, L_2 L_1 A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ 0 & a_{22}^1 & a_{23}^1 & \dots & a_{2n}^1 \\ 0 & 0 & a_{33}^2 & \dots & a_{3n}^2 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & a_{n3}^2 & \dots & a_{nn}^2 \end{bmatrix}$$

- Final Step:  $L_{n-1} \cdot \dots \cdot L_1 A = U$ .

- $L_{n-1} \cdot \dots \cdot L_1 \cdot L = I_n$ .

- $L_{n-1} \cdot \dots \cdot L_1 \cdot [x_1 \ x_2 \ \dots \ x_n] = [I_{*1} \ I_{*2} \ \dots \ I_{*n}]$ .

- How to eliminate variable 2 from Eqns 3 to  $n$ ?

$$\text{► } L_2 = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & -\frac{a_{32}^1}{a_{22}^1} & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & -\frac{a_{n2}^1}{a_{22}^1} & 0 & \dots & 1 \end{bmatrix}, L_2 L_1 A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ 0 & a_{22}^1 & a_{23}^1 & \dots & a_{2n}^1 \\ 0 & 0 & a_{33}^2 & \dots & a_{3n}^2 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & a_{n3}^2 & \dots & a_{nn}^2 \end{bmatrix}$$

- Final Step:  $L_{n-1} \cdot \dots \cdot L_1 A = U$ .

$$\text{► } L_{n-1} \cdot \dots \cdot L_1 \cdot L = I_n.$$

$$\text{► } L_{n-1} \cdot \dots \cdot L_1 \cdot [x_1 \ x_2 \ \dots \ x_n] = [I_{*1} \ I_{*2} \ \dots \ I_{*n}].$$

$$\text{► If } L_1 x_1 = e_1, \text{ then } L_2 L_1 x_1 =$$



- How to eliminate variable 2 from Eqns 3 to  $n$ ?

$$\text{► } L_2 = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & -\frac{a_{32}^1}{a_{22}^1} & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & -\frac{a_{n2}^1}{a_{22}^1} & 0 & \dots & 1 \end{bmatrix}, L_2 L_1 A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ 0 & a_{22}^1 & a_{23}^1 & \dots & a_{2n}^1 \\ 0 & 0 & a_{33}^2 & \dots & a_{3n}^2 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & a_{n3}^2 & \dots & a_{nn}^2 \end{bmatrix}$$

- Final Step:  $L_{n-1} \cdot \dots \cdot L_1 A = U$ .

- $L_{n-1} \cdot \dots \cdot L_1 \cdot L = I_n$ .

- $L_{n-1} \cdot \dots \cdot L_1 \cdot [x_1 \ x_2 \ \dots \ x_n] = [I_{*1} \ I_{*2} \ \dots \ I_{*n}]$ .

- If  $L_1 x_1 = e_1$ , then  $L_2 L_1 x_1 = e_1$ .

- How to eliminate variable 2 from Eqns 3 to  $n$ ?

$$\text{► } L_2 = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & -\frac{a_{32}}{a_{22}} & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & -\frac{a_{n2}}{a_{22}} & 0 & \dots & 1 \end{bmatrix}, L_2 L_1 A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ 0 & a_{22}^1 & a_{23}^1 & \dots & a_{2n}^1 \\ 0 & 0 & a_{33}^2 & \dots & a_{3n}^2 \\ \vdots & \vdots & \vdots & \dots & \vdots \\ 0 & 0 & a_{n3}^2 & \dots & a_{nn}^2 \end{bmatrix}$$

- Final Step:  $L_{n-1} \cdot \dots \cdot L_1 A = U$ .

$$\text{► } L_{n-1} \cdot \dots \cdot L_1 \cdot L = I_n.$$

$$\text{► } L_{n-1} \cdot \dots \cdot L_1 \cdot [x_1 \ x_2 \ \dots \ x_n] = [I_{*1} \ I_{*2} \ \dots \ I_{*n}].$$

- If  $L_1 x_1 = e_1$ , then  $L_2 L_1 x_1 = e_1$ .

$$\text{► Thus, } L = \begin{bmatrix} 1 & 0 & \dots & 0 \\ \frac{a_{21}}{a_{11}} & 1 & \dots & 0 \\ \frac{a_{31}}{a_{11}} & \frac{a_{32}}{a_{22}} & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots \\ \frac{a_{n1}}{a_{11}} & \frac{a_{n2}}{a_{22}} & \dots & 1 \end{bmatrix}$$

# Computing Matrix Inverses

- $A^{-1}$  is the unique matrix such that  $A^{-1}A = AA^{-1} = I$ .

# Computing Matrix Inverses

- $A^{-1}$  is the unique matrix such that  $A^{-1}A = AA^{-1} = I$ .
- Simultaneously solve  $n$  linear equations:

$$\underbrace{\begin{bmatrix} a_{*1} & \dots & a_{*n} \\ a_{*1} & \dots & a_{*n} \end{bmatrix}}_{AM} \begin{bmatrix} x_1 & \dots & x_n \end{bmatrix} = I, x_i \in \mathbb{R}^n, i = 1, \dots, n$$
$$\begin{bmatrix} LU & | & I \end{bmatrix} \Rightarrow LUx_1 = I_{*1}, \dots, LUx_n = I_{*n}.$$

# Computing Matrix Inverses

- $A^{-1}$  is the unique matrix such that  $A^{-1}A = AA^{-1} = I$ .
- Simultaneously solve  $n$  linear equations:

$$\underbrace{\begin{bmatrix} a_{*1} & \dots & a_{*n} \end{bmatrix}}_{AM} \begin{bmatrix} x_1 & \dots & x_n \end{bmatrix} = I, x_i \in \mathbb{R}^n, i = 1, \dots, n$$
$$\begin{bmatrix} LU & | & I \end{bmatrix} \Rightarrow LUx_1 = I_{*1}, \dots, LUx_n = I_{*n}.$$

- Gauss-Jordan Method: Let  $R_1, \dots, R_k$  represent row transformation matrices, not necessarily lower triangular, such that  $R_k \cdot R_{k-1} \cdot \dots \cdot R_1 A = I$ , then  $A^{-1} =$

# Computing Matrix Inverses

- $A^{-1}$  is the unique matrix such that  $A^{-1}A = AA^{-1} = I$ .
- Simultaneously solve  $n$  linear equations:

$$\underbrace{\begin{bmatrix} a_{*1} & \dots & a_{*n} \end{bmatrix}}_{AM} \begin{bmatrix} x_1 & \dots & x_n \end{bmatrix} = I, x_i \in \mathbb{R}^n, i = 1, \dots, n$$
$$\begin{bmatrix} a_{*1} & \dots & a_{*n} & | & I \end{bmatrix}$$

$$\begin{bmatrix} LU & | & I \end{bmatrix} \Rightarrow LUx_1 = I_{*1}, \dots, LUx_n = I_{*n}.$$

- Gauss-Jordan Method: Let  $R_1, \dots, R_k$  represent row transformation matrices, not necessarily lower triangular, such that  $R_k \cdot R_{k-1} \cdot \dots \cdot R_1 A = I$ , then  $A^{-1} = R_k \cdot R_{k-1} \cdot \dots \cdot R_1$ .

# Computing Matrix Inverses

- $A^{-1}$  is the unique matrix such that  $A^{-1}A = AA^{-1} = I$ .
- Simultaneously solve  $n$  linear equations:

$$\underbrace{\begin{bmatrix} a_{*1} & \dots & a_{*n} \end{bmatrix}}_{AM} \begin{bmatrix} x_1 & \dots & x_n \end{bmatrix} = I, x_i \in \mathbb{R}^n, i = 1, \dots, n$$
$$\begin{bmatrix} LU & | & I \end{bmatrix} \Rightarrow LUx_1 = I_{*1}, \dots, LUx_n = I_{*n}.$$

- Gauss-Jordan Method: Let  $R_1, \dots, R_k$  represent row transformation matrices, not necessarily lower triangular, such that  $R_k \cdot R_{k-1} \cdot \dots \cdot R_1 A = I$ , then  $A^{-1} = R_k \cdot R_{k-1} \cdot \dots \cdot R_1$ .
- Thus,

$$R_k \cdot R_{k-1} \cdot \dots \cdot R_1 \begin{bmatrix} A & | & I \end{bmatrix} =$$

# Computing Matrix Inverses

- $A^{-1}$  is the unique matrix such that  $A^{-1}A = AA^{-1} = I$ .
- Simultaneously solve  $n$  linear equations:

$$\underbrace{\begin{bmatrix} a_{*1} & \dots & a_{*n} \end{bmatrix}}_{AM} \begin{bmatrix} x_1 & \dots & x_n \end{bmatrix} = I, x_i \in \mathbb{R}^n, i = 1, \dots, n$$
$$\begin{bmatrix} LU & | & I \end{bmatrix} \Rightarrow LUx_1 = I_{*1}, \dots, LUx_n = I_{*n}.$$

- Gauss-Jordan Method: Let  $R_1, \dots, R_k$  represent row transformation matrices, not necessarily lower triangular, such that  $R_k \cdot R_{k-1} \cdot \dots \cdot R_1 A = I$ , then  $A^{-1} = R_k \cdot R_{k-1} \cdot \dots \cdot R_1$ .
- Thus,

$$R_k \cdot R_{k-1} \cdot \dots \cdot R_1 \begin{bmatrix} A & | & I \end{bmatrix} = \begin{bmatrix} I & | & A^{-1} \end{bmatrix}$$



# Computational Cost

- Additions and multiplications in worst case in computing the  $LU$  decomposition?

# Computational Cost

- Additions and multiplications in worst case in computing the  $LU$  decomposition?
- $\mathcal{O}(n^3)$

# Computational Cost

- Additions and multiplications in worst case in computing the  $LU$  decomposition?
- $\mathcal{O}(n^3)$
- Cost involved in Back-substitution?

# Computational Cost

- Additions and multiplications in worst case in computing the  $LU$  decomposition?
- $\mathcal{O}(n^3)$
- Cost involved in Back-substitution?
- $\mathcal{O}(n^2)$

# Computational Cost

- Additions and multiplications in worst case in computing the  $LU$  decomposition?
- $\mathcal{O}(n^3)$
- Cost involved in Back-substitution?
- $\mathcal{O}(n^2)$
- Cost for computing  $A^{-1}$  using Gauss-Jordan method?

# Computational Cost

- Additions and multiplications in worst case in computing the  $LU$  decomposition?
- $\mathcal{O}(n^3)$
- Cost involved in Back-substitution?
- $\mathcal{O}(n^2)$
- Cost for computing  $A^{-1}$  using Gauss-Jordan method?
- $\mathcal{O}(n^3)$

# Computational Cost

- Additions and multiplications in worst case in computing the  $LU$  decomposition?
- $\mathcal{O}(n^3)$
- Cost involved in Back-substitution?
- $\mathcal{O}(n^2)$
- Cost for computing  $A^{-1}$  using Gauss-Jordan method?
- $\mathcal{O}(n^3)$
- Why should one use  $LU$  decomposition?