Residue-pair PMFs (6×6)

each mini panel: x 0.3–1.8 nm, y −1.75…2.5 kcal/mol

Three-residue window PMFs

same axes; rows/cols = window on pep 1 / pep 2

LIVA–LIVA fragment PMF

mean of halves ± |PMF₁−PMF₂|/2

H-bond occupancy (LIVA), by direction

rows = donor, cols = acceptor; per contact frame, mean ± |h₁−h₂|/2

H-bond network

filled = pep 1, open = pep 2; arrow = donor N–H → acceptor O

pep 1 donates (system colour) pep 2 donates (gray)

pep1–pep2 angle

mean of half densities ± |h₁−h₂|/2

End-to-end distance (pep 1)

mean of half densities ± |h₁−h₂|/2

LIVAGE (E⁻/E⁻)

GLU/GLU

−1.31 ± 0.10 at 0.42 nm

mean 106.6

mean 13.0

LIVAGE (E⁻/Eº)

GLU/GLH

−1.08 ± 0.04 at 0.42 nm

mean 98.5

mean 12.9

LIVAGE (Eº/Eº)

GLH/GLH

−0.96 ± 0.15 at 0.43 nm

mean 99.1

mean 12.2

LIVAGG

GLY

−1.13 ± 0.13 at 0.42 nm

mean 105.1

mean 12.7

LIVAGH (H⁺)

HIP

−0.85 ± 0.09 at 0.42 nm

mean 105.0

mean 12.5

LIVAGK

LYS

−1.13 ± 0.03 at 0.52 nm

mean 105.9

mean 12.8

LIVAGQ

GLN

−1.04 ± 0.02 at 0.52 nm

mean 100.7

mean 12.3

LIVAGR

ARG

−0.93 ± 0.05 at 0.42 nm

mean 105.2

mean 12.4

PMF (kcal/mol)

COM distance (nm)

angle (deg)

end-to-end distance (Å)

density

donor (pep 1)

acceptor (pep 2)

donor (pep 2)

acceptor (pep 1)

pep 1 → pep 2 occupancy (each row in its system colour)

pep 2 → pep 1 occupancy