

Hartshorne: Varieties

February 2024

February 9, 2024

Contents

1	Affine Varieties	2
1.1	Familiar Definitions	2
1.2	The Zariski Topology	2

1 Affine Varieties

1.1 Familiar Definitions

Let k be an algebraically closed field. We define **affine space** over k , denoted \mathbb{A}_k^n or \mathbb{A}^n , as the set of all n -tuples with components in k . The elements $P \in \mathbb{A}^n$ are called **points** — and if $P = (a_1, \dots, a_n)$, the elements a_1, \dots, a_n are called **components**.

The set $R = k[x_1, \dots, x_n]$ denotes the commutative ring of polynomials with variables x_1, \dots, x_n with coefficients in k . We may interpret each $f \in R$ as a function from \mathbb{A}^n to k , defined by $f(P) = f(a_1, \dots, a_n)$. We may thus define the **zeroes** of f , given by the set $Z(f) = \{P \in \mathbb{A}_k^n \mid f(P) = 0\}$. More generally, for any subset T of polynomials R , its **zeroes** are given by

$$Z(T) = \{P \in \mathbb{A}^n \mid f(P) = 0 \text{ for each } f \in T\}.$$

Ideals in R are quite elegant: since R is Noetherian, each ideal \mathfrak{a} has a finite set of generators f_1, \dots, f_n . Thus \mathfrak{a} may be expressed as the common zeroes of f_1, \dots, f_n . It is easy to verify that if \mathfrak{b} is the ideal generated by T , then $Z(T) = Z(\mathfrak{b})$.

1.2 The Zariski Topology

A subset $Y \subseteq \mathbb{A}^n$ is an **algebraic set** if there exists a subset $T \subseteq R$ such that $Y = Z(T)$.

Theorem 1. *The following two results hold:*

1. *The union of two algebraic sets $X, Y \subseteq \mathbb{A}^n$ is algebraic.*
2. *The intersection of any family of algebraic sets Y_α is an algebraic set.*

Proof. For (1), let $X = Z(T)$ and $Y = Z(S)$. We claim that $Z(T \cup S) = Z(TS)$, where TS is the set of all ts with $t \in T$ and $s \in S$.

1. Suppose $P \in Z(TS)$ — that is, $ts(P) = 0$ for all $ts \in TS$. Since R is an integral domain, we have $t(P) = 0$ or $s(P) = 0$, so $P \in Z(T)$ or $P \in Z(S)$. Hence $P \in Z(T) \cup Z(S)$.
2. Suppose $P \in Z(T) \cup Z(S)$. Then $P \in Z(T)$ or $P \in Z(S)$ — in which case, $t(P) = 0$ for all t or $s(P) = 0$ for all s . In either case, $ts(P) = 0$ for all $ts \in TS$, so $P \in Z(TS)$.

Thus $X \cup Y = Z(T) \cup Z(S) = Z(TS)$. We deduce that $X \cup Y$ is an algebraic set. For (2), let $Y_\alpha = Z(T_\alpha)$. It is easy to verify that $\bigcap_{\alpha \in A} Z(T_\alpha) = Z(\bigcup_{\alpha \in A} T_\alpha)$; hence $\bigcup_{\alpha \in A} Y_\alpha$ is an algebraic set. \square

Noting that $\{0\}$ and \mathbb{A}^n are algebraic sets (since $\{0\} = Z(1)$ and $\mathbb{A}^n = Z(0)$), we deduce that algebraic sets in R satisfy the closed set axioms to be a topological space. The ensuing topology is called the **Zariski topology**.