

Centre for Metamaterial Research and Innovation


EPSRC Centre for Doctoral Training in Metamaterials




www.metamaterials.center


## The Adjoint Method

Sandro Mignuzzi, Stefano Vezzoli, Simon A. R. Horsley, William L. Barnes, Stefan A. Maier, and Riccardo Sapienza "Nanoscale Design of the Local Density of Optical States", Nano Lett. 19, 3, 1613–1617 (2019) Owen Miller, "Photonic Design: From Fundamental Solar Cell Physics to Computational Inverse Design", PhD thesis (2012)



 $\delta P = \frac{1}{2} \text{Im} \left[ \mathbf{E}_*(\mathbf{r}) \cdot \mathbf{E}(\mathbf{r}) \right] \delta \varepsilon$ 








## The Adjoint Method

$$\delta P = \frac{1}{2} \operatorname{Im} \left[ \mathbf{E}_*(\mathbf{r}) \cdot \mathbf{E}(\mathbf{r}) \right] \delta \varepsilon$$



- Sandro Mignuzzi, Stefano Vezzoli, Simon A. R. Horsley, William L. Barnes, Stefan A. Maier, and Riccardo Sapienza "Nanoscale Design of the Local Density of Optical States", Nano Lett. 19, 3, 1613–1617 (2019)
  - Owen Miller, "Photonic Design: From Fundamental Solar Cell Physics to Computational Inverse Design", PhD thesis (2012)




Centre for Metamaterial Research and Innovation


EPSRC Centre for Doctoral Training in Metamaterials




www.metamaterials.center


## Continuous → Discrete





$$kR \leq 1$$





$$\begin{pmatrix} \mathbf{p} \\ \mathbf{m} \end{pmatrix} = \begin{pmatrix} \boldsymbol{\alpha}_E & 0 \\ 0 & \boldsymbol{\alpha}_H \end{pmatrix} \begin{pmatrix} \mathbf{E} \\ \mathbf{H} \end{pmatrix}$$

$$\begin{pmatrix} \mathbf{E}(\mathbf{r}) \\ \mathbf{H}(\mathbf{r}) \end{pmatrix} = \begin{pmatrix} \mathbf{E}_{s}(\mathbf{r}) \\ \mathbf{H}_{s}(\mathbf{r}) \end{pmatrix} + \sum_{n} \begin{pmatrix} \xi^{2}\mathbf{G}(\mathbf{r}, \mathbf{r}_{n})\boldsymbol{\alpha}_{E} & i\xi\nabla\times\mathbf{G}(\mathbf{r}, \mathbf{r}_{n})\boldsymbol{\alpha}_{H} \\ -i\xi\nabla\times\mathbf{G}(\mathbf{r}, \mathbf{r}_{n})\boldsymbol{\alpha}_{E} & \xi^{2}\mathbf{G}(\mathbf{r}, \mathbf{r}_{n})\boldsymbol{\alpha}_{H} \end{pmatrix} \begin{pmatrix} \mathbf{E}(\mathbf{r}_{n}) \\ \mathbf{H}(\mathbf{r}_{n}) \end{pmatrix}$$

Scattered

$$\delta \mathbf{E}(\mathbf{r}) = -\left[\xi^2 \mathbf{G}(\mathbf{r}, \mathbf{r}_n) \alpha_E \nabla \mathbf{E}(\mathbf{r}_n) + i \xi \nabla \times \mathbf{G}(\mathbf{r}, \mathbf{r}_n) \alpha_H \nabla \mathbf{H}(\mathbf{r}_n)\right] \delta \mathbf{r}_n$$

Purcell, E. M. and Pennypacker, C. R., "Scattering and absorption of light by nonspherical dielectric grains" Astrophysical Journal 7