

* Solutions *

- $$\begin{bmatrix} 1 & 1 & 0 & 0 & 2 & 4b_2 - 3b_1 - 4b_3 \\ 0 & 0 & 1 & 0 & 1 & b_2 - 2b_1 \\ 0 & 0 & 0 & 1 & 0 & b_1 - b_2 + b_3 \\ 0 & 0 & 0 & 0 & 0 & b_4 - b_1 - b_3 \\ 0 & 0 & 0 & 0 & 0 & b_5 - 4b_1 \end{bmatrix}, \text{ where } \mathbf{b} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \\ b_5 \end{bmatrix}.$$
 Answer the following questions about this system.

- ☒ (a) x_1
☐ (b) x_2
☒ (c) x_3
- ☒ (d) x_4
☐ (e) x_5
☐ (f) There are no leading variables

- ☐ (a) x_1
☒ (b) x_2
☐ (c) x_3
- ☐ (d) x_4
☒ (e) x_5
☐ (f) There are no free variables

- If yes, list the condition(s) on b_1, b_2, b_3, b_4, b_5 : $b_4 - b_1 - b_3 = 0$
 $b_5 - 4b_1 = 0$

(1.4) For which of the following $\mathbf{b}$, if any, is the system $A\mathbf{x} = \mathbf{b}$ consistent? **Select ALL that apply.**

- ☒ (a) $\mathbf{b} = \begin{bmatrix} 1 \\ 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}$ ☐ (b) $\mathbf{b} = \begin{bmatrix} 1 \\ 1 \\ 5 \\ 3 \\ 4 \end{bmatrix}$ ☐ (c) $\mathbf{b} = \begin{bmatrix} 1 \\ 1 \\ 4 \\ 3 \\ 1 \end{bmatrix}$ ☒ (d) $\mathbf{b} = \begin{bmatrix} 0 \\ 5 \\ -2 \\ -2 \\ 0 \end{bmatrix}$
- $4 - 4(1) = 0 \checkmark$
 $3 - 1 - 2 = 0 \checkmark$
 $4 - 4(1) = 0 \checkmark$
 $3 - 1 - 5 \neq 0$
 $1 - 4(1) \neq 0$
 $0 - 4(0) = 0 \checkmark$
 $-2 - 0 - (-2) = 0 \checkmark$
- ☐ (e) $\mathbf{b} = \begin{bmatrix} 2 \\ 1 \\ 4 \\ 1 \\ 8 \end{bmatrix}$ ☒ (f) $\mathbf{b} = \begin{bmatrix} 1 \\ 2 \\ 4 \\ 5 \\ 4 \end{bmatrix}$ ☒ (g) $\mathbf{b} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$ ☐ (h) none
- $8 - 4(2) = 0 \checkmark$
 $1 - 2 - 4 \neq 0$
 $4 - 4(1) = 0 \checkmark$
 $5 - 1 - 4 = 0 \checkmark$

- If yes, list the condition(s) on b_1, b_2, b_3, b_4, b_5 ?

There are free variables. The system is either inconsistent or it will be consistent with infinitely many solutions.

2. (12 points) In each of the following cases, decide if the linear system corresponding to the augmented matrix is consistent or inconsistent by filling in the bubble corresponding to your answer. If the system is consistent, write the **vector form of the solution**.

(2.1)
$$\left[\begin{array}{cccc|c} 1 & 0 & 0 & 3 & 2 \\ 0 & 1 & 0 & 1 & 4 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right]$$
 ☐ consistent ☒ inconsistent

If consistent, solution $\mathbf{x} =$

(2.2)
$$\left[\begin{array}{cccc|c} 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$
 ☒ consistent ☐ inconsistent

$x_1 + x_2 = 0$
 $x_3 = 0 \Rightarrow \begin{cases} x_1 = -x_2 \\ x_2 = x_2 \\ x_3 = 0 \end{cases}$

If consistent, solution $\mathbf{x} = x_2 \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$

(2.3)
$$\left[\begin{array}{cccc|c} 1 & 0 & 0 & 0 & 6 \\ 0 & 1 & 0 & 0 & -5 \\ 0 & 0 & 1 & 0 & 2 \\ 0 & 0 & 0 & 1 & 0 \end{array} \right]$$
 ☒ consistent ☐ inconsistent

If consistent, solution $\mathbf{x} = \begin{bmatrix} 6 \\ -5 \\ 2 \\ 0 \end{bmatrix}$

(2.4)
$$\left[\begin{array}{ccccc|c} 1 & 0 & 1 & -4 & 0 & -3 \\ 0 & 1 & 2 & 1 & 0 & 8 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{array} \right]$$
 ☒ consistent ☐ inconsistent

$x_1 + x_3 - 4x_4 = -3 \Rightarrow x_1 = -3 - x_3 + 4x_4$
 $x_2 + 2x_3 + x_4 = 8 \Rightarrow x_2 = 8 - 2x_3 - x_4$
 $x_5 = 0 \Rightarrow \begin{cases} x_1 = -3 - x_3 + 4x_4 \\ x_2 = 8 - 2x_3 - x_4 \\ x_3 = x_3 \\ x_4 = x_4 \\ x_5 = 0 \end{cases}$

If consistent, solution $\mathbf{x} = \begin{bmatrix} -3 \\ 8 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} -1 \\ -2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_4 \begin{bmatrix} 4 \\ -1 \\ 0 \\ 1 \\ 0 \end{bmatrix}$

3. (18 points) Let $A = \begin{bmatrix} 1 & 4 \\ 2 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 1 & -2 \\ 3 & 2 \end{bmatrix}$, $C = \begin{bmatrix} 1 & -2 \\ 1 & 0 \\ 2 & -1 \end{bmatrix}$, $\mathbf{u} = \begin{bmatrix} 1 \\ 3 \\ -2 \end{bmatrix}$, $\mathbf{w} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$. Compute the following expressions when they are defined or state they are undefined and explain why.

(3.1) $A^T + 3B = \begin{bmatrix} 1 & 2 \\ 4 & 0 \end{bmatrix} + 3 \begin{bmatrix} 1 & -2 \\ 3 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 4 & 0 \end{bmatrix} + \begin{bmatrix} 3 & -6 \\ 9 & 6 \end{bmatrix} = \begin{bmatrix} 4 & -4 \\ 13 & 6 \end{bmatrix}$

(3.2) $B + AC^T$

(2×2)
 $(2 \times 2) (2 \times 3)$
 (2×3)

Undefined, B is (2×2) and AC^T is (2×3)
 cannot add matrices that are different sizes.

- (3.3) BC Undefined, the # of columns of B is not equal to the # of rows of C . (Inner dimensions do not match)

(3.4) $CB = \begin{bmatrix} 1 & -2 \\ 1 & 0 \\ 2 & -1 \end{bmatrix} \begin{bmatrix} 1 & -2 \\ 3 & 2 \end{bmatrix} = \begin{bmatrix} -5 & -6 \\ 1 & -2 \\ -1 & -6 \end{bmatrix}$

$(3 \times 2) (2 \times 2)$

- (3.5) $C\mathbf{u}$
- $(3 \times 2) (3 \times 1)$
 $\neq$
- Undefined, the # of columns of C is not equal to the # of rows of $\mathbf{u}$. (Inner dimensions do not match)

(3.6) $C\mathbf{w} = \begin{bmatrix} 1 & -2 \\ 1 & 0 \\ 2 & -1 \end{bmatrix} \begin{bmatrix} 3 \\ 1 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix} + 1 \begin{bmatrix} -2 \\ 0 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix}$

$(3 \times 2) (2 \times 1)$

4. (10 points) Identify whether each one of the given sets is linearly dependent or linearly independent by filling in the bubble corresponding to your answers. Explanations are not necessary. *Hint:* all can be done by inspection.

(4.1) $\left\{ \begin{bmatrix} -2 \\ 4 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 3 \\ 3 \end{bmatrix} \right\}$ ☒ linearly dependent ☐ linearly independent

(4.2) $\left\{ \begin{bmatrix} 1 \\ 3 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \end{bmatrix} \right\}$ ☐ linearly dependent ☐ linearly independent

(4.3) $\left\{ \begin{bmatrix} -4 \\ 2 \end{bmatrix}, \begin{bmatrix} 8 \\ -4 \end{bmatrix} \right\}$ ☒ linearly dependent ☐ linearly independent

(4.4) $\left\{ \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}$ ☒ linearly dependent ☐ linearly independent

(4.5) $\left\{ \begin{bmatrix} 4 \\ 4 \end{bmatrix} \right\}$ ☐ linearly dependent ☒ linearly independent

(4.6) $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \right\}$ ☒ linearly dependent ☐ linearly independent

(4.7) $\left\{ \begin{bmatrix} 3 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 6 \\ 4 \\ 4 \end{bmatrix} \right\}$ ☐ linearly dependent ☒ linearly independent

(4.8) $\left\{ \begin{bmatrix} 3 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right\}$ ☒ linearly dependent ☐ linearly independent

(4.9) $\left\{ \begin{bmatrix} 3 \\ 6 \\ 9 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \right\}$ ☐ linearly dependent ☐ linearly independent

(4.10) $\left\{ \begin{bmatrix} 2 \\ 1 \\ -3 \end{bmatrix}, \begin{bmatrix} 6 \\ 3 \\ -9 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \right\}$ ☒ linearly dependent ☐ linearly independent

5. (8 points) Let $A = \begin{bmatrix} 1 & -3 & 4 \\ 0 & 1 & -3 \\ -2 & 7 & a \end{bmatrix}$ and $B = \begin{bmatrix} 1 & -3 & 4 \\ 0 & 1 & -3 \\ -2 & 7 & a+1 \end{bmatrix}$.

(5.1) Find the value(s) of a so that the vectors $\begin{bmatrix} 1 \\ 0 \\ -2 \end{bmatrix}$, $\begin{bmatrix} -3 \\ 1 \\ 7 \end{bmatrix}$, and $\begin{bmatrix} 4 \\ -3 \\ a \end{bmatrix}$ are linearly dependent.

$$\left[\begin{array}{ccc|c} 1 & -3 & 4 & 0 \\ 0 & 1 & -3 & 0 \\ -2 & 7 & a & 0 \end{array} \right] \xrightarrow{R_3 \rightarrow R_3 + 2R_2} \left[\begin{array}{ccc|c} 1 & -3 & 4 & 0 \\ 0 & 1 & -3 & 0 \\ 0 & 1 & a+8 & 0 \end{array} \right] \xrightarrow{R_3 \rightarrow R_3 - R_2} \left[\begin{array}{ccc|c} 1 & -3 & 4 & 0 \\ 0 & 1 & -3 & 0 \\ 0 & 0 & a+11 & 0 \end{array} \right]$$

$a+11=0$
 $\Rightarrow a=-11$

(5.2) For the value of a you found in 5.1, is the matrix A nonsingular (i.e., invertible)? ☐ yes ☒ no

(5.3) For the value of a you found in 5.1, is the matrix B nonsingular? ☒ yes ☐ no

6. (5 points) For each true/false question, completely fill in the answer bubble that corresponds to your answer. Explanations are not necessary.

(6.1) ☐ (T) ☒ (F) It is possible for a (5×5) system of linear equations to have exactly 5 solutions.

(6.2) ☐ (T) ☒ (F) If A is a (4×5) matrix and $\mathbf{0}$ is the zero vector in $\mathbb{R}^5$, then it is possible that $A\mathbf{x} = \mathbf{0}$ has a unique solution.

(6.3) ☒ (T) ☐ (F) If A is $(n \times n)$ and nonsingular then $A\mathbf{x} = \mathbf{b}$ has a unique solution in $\mathbb{R}^n$.

(6.4) ☐ (T) ☒ (F) If A is a (5×4) matrix and $\mathbf{b}$ is any vector in $\mathbb{R}^4$, then $A\mathbf{x} = \mathbf{b}$ always has at least one solution.

(6.5) ☒ (T) ☐ (F) If A is an $(m \times n)$ matrix and C is an $(n \times p)$ matrix then $(AC)^T = C^T A^T$.

7. (10 points) Let $A = \begin{bmatrix} 1 & 0 & 2 \\ -3 & 1 & -6 \\ 2 & 3 & 5 \end{bmatrix}$.

(7.1) The matrix A **is invertible**. Find its inverse A^{-1} . Show all your work and write your final answer in the space provided.

$$A^{-1} = \begin{bmatrix} 23 & 6 & -2 \\ 3 & 1 & 0 \\ -11 & -3 & 1 \end{bmatrix}$$

$$\begin{aligned} & \left[\begin{array}{ccc|ccc} 1 & 0 & 2 & 1 & 0 & 0 \\ -3 & 1 & -6 & 0 & 1 & 0 \\ 2 & 3 & 5 & 0 & 0 & 1 \end{array} \right] \begin{array}{l} R_2 \rightarrow R_2 + 3R_1 \\ R_3 \rightarrow R_3 - 2R_1 \end{array} \quad \left[\begin{array}{ccc|ccc} 1 & 0 & 2 & 1 & 0 & 0 \\ 0 & 1 & 0 & 3 & 1 & 0 \\ 0 & 3 & 1 & -2 & 0 & 1 \end{array} \right] \\ & \quad R_3 \rightarrow R_3 - 3R_2 \quad \left[\begin{array}{ccc|ccc} 1 & 0 & 2 & 1 & 0 & 0 \\ 0 & 1 & 0 & 3 & 1 & 0 \\ 0 & 0 & 1 & -11 & -3 & 1 \end{array} \right] \\ & \quad R_1 \rightarrow R_1 - 2R_3 \quad \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 23 & 6 & -2 \\ 0 & 1 & 0 & 3 & 1 & 0 \\ 0 & 0 & 1 & -11 & -3 & 1 \end{array} \right] \end{aligned}$$

(7.2) Use your answer from 7.1 to solve the system $A\mathbf{x} = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}$.

$$\vec{x} = A^{-1} \vec{b} = \begin{bmatrix} 23 & 6 & -2 \\ 3 & 1 & 0 \\ -11 & -3 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix} = \begin{bmatrix} 13 \\ 2 \\ -6 \end{bmatrix}$$

8. (5 points) Let A and B be the matrices such that $A^{-1} = \begin{bmatrix} 1 & 1 & 5 \\ 0 & 1 & 6 \\ 0 & -2 & -4 \end{bmatrix}$ and $B^{-1} = \begin{bmatrix} 1 & 3 & -1 \\ 0 & 3 & 1 \\ -1 & 1 & 1 \end{bmatrix}$.
 Compute $(AB)^{-1}$, or explain why it is not possible.

$$\begin{aligned} (AB)^{-1} &= B^{-1} A^{-1} = \begin{bmatrix} 1 & 3 & -1 \\ 0 & 3 & 1 \\ -1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 5 \\ 0 & 1 & 6 \\ 0 & -2 & -4 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 6 & 27 \\ 0 & 1 & 14 \\ -1 & -2 & -3 \end{bmatrix} \end{aligned}$$

9. (4 points) Which of the following matrices are symmetric? Select **ALL** answers that apply. No credit will be given if all answers are left blank or all answers are selected. No work required.

☒ (a) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$

☐ (b) $\begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$

☐ (c) $\begin{bmatrix} 1 & -2 & -3 \\ 2 & 3 & -4 \\ 3 & 4 & 1 \end{bmatrix}$

☒ (d) $\begin{bmatrix} 1 & 3 & 2 \\ 3 & 6 & 8 \\ 2 & 8 & 7 \end{bmatrix}$

☐ (e) $\begin{bmatrix} -3 & 7 & 4 \\ 7 & -3 & 5 \\ 4 & 6 & -3 \end{bmatrix}$

☒ (f) $\begin{bmatrix} 2 & -1 & 3 \\ -1 & 7 & -4 \\ 3 & -4 & 0 \end{bmatrix}$

☐ (g) $\begin{bmatrix} 1 & 3 & 1 \\ 3 & 1 & 3 \end{bmatrix}$

☐ (h) $\begin{bmatrix} 2 & -3 & 4 \\ -1 & 7 & -3 \\ 4 & -1 & 2 \end{bmatrix}$

10. (12 points) Let $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$. Indicate if the subset W of $\mathbb{R}^3$ is a subspace of $\mathbb{R}^3$ by filling in the bubble corresponding to your answers. In the cases where W is **not** a subspace, give a **specific** counterexample that **clearly shows which property fails**. In the cases where W is a subspace of $\mathbb{R}^3$, give a geometric description of W . (You do NOT need to verify in general.)

(10.1) $W = \{\mathbf{x} : 2x_1 - 5x_2 + 8x_3 = 0\}$ ☐ Not a subspace of $\mathbb{R}^3$ ☒ W is a subspace of $\mathbb{R}^3$

Counterexample OR geometric description of W :

Plane through origin with normal vector $\vec{n} = \langle 2, -5, 8 \rangle$
must be stated

(10.2) $W = \{\mathbf{x} : x_1 \leq 0\}$ ☐ Not a subspace of $\mathbb{R}^3$ ☐ W is a subspace of $\mathbb{R}^3$

Counterexample OR geometric description of W :

$\vec{x} = \begin{bmatrix} -1 \\ 0 \\ 0 \end{bmatrix} \in W$ but if $a = -1$, then $a\vec{x} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \notin W$ since $x_1 > 0$

(10.3) $W = \{\mathbf{x} : x_3 = x_2 = 5x_1\}$ ☐ Not a subspace of $\mathbb{R}^3$ ☒ W is a subspace of $\mathbb{R}^3$

Counterexample OR geometric description of W :

$\vec{x} = \begin{bmatrix} x_1 \\ 5x_1 \\ 5x_1 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 5 \\ 5 \end{bmatrix}$
line through the origin with direction vector $\begin{bmatrix} 1 \\ 5 \\ 5 \end{bmatrix}$
must be stated

(10.4) $W = \{\mathbf{x} : x_1x_3 = 0\}$ ☒ Not a subspace of $\mathbb{R}^3$ ☐ W is a subspace of $\mathbb{R}^3$

Counterexample OR geometric description of W :

Let $\vec{x} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \in W$ and $\vec{y} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \in W$
since $0(1) = 0$ since $1(0) = 0$

$\vec{x} + \vec{y} = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \notin W$ since $1(1) = 1 \neq 0$