

Elliptic PDEs of 2nd Order, Gilbarg and Trudinger

Chapter 7 Sobolev Spaces*

Yung-Hsiang Huang[†]

1. *Proof.* □
2. *Proof.* □
3. *Proof.* □
4. **Prove that the product formula**

$$D(uv) = vDu + uDv$$

holds for all $u, v \in W^1(\Omega)$ **such that** $uv, uDv + vDu \in L^1_{loc}(\Omega)$.

Remark 1. The key idea to define U_n comes from <http://math.stackexchange.com/questions/443425/product-rule-of-weak-derivatives>

Proof. Step 1 is to prove the theorem under the additional assumption $v \in C^1$. This is standard, so we just state it without proof.

Step 2 is to prove the theorem under the additional assumption $u \in W^1(\Omega) \cap L^\infty(\Omega)$. This is proved by Step 1 and Theorem 7.4, a characterization of weakly differentiability. This is also standard.

Step 3 is consider the problem under the assumption $u, v \geq 0$: we first assume $v \geq 1$ and define $U_n := \min\{u, \frac{n}{v}\}$. By Lemma 7.6, $U_n \in W^1(\Omega) \cap L^\infty(\Omega)$, and $D(U_nv) = Du\chi_{\{uv < n\}} + (-\frac{nDv}{v^2})\chi_{\{uv \geq n\}}$. Hence $U_nDv + vDU_n = (uDv + vDu)\chi_{\{uv < n\}} \in L^1_{loc}(\Omega)$. Step 2 implies that, for each $\phi \in C_0^1(\Omega)$,

$$\int_{\Omega} U_nvD\phi = - \int_{\Omega} [U_nDv + vDU_n]\phi.$$

*Last Modified: 2017/07/15.

[†]Department of Math., National Taiwan University. Email: d04221001@ntu.edu.tw

Note that $U_n v = \min\{uv, n\} \nearrow uv$, $U_n Dv + v D U_n \rightarrow u Dv + v Du$ a.e. and $|U_n Dv + v D U_n| \leq |u Dv + v Du| \in L^1(\text{supp } \phi)$, MCT and LDCT imply that

$$\begin{aligned} \int_{\Omega} uv D\phi &= \left(\int_{\{D\phi \geq 0\}} + \int_{\{D\phi < 0\}} \right) uv D\phi = \lim_{n \rightarrow \infty} \left(\int_{\{D\phi \geq 0\}} + \int_{\{D\phi < 0\}} \right) U_n v D\phi \\ &= \lim_{n \rightarrow \infty} \int_{\Omega} U_n v D\phi = \lim_{n \rightarrow \infty} - \int_{\Omega} [U_n Dv + v D U_n] \phi = - \int_{\Omega} (u Dv + v Du) \phi. \end{aligned}$$

The above formula is also true for general $v \geq 0$, since

$$\int_{\Omega} uv D\phi + \int_{\Omega} u D\phi = \int_{\Omega} u(v+1) D\phi = - \int_{\Omega} (u D(v+1) + (v+1) Du) \phi = - \int_{\Omega} (u Dv + v Du) \phi - \int_{\Omega} Du \phi.$$

Finally, we decompose $u = u^+ - u^-$, $v = v^+ - v^-$. Since $u^+ v^+ = uv \chi_{\{u>0, v>0\}}$ and $u^+ Dv^+ + v^+ Du^+ = (u Dv + v Du) \chi_{\{u>0, v>0\}}$, they are in $L^1_{loc}(\Omega)$. Other similar functions also in $L^1_{loc}(\Omega)$ without proof. Then by Step 3, for each $\phi \in C_0^1(\Omega)$,

$$\begin{aligned} \int_{\Omega} uv D\phi &= \int_{\Omega} (u^+ v^+ - u^+ v^- - u^- v^+ + u^- v^-) D\phi \\ &= - \int_{\Omega} \left((u^+ Dv^+ + v^+ Du^+) - (u^+ Dv^- + v^- Du^+) - (u^- Dv^+ + v^+ Du^-) + (u^- Dv^- + v^- Du^-) \right) \phi \\ &= - \int_{\Omega} (u Dv + v Du) \phi. \end{aligned}$$

□

Remark 2. In the same website, Mateusz Kwanicki (from Wrocław University of Science and Technology?) informed me that this problem may be solved by the characterization of absolutely continuity along any line. He said:

”How about using the ACL characterisation of differentiability? I mean, u has weak partial derivatives if and only if it is absolutely continuous on almost every line in any cardinal direction, and the classical partial derivatives (defined thus almost everywhere) are integrable.

Since the product of absolutely continuous functions is absolutely continuous, with $(fg)' = f'g + fg'$, it follows that uv has the ACL property, and the classical partial derivatives of uv obey the product rule almost everywhere. In order that uv has weak partial derivatives it is therefore sufficient to assume that the classical partial derivatives of uv are locally integrable.”

5. *Proof.*

□

6. *Proof.*

□

7. *Proof.*

□

- 8. *Proof.* □
- 9. *Proof.* □
- 10. *Proof.* □
- 11. *Proof.* □
- 12. *Proof.* □
- 13. *Proof.* □
- 14. *Proof.* □
- 15. *Proof.* □
- 16. *Proof.* □
- 17. *Proof.* □
- 18. *Proof.* □