

Math 480 Homework SVD updates

James Pak

May 22, 2013

1 Introduction

Consider the real matrix $X \in \mathbb{R}^{p \times q}$. Then the singular value decomposition (SVD) diagonalizes X with orthonormal matrices $U \in \mathbb{R}^{p \times p}$ and $V \in \mathbb{R}^{q \times q}$. Note that $S = U^T X V$.

1.1 Formula's

For Additive Modification, see Section 2. For Rank-1 Modification, see Section 3.

2 Additive Modification

Let $A \in \mathbb{R}^{p \times c}$ and $B \in \mathbb{R}^{p \times c}$. Then

$$X + AB^T = [U \ A] \begin{bmatrix} S & 0 \\ 0 & I \end{bmatrix} [V \ B]^T$$

Furthermore, let P be the orthogonal basis of the column space of $(I - UU^T)A$ and $R_A = P^T(I - UU^T)A$

$$[U \ A] = [U \ P] \begin{bmatrix} I & U^T A \\ 0 & R_A \end{bmatrix}$$

Let $QR_B = (I - VV^T)B$. Then

$$X + AB^T = [U \ P] K [V \ Q]^T$$

and

$$K = \begin{bmatrix} I & U^T A \\ 0 & R_A \end{bmatrix} \begin{bmatrix} S & 0 \\ 0 & I \end{bmatrix} \begin{bmatrix} I & V^T B \\ 0 & R_B \end{bmatrix}^T = \begin{bmatrix} S & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} U^T A \\ R_A \end{bmatrix} \begin{bmatrix} V^T B \\ R_B \end{bmatrix}^T$$

Diagonalizing K as $U'^T K V' = S'$ give the rotations U' and V' of the extended subspaces $[U \ P]$ and $[V \ Q]$ such that

$$X + AB^T = ([U \ P] U') S' ([V \ Q] V')^T$$

is the desired SVD.

3 Rank-1 Modification

3.1 Update

3.2 Downdate

3.3 Revise

3.4 Recenter

For each operate in subsection 3.1 - 3.4

1. Known for 3.1:

$$US [V^T \ 0] = [X \ 0]$$

2. Known for 3.2:

$$USV^T = [X \ c]$$

3. Known for 3.3:

$$USV^T = [X \ c]$$

4. Known for 3.4:

$$USV^T = X$$

4 Reference

All work within this paper is taken from Matthew Brand's article on "Fast low-rank modifications of the thing singular value decomposition."