

More Math on Markov Chains

September 29, 2020

Definition 1. The n -step transition probability, p_{ij}^n , is defined as

$$p_{ij}^n = \Pr(X_n = j | X_0 = i)$$

Definition 2. We say that state i **leads to** state j (written as $i \rightarrow j$) if $\exists n \geq 0$ such that $p_{ij}^n > 0$.

Therefore, i communicates with j iff $i \rightarrow j$ and $j \rightarrow i$, written as $i \longleftrightarrow j$. This relation partitions the state space into disjoint sets of states, called communication classes.

Definition 3. A Markov Chain is **irreducible** if it only has one communication class.

0.1 Transient and Recurrent States

We define the first passage time to state i by $T_i = \min_n \{n \geq 1 : X_n = i\}$. Then we define

$$f_{ij}^m \equiv \Pr_i(T_j = m) = \Pr(X_m = j, X_{m-1} \neq j, \dots, X_2 \neq j, X_1 \neq j | X_0 = i)$$

and

$$f_{ij} = \sum_{m=1}^{\infty} f_{ij}^m$$

Definition 4. A state i is **recurrent** if $f_{ii} = 1$ and **transient** if $f_{ii} < 1$.

In a finite irreducible Markov chain, all the states are recurrent.

0.2 Periodicity