

$r: X \rightarrow A; r|_A = \text{id}_A$
 $j: A \rightarrow X; j|_A = \text{id}_A$

$r \circ j: A \rightarrow A$

NOTE:
 $r_*: \pi_1(X, x) \rightarrow \pi_1(A, a)$
 $j_*: \pi_1(A, a) \rightarrow \pi_1(X, x)$

Thm 52.4: $(r \circ j)_*: \pi_1(A, a) \rightarrow \pi_1(A, a)$
 $L \mapsto r_* \circ j_*$ is identity map of fund. groups!!

WTS j_* is injective.

Well. Let $j_*([f]) = j_*([g])$
 $\Rightarrow (r_* \circ j_*)([f]) = (r_* \circ j_*)([g])$
 $\Rightarrow [f] = [g]. \text{ epic.}$

Theorem 2. No retraction Theorem

There is no retraction of B^2 onto S^1 , aka the unit disk onto the unit circle.

This is a silly little definition-based proof. Let's look at the disk; it is star-convex (recall Chapter 52??). This means it must be simply connected, and thus its fundamental group is trivial (page 333).

Now let's look at S^1 and its fundamental group. By Theorem 54.5, we know the fundamental group of S^1 is isomorphic to the additive set of integers. Great. It is nontrivial.

Okay, now that those observations are out of the way, let's suppose towards a contradiction that there is a retraction of B^2 onto S^1 . By Lemma 55.1 (above), this means that inclusion $j : S^1 \rightarrow B^2$ must induce an injective j_* . But guys. Horrible news ahead. j_* maps the fundamental group of S^1 , which is nontrivial, to the fundamental group of B^2 , which is trivial. So it can't be injective. Oh my god gasp a contradiction. Crazy time! I wanted to also briefly mention a generalization about this; the source I will link to Piazza.

Theorem 3. *Let X be a topological space homeomorphic to the underlying space of a 2-dimensional simplicial complex. Then there exists no retraction $r : X \rightarrow \partial X$*

Several comments on this. Given my brief and hasty research on what a "simplicial complex" means, I am very unsure. But, given our knowledge of the proof, I assume we can make the less stronger statement that


Theorem 4. *Let X be a topological space homeomorphic to a simply connected space. Then there exists no retraction $r : X \rightarrow \partial X$*

Again, I am very unsure. But this is cool!

Theorem 5. (*Brouwer fixed-point theorem for disc*) If $f : B^2 \rightarrow B^2$ is continuous, then there exists a point $x \in B^2$ such that $f(x) = x$.

Proof Sketch (Hatcher):

Suppose for a contradiction...



We can define $r : B^2 \rightarrow S^1$, which is a retraction!

Note that $w_0 \simeq w_1$ in B^2 , but $w_0 \not\simeq w_1$ in S^1 . Yet we can force a composition that reaches a contradiction.

Corollary: Let A be a 3 by 3 matrix of positive real numbers. Then A has a positive real eigenvalue.

How is B homeomorphic to B^2 in the textbook proof?