

Majorization-Minimization Algorithm

Theory and Applications

Ying Sun and Daniel P. Palomar

Department of Electronic and Computer Engineering
The Hong Kong University of Science and Technology

ELEC 5470 - Convex Optimization
Fall 2017-18, HKUST, Hong Kong

Slides of this lecture are majorly based on the following works:

- [\[Hun-Lan'J04\]](#) D. R. Hunter, and K. Lange, "A Tutorial on MM Algorithms", *Amer. Statistician*, pp. 30-37, 2004.
- [\[Raz-Hon-Luo'J13\]](#) M. Razaviyayn, M. Hong, and Z. Luo, "A Unified Convergence Analysis of Block Successive Minimization Methods for Nonsmooth Optimization", *SIAM J. Optim.*, pp. 1126-1153, 2013.
- [\[Scu-Fac-Son-Pal-Pan'J14\]](#) G. Scutari, F. Facchinei, Peiran Song, D. P. Palomar, and Jong-Shi Pang, "Decomposition by Partial Linearization: Parallel Optimization of Multi-Agent Systems", *IEEE Trans. Signal Processig*, vol. 62, no. 3, pp. 641-656, Feb. 2014.
- [\[Sun-Bab-Pal'J17\]](#) Y. Sun, P. Babu, and D. P. Palomar, "Majorization-Minimization Algorithms in Signal Processing, Communications, and Machine Learning", *IEEE Trans. Signal Process*, vol. 65, no. 3, pp. 794-816, Feb. 2017.

- 1 The Majorization-Minimization Algorithm
 - Introduction
 - Construction Techniques
 - Example Algorithms
 - Applications
- 2 Block Successive Majorization-Minimization
 - Introduction
 - Block Coordinate Descent
 - Block Successive Majorization-Minimization
 - Example Algorithms
 - Applications
- 3 Distributed Algorithm for Nonlinear Programming
 - Exact Jacobi Successive Convex Approximation
 - Extensions

- 1 The Majorization-Minimization Algorithm
 - Introduction
 - Construction Techniques
 - Example Algorithms
 - Applications
- 2 Block Successive Majorization-Minimization
 - Introduction
 - Block Coordinate Descent
 - Block Successive Majorization-Minimization
 - Example Algorithms
 - Applications
- 3 Distributed Algorithm for Nonlinear Programming
 - Exact Jacobi Successive Convex Approximation
 - Extensions

Problem Statement

- Consider the following optimization problem

$$\begin{array}{ll}\underset{\mathbf{x}}{\text{minimize}} & f(\mathbf{x}) \\ \text{subject to} & \mathbf{x} \in \mathcal{X},\end{array}$$

with \mathcal{X} being a closed convex set and $f(\mathbf{x})$ being continuous.

- $f(\mathbf{x})$ is too complicated to manipulate.
- Idea: successively minimize an approximating function $u(\mathbf{x}, \mathbf{x}^k)$

$$\mathbf{x}^{k+1} = \arg \min_{\mathbf{x} \in \mathcal{X}} u(\mathbf{x}, \mathbf{x}^k),$$

hoping the sequence of minimizers $\{\mathbf{x}^k\}$ will converge to optimal \mathbf{x}^* .

- Question: how to construct $u(\mathbf{x}, \mathbf{x}^k)$?

Problem Statement

- Consider the following optimization problem

$$\begin{array}{ll}\underset{\mathbf{x}}{\text{minimize}} & f(\mathbf{x}) \\ \text{subject to} & \mathbf{x} \in \mathcal{X},\end{array}$$

with \mathcal{X} being a closed convex set and $f(\mathbf{x})$ being continuous.

- $f(\mathbf{x})$ is too complicated to manipulate.
- Idea: successively minimize an approximating function $u(\mathbf{x}, \mathbf{x}^k)$

$$\mathbf{x}^{k+1} = \arg \min_{\mathbf{x} \in \mathcal{X}} u(\mathbf{x}, \mathbf{x}^k),$$

hoping the sequence of minimizers $\{\mathbf{x}^k\}$ will converge to optimal \mathbf{x}^* .

- Question: how to construct $u(\mathbf{x}, \mathbf{x}^k)$?

Problem Statement

- Consider the following optimization problem

$$\begin{array}{ll}\underset{\mathbf{x}}{\text{minimize}} & f(\mathbf{x}) \\ \text{subject to} & \mathbf{x} \in \mathcal{X},\end{array}$$

with \mathcal{X} being a closed convex set and $f(\mathbf{x})$ being continuous.

- $f(\mathbf{x})$ is too complicated to manipulate.
- Idea: successively minimize an approximating function $u(\mathbf{x}, \mathbf{x}^k)$

$$\mathbf{x}^{k+1} = \arg \min_{\mathbf{x} \in \mathcal{X}} u(\mathbf{x}, \mathbf{x}^k),$$

hoping the sequence of minimizers $\{\mathbf{x}^k\}$ will converge to optimal \mathbf{x}^* .

- Question: how to construct $u(\mathbf{x}, \mathbf{x}^k)$?

Terminology

- Distance from a point to a set:

$$d(\mathbf{x}, \mathcal{S}) = \inf_{\mathbf{s} \in \mathcal{S}} \|\mathbf{x} - \mathbf{s}\|.$$

- Directional derivative:

$$f'(\mathbf{x}; \mathbf{d}) \triangleq \liminf_{\lambda \downarrow 0} \frac{f(\mathbf{x} + \lambda \mathbf{d}) - f(\mathbf{x})}{\lambda}.$$

- Stationary point: \mathbf{x} is a stationary point if

$$f'(\mathbf{x}; \mathbf{d}) \geq 0, \quad \forall \mathbf{d} \text{ such that } \mathbf{x} + \mathbf{d} \in \mathcal{X}.$$

Majorization-Minimization

- Construction rule:


$$u(\mathbf{y}, \mathbf{y}) = f(\mathbf{y}), \quad \forall \mathbf{y} \in \mathcal{X} \quad (\text{A1})$$

$$u(\mathbf{x}, \mathbf{y}) \geq f(\mathbf{x}), \quad \forall \mathbf{x}, \mathbf{y} \in \mathcal{X} \quad (\text{A2})$$

$$u'(\mathbf{x}, \mathbf{y}; \mathbf{d})|_{\mathbf{x}=\mathbf{y}} = f'(\mathbf{y}; \mathbf{d}), \quad \forall \mathbf{d} \text{ with } \mathbf{y} + \mathbf{d} \in \mathcal{X} \quad (\text{A3})$$

$$u(\mathbf{x}, \mathbf{y}) \text{ is continuous in } \mathbf{x} \text{ and } \mathbf{y} \quad (\text{A4})$$

- Pictorially:



Algorithm

- Majorization-Minimization (Successive Upper-Bound Minimization):

- 1: Find a feasible point $\mathbf{x}^0 \in \mathcal{X}$ and set $k = 0$
- 2: **repeat**
- 3: $\mathbf{x}^{k+1} = \arg \min_{\mathbf{x} \in \mathcal{X}} u(\mathbf{x}, \mathbf{x}^k)$ (global minimum)
- 4: $k \leftarrow k + 1$
- 5: **until** some convergence criterion is met

Convergence

- Under assumptions A1-A4, every limit point of the sequence $\{\mathbf{x}^k\}$ is a stationary point of the original problem.
- If further assume that the level set $\mathcal{X}^0 = \{\mathbf{x} | f(\mathbf{x}) \leq f(\mathbf{x}^0)\}$ is compact, then

$$\lim_{k \rightarrow \infty} d(\mathbf{x}^k, \mathcal{X}^*) = 0,$$

where \mathcal{X}^* is the set of stationary points.

- 1 The Majorization-Minimization Algorithm
 - Introduction
 - Construction Techniques
 - Example Algorithms
 - Applications
- 2 Block Successive Majorization-Minimization
 - Introduction
 - Block Coordinate Descent
 - Block Successive Majorization-Minimization
 - Example Algorithms
 - Applications
- 3 Distributed Algorithm for Nonlinear Programming
 - Exact Jacobi Successive Convex Approximation
 - Extensions

Construct Surrogate Function

- The performance of Majorization-Minimization algorithm depends crucially on the surrogate function $u(\mathbf{x}, \mathbf{x}^k)$.
- Guideline: the global minimizer of $u(\mathbf{x}, \mathbf{x}^k)$ should be easy to find.
- Suppose $f(\mathbf{x}) = f_1(\mathbf{x}) + \kappa(\mathbf{x})$, where $f_1(\mathbf{x})$ is some “nice” function and $\kappa(\mathbf{x})$ is the one needed to be approximated.

Construction by Convexity

- Suppose $\kappa(t)$ is convex, then

$$\kappa\left(\sum_i \alpha_i t_i\right) \leq \sum_i \alpha_i \kappa(t_i)$$

with $\alpha_i \geq 0$ and $\sum \alpha_i = 1$.

- For example:

$$\begin{aligned}\kappa(\mathbf{w}^T \mathbf{x}) &= \kappa(\mathbf{w}^T (\mathbf{x} - \mathbf{x}^k) + \mathbf{w}^T \mathbf{x}^k) \\ &= \kappa\left(\sum_i \alpha_i \left(\frac{w_i (x_i - x_i^k)}{\alpha_i} + \mathbf{w}^T \mathbf{x}^k\right)\right) \\ &\leq \sum_i \alpha_i \kappa\left(\frac{w_i (x_i - x_i^k)}{\alpha_i} + \mathbf{w}^T \mathbf{x}^k\right)\end{aligned}$$

- If further assume that \mathbf{w} and \mathbf{x} are positive ($\alpha_i = w_i x_i^k / \mathbf{w}^T \mathbf{x}^k$):

$$\kappa(\mathbf{w}^T \mathbf{x}) \leq \sum_i \frac{w_i x_i^k}{\mathbf{w}^T \mathbf{x}^k} \kappa\left(\frac{\mathbf{w}^T \mathbf{x}^k}{x_i^k} x_i\right)$$

- The surrogate functions are separable (parallel algorithm).

Construction by Taylor Expansion

- Suppose $\kappa(\mathbf{x})$ is concave and differentiable, then

$$\kappa(\mathbf{x}) \leq \kappa(\mathbf{x}^k) + \nabla \kappa(\mathbf{x}^k)^T (\mathbf{x} - \mathbf{x}^k),$$

which is a linear upper-bound.

- Suppose $\kappa(\mathbf{x})$ is convex and twice differentiable, then

$$\kappa(\mathbf{x}) \leq \kappa(\mathbf{x}^k) + \nabla \kappa(\mathbf{x}^k)^T (\mathbf{x} - \mathbf{x}^k) + \frac{1}{2} (\mathbf{x} - \mathbf{x}^k)^T \mathbf{M} (\mathbf{x} - \mathbf{x}^k)$$

if $\mathbf{M} - \nabla^2 \kappa(\mathbf{x}) \succeq \mathbf{0}, \forall \mathbf{x}$.

Construction by Inequalities

- Arithmetic-Geometric Mean Inequality:

$$\left(\prod_{i=1}^n x_i \right)^{1/n} \leq \frac{1}{n} \sum_{i=1}^n x_i$$

- Cauchy-Schwartz Inequality:

$$\|\mathbf{x}\| \geq \frac{\mathbf{x}^T \mathbf{x}^k}{\|\mathbf{x}^k\|}$$

- Jensen's Inequality:

$$\kappa(\mathbb{E}\mathbf{x}) \leq \mathbb{E}\kappa(\mathbf{x})$$

with $\kappa(\cdot)$ being convex.

- 1 The Majorization-Minimization Algorithm
 - Introduction
 - Construction Techniques
 - Example Algorithms
 - Applications
- 2 Block Successive Majorization-Minimization
 - Introduction
 - Block Coordinate Descent
 - Block Successive Majorization-Minimization
 - Example Algorithms
 - Applications
- 3 Distributed Algorithm for Nonlinear Programming
 - Exact Jacobi Successive Convex Approximation
 - Extensions

EM Algorithm

- Assume the complete data set $\{\mathbf{x}, \mathbf{z}\}$ consists of observed variable \mathbf{x} and latent variable \mathbf{z} .
- Objective: estimate parameter $\theta \in \Theta$ from \mathbf{x} .
- Maximum likelihood estimator: $\hat{\theta} = \arg \min_{\theta \in \Theta} -\log p(\mathbf{x}|\theta)$
- EM (Expectation Maximization) algorithm:
 - E-step: evaluate $p(\mathbf{z}|\mathbf{x}, \theta^k)$
“guess” \mathbf{z} from current estimate of θ
 - M-step: update θ as $\theta^{k+1} = \arg \min_{\theta \in \Theta} u(\theta, \theta^k)$, where

$$u(\theta, \theta^k) = -\mathbb{E}_{\mathbf{z}|\mathbf{x}, \theta^k} \log p(\mathbf{x}, \mathbf{z}|\theta)$$

update θ from “guessed” complete data set

An MM Interpretation of EM

- The objective function can be written as

$$\begin{aligned}
 & -\log p(\mathbf{x}|\theta) \\
 &= -\log E_{\mathbf{z}|\theta} p(\mathbf{x}|\mathbf{z}, \theta) \\
 &= -\log E_{\mathbf{z}|\theta} \left(\frac{p(\mathbf{z}|\mathbf{x}, \theta^k) p(\mathbf{x}|\mathbf{z}, \theta)}{p(\mathbf{z}|\mathbf{x}, \theta^k)} \right) \\
 &= -\log E_{\mathbf{z}|\mathbf{x}, \theta^k} \left(\frac{p(\mathbf{x}|\mathbf{z}, \theta)}{p(\mathbf{z}|\mathbf{x}, \theta^k)} p(\mathbf{z}|\theta) \right) \\
 &\leq -E_{\mathbf{z}|\mathbf{x}, \theta^k} \log \left(\frac{p(\mathbf{x}|\mathbf{z}, \theta)}{p(\mathbf{z}|\mathbf{x}, \theta^k)} p(\mathbf{z}|\theta) \right) \quad (\text{Jensen's Inequality}) \\
 &= \underbrace{-E_{\mathbf{z}|\mathbf{x}, \theta^k} \log p(\mathbf{x}, \mathbf{z}|\theta)}_{u(\theta, \theta^k)} + E_{\mathbf{z}|\mathbf{x}, \theta^k} p(\mathbf{z}|\mathbf{x}, \theta^k)
 \end{aligned}$$

Proximal Minimization

- $f(\mathbf{x})$ is convex. Solve $\min_{\mathbf{x}} f(\mathbf{x})$ by solving the equivalent problem

$$\underset{\mathbf{x} \in \mathcal{X}, \mathbf{y} \in \mathcal{X}}{\text{minimize}} \quad f(\mathbf{x}) + \frac{1}{2c} \|\mathbf{x} - \mathbf{y}\|^2.$$

- Objective function is strongly convex in both \mathbf{x} and \mathbf{y} .
- Algorithm:

$$\begin{aligned} \mathbf{x}^{k+1} &= \arg \min_{\mathbf{x} \in \mathcal{X}} \left\{ f(\mathbf{x}) + \frac{1}{2c} \|\mathbf{x} - \mathbf{y}^k\|^2 \right\} \\ \mathbf{y}^{k+1} &= \mathbf{x}^{k+1}. \end{aligned}$$

- An MM interpretation:

$$\mathbf{x}^{k+1} = \arg \min_{\mathbf{x} \in \mathcal{X}} \left\{ f(\mathbf{x}) + \frac{1}{2c} \|\mathbf{x} - \mathbf{x}^k\|^2 \right\}$$

DC Programming

- Consider the unconstrained problem

$$\underset{\mathbf{x} \in \mathbb{R}^n}{\text{minimize}} \quad f(\mathbf{x}),$$

where $f(\mathbf{x}) = g(\mathbf{x}) + h(\mathbf{x})$ with $g(\mathbf{x})$ convex and $h(\mathbf{x})$ concave.

- DC (Difference of Convex) Programming generates $\{\mathbf{x}^k\}$ by solving

$$\nabla g(\mathbf{x}^{k+1}) = -\nabla h(\mathbf{x}^k).$$

- An MM interpretation:

$$\mathbf{x}^{k+1} = \arg \min_{\mathbf{x}} \left\{ g(\mathbf{x}) + \nabla h(\mathbf{x}^k)^T (\mathbf{x} - \mathbf{x}^k) \right\}.$$

- 1 The Majorization-Minimization Algorithm
 - Introduction
 - Construction Techniques
 - Example Algorithms
 - Applications
- 2 Block Successive Majorization-Minimization
 - Introduction
 - Block Coordinate Descent
 - Block Successive Majorization-Minimization
 - Example Algorithms
 - Applications
- 3 Distributed Algorithm for Nonlinear Programming
 - Exact Jacobi Successive Convex Approximation
 - Extensions

Power Control by GP

- [Chi-Tan-Pal-O'Ne-Jul'J07] Problem: maximize system throughput. Essentially we need to solve the following problem:

$$\underset{\mathbf{P} \in \mathcal{P}}{\text{minimize}} \quad \frac{\sum_{j \neq i} G_{ij} P_j + n_i}{\sum_j G_{ij} P_j + n_i}$$

- Objective function is the ratio of two posynomials.
- Minorize a posynomial, denoted by $g(\mathbf{x}) = \sum_i m_i(\mathbf{x})$, by monomial:

$$g(\mathbf{x}) \geq \prod_i \left(\frac{m_i(\mathbf{x})}{\alpha_i} \right)^{\alpha_i}$$

where $\alpha_i = \frac{m_i(\mathbf{x}^k)}{g(\mathbf{x}^k)}$. (Arithmetic-Geometric Mean Inequality)

- Solution: approximate the denominator posynomial $\sum_j G_{ij} P_j + n_i$ by monomial.

Reweighted ℓ_1 -norm

- Sparsity signal recovery problem

$$\begin{array}{ll}\underset{\mathbf{x}}{\text{minimize}} & \|\mathbf{x}\|_0 \\ \text{subject to} & \mathbf{Ax} = \mathbf{b}\end{array}$$

- ℓ_1 -norm approximation


$$\begin{array}{ll}\underset{\mathbf{x}}{\text{minimize}} & \|\mathbf{x}\|_1 \\ \text{subject to} & \mathbf{Ax} = \mathbf{b}\end{array}$$

- General form

$$\begin{array}{ll}\underset{\mathbf{x}}{\text{minimize}} & \sum_{i=1}^n \phi(|x_i|) \\ \text{subject to} & \mathbf{Ax} = \mathbf{b}\end{array}$$

- [Can-Wak-Boy'J08] Assume $\phi(t)$ is concave nondecreasing, at x_i^k , $\phi(|x_i|)$ is majorized by $w_i^k |x_i|$ with $w_i^k = \phi'(t)|_{t=|x_i|}$.
- At each iteration a weighted ℓ_1 -norm is solved

$$\begin{array}{ll} \underset{\mathbf{x}}{\text{minimize}} & \sum w_i^k |x_i| \\ \text{subject to} & \mathbf{Ax} = \mathbf{b} \end{array}$$



Sparse Generalized Eigenvalue Problem

- ℓ_0 -norm regularized generalized eigenvalue problem

$$\begin{aligned} & \underset{\mathbf{x}}{\text{maximize}} && \mathbf{x}^T \mathbf{A} \mathbf{x} - \rho \|\mathbf{x}\|_0 \\ & \text{subject to} && \mathbf{x}^T \mathbf{B} \mathbf{x} = 1. \end{aligned}$$

- Replace $\|x_i\|_0$ by some nicely behaved function $g_p(x_i)$
 - $|x_i|^p$, $0 < p \leq 1$
 - $\log(1 + |x_i|/p) / \log(1 + 1/p)$, $p > 0$
 - $1 - e^{-|x_i|/p}$, $p > 0$.
- Take $g_p(x_i) = |x_i|^p$ for example.

- [Son-Bab-Pal'J15a] Majorize $g_p(x_i)$ at x_i^k by quadratic function $w_i^k x_i^2 + c_i^k$.
- The surrogate function for $g_p(x_i) = |x_i|^p$ is defined as

$$u(x_i, x_i^k) = \frac{p}{2} |x_i^k|^{p-2} x_i^2 + \left(1 - \frac{p}{2}\right) |x_i^k|^p.$$

- Solve at each iteration the following GEVP:


$$\begin{aligned} & \underset{\mathbf{x}}{\text{maximize}} && \mathbf{x}^T \mathbf{A} \mathbf{x} - \rho \mathbf{x}^T \text{diag}(\mathbf{w}^k) \mathbf{x} \\ & \text{subject to} && \mathbf{x}^T \mathbf{B} \mathbf{x} = 1 \end{aligned}$$

- However, as $|x_i| \rightarrow 0$, $w_i \rightarrow +\infty \dots$

- Smooth approximation of $g_p(x)$:

$$g_p^\varepsilon(x) = \begin{cases} \frac{p}{2} \varepsilon^{p-2} x^2, & |x| \leq \varepsilon \\ |x|^p - \left(1 - \frac{p}{2}\right) \varepsilon^p, & |x| > \varepsilon \end{cases}$$

- When $|x| \leq \varepsilon$, w remains to be a constant.



Sequence Design

- Complex unimodular sequence $\{x_n \in \mathbb{C}\}_{n=1}^N$.
- Autocorrelation: $r_k = \sum_{n=k+1}^N x_n x_{n-k}^* = r_{-k}^*$, $k = 0, \dots, N-1$.
- Integrated sidelobe level (ISL):

$$\text{ISL} = \sum_{k=1}^{N-1} |r_k|^2.$$

- Problem formulation:

$$\begin{array}{ll} \text{minimize} & \text{ISL} \\ & \{x_n\}_{n=1}^N \\ \text{subject to} & |x_n| = 1, \quad n = 1, \dots, N. \end{array}$$

- By Fourier transform:

$$\text{ISL} \propto \sum_{p=1}^{2N} \left[\left| \mathbf{a}_p^H \mathbf{x} \right|^2 - N \right]^2$$

with $\mathbf{x} = [x_1, \dots, x_N]^T$, $\mathbf{a}_p = [1, e^{j\omega_p}, \dots, e^{j\omega_p(N-1)}]^T$ and $\omega_p = \frac{2\pi}{2N}(p-1)$.

- Equivalent problem:

$$\begin{aligned} & \underset{\mathbf{x}}{\text{minimize}} && \sum_{p=1}^{2N} (\mathbf{a}_p^H \mathbf{x} \mathbf{x}^H \mathbf{a}_p)^2 \\ & \text{subject to} && |x_n| = 1, \forall n. \end{aligned}$$

- [Son-Bab-Pal'J15b] Define $\mathbf{A} = [\mathbf{a}_1, \dots, \mathbf{a}_{2N}]$,
 $\mathbf{p}^k = \left[|\mathbf{a}_1^H \mathbf{x}^k|^2, \dots, |\mathbf{a}_{2N}^H \mathbf{x}^k|^2 \right]^T$, $\tilde{\mathbf{A}} = \mathbf{A} (\text{diag}(\mathbf{p}^k) - p_{\max}^k \mathbf{I}) \mathbf{A}^H$.
- Quadratic surrogate function:

$$\underbrace{p_{\max}^k \mathbf{x}^H \mathbf{A} \mathbf{A}^H \mathbf{x}}_{\text{const.}} + 2\text{Re} \left(\mathbf{x}^H \left(\tilde{\mathbf{A}} - 2N^2 \mathbf{x}^k (\mathbf{x}^k)^H \right) \mathbf{x}^k \right)$$


- Equivalent to

$$\begin{aligned} & \underset{\mathbf{x}}{\text{minimize}} && \|\mathbf{x} - \mathbf{y}\|_2 \\ & \text{subject to} && |x_n| = 1, \forall n \end{aligned}$$

with

$$\mathbf{y} = - \left(\tilde{\mathbf{A}} - 2N^2 \mathbf{x}^k (\mathbf{x}^k)^H \right) \mathbf{x}^k$$

- Closed-form solution: $x_n = e^{j \arg(y_n)}$.



Covariance Estimation

- $\mathbf{x}_i \sim \text{elliptical}(\mathbf{0}, \mathbf{\Sigma})$
- Fitting normalized sample $\mathbf{s}_i = \frac{\mathbf{x}_i}{\|\mathbf{x}_i\|_2}$ to Angular Central Gaussian distribution

$$f(\mathbf{s}_i) \propto \det(\mathbf{\Sigma})^{-1/2} \left(\mathbf{s}_i^T \mathbf{\Sigma}^{-1} \mathbf{s}_i \right)^{-K/2}$$

- [Sun-Bab-Pal' J14] Shrinkage penalty

$$h(\mathbf{\Sigma}) = \log \det(\mathbf{\Sigma}) + \text{Tr}(\mathbf{\Sigma}^{-1} \mathbf{T})$$

- Solve the following problem:


$$\begin{aligned} & \underset{\mathbf{\Sigma}}{\text{minimize}} && \log \det(\mathbf{\Sigma}) + \frac{K}{N} \sum \log(\mathbf{x}_i^T \mathbf{\Sigma}^{-1} \mathbf{x}_i) + \alpha h(\mathbf{\Sigma}) \\ & \text{subject to} && \mathbf{\Sigma} \succeq \mathbf{0} \end{aligned}$$

- At Σ^k , the objective function is majorized by

$$(1 + \alpha) \log \det(\Sigma) + \frac{K}{N} \sum_{i=1}^N \frac{\mathbf{x}_i^T \Sigma^{-1} \mathbf{x}_i}{\mathbf{x}_i^T (\Sigma^k)^{-1} \mathbf{x}_i} + \alpha \text{Tr}(\Sigma^{-1} \mathbf{T})$$

- Surrogate function is convex in Σ^{-1} .
- Setting the gradient to zero leads to the weighted sample average

$$\Sigma^{k+1} = \frac{1}{1 + \alpha} \frac{K}{N} \sum \frac{\mathbf{x}_i \mathbf{x}_i^T}{\mathbf{x}_i^T (\Sigma^k)^{-1} \mathbf{x}_i} + \frac{\alpha}{1 + \alpha} \mathbf{T}$$



- 1 The Majorization-Minimization Algorithm
 - Introduction
 - Construction Techniques
 - Example Algorithms
 - Applications
- 2 Block Successive Majorization-Minimization
 - Introduction
 - Block Coordinate Descent
 - Block Successive Majorization-Minimization
 - Example Algorithms
 - Applications
- 3 Distributed Algorithm for Nonlinear Programming
 - Exact Jacobi Successive Convex Approximation
 - Extensions

Problem Statement

- Consider the following problem

$$\underset{\mathbf{x} \in \mathcal{X}}{\text{minimize}} \quad f(\mathbf{x})$$

- Set \mathcal{X} possesses Cartesian product structure $\mathcal{X} = \prod_{i=1}^m \mathcal{X}_i$.
- Observation: the problem

$$\underset{\mathbf{x}_i \in \mathcal{X}_i}{\text{minimize}} \quad f(\mathbf{x}_1^0, \dots, \mathbf{x}_{i-1}^0, \mathbf{x}_i, \mathbf{x}_{i+1}^0, \dots, \mathbf{x}_m^0)$$

with \mathbf{x}_{-i}^0 taking some feasible value, is easy to solve.

- 1 The Majorization-Minimization Algorithm
 - Introduction
 - Construction Techniques
 - Example Algorithms
 - Applications
- 2 Block Successive Majorization-Minimization
 - Introduction
 - **Block Coordinate Descent**
 - Block Successive Majorization-Minimization
 - Example Algorithms
 - Applications
- 3 Distributed Algorithm for Nonlinear Programming
 - Exact Jacobi Successive Convex Approximation
 - Extensions

Block Coordinate Descent (BCD)

- Denote $\mathbf{x} \triangleq (\mathbf{x}_1, \dots, \mathbf{x}_m)$,
 $f(\mathbf{x}_1^0, \dots, \mathbf{x}_{i-1}^0, \mathbf{x}_i, \mathbf{x}_{i+1}^0, \dots, \mathbf{x}_m^0) \triangleq f(\mathbf{x}_i, \mathbf{x}^0)$
- Block Coordinate Descent (nonlinear Gauss-Seidel)
 - 1: Initialize $\mathbf{x}^0 \in \mathcal{X}$ and set $k = 0$.
 - 2: **repeat**
 - 3: $k = k + 1, i = (k \bmod n) + 1$
 - 4: $\mathbf{x}_i^k = \arg \min_{\mathbf{x}_i \in \mathcal{X}_i} f(\mathbf{x}_i, \mathbf{x}^{k-1})$
 - 5: $\mathbf{x}_i^k \leftarrow \mathbf{x}_i^{k-1}, \forall k \neq i$
 - 6: **until** some convergence criterion is met

Convergence

- [Ber'B99] Assume that
 - $f(\mathbf{x})$ is **continuously differentiable** over the set \mathcal{X} .
 - $\mathbf{x}_i^k = \arg \min_{\mathbf{x}_i \in \mathcal{X}_i} f(\mathbf{x}_i, \mathbf{x}^{k-1})$ has a **unique** solution.

Then every limit point of the sequence $\{\mathbf{x}^k\}$ is a stationary point.

- [Gri-Sci'J00] Generalizations
 - globally convergent for $m = 2$.
 - f is component-wise strictly quasi-convex w.r.t. $m - 2$ components.
 - f is pseudo-convex.

- 1 The Majorization-Minimization Algorithm
 - Introduction
 - Construction Techniques
 - Example Algorithms
 - Applications
- 2 Block Successive Majorization-Minimization
 - Introduction
 - Block Coordinate Descent
 - **Block Successive Majorization-Minimization**
 - Example Algorithms
 - Applications
- 3 Distributed Algorithm for Nonlinear Programming
 - Exact Jacobi Successive Convex Approximation
 - Extensions

BS-MM Algorithm

- Combination of MM and BCD
- Block Successive Majorization-Minimization (BS-MM):
 - 1: Initialize $\mathbf{x}^0 \in \mathcal{X}$ and set $k = 0$.
 - 2: **repeat**
 - 3: $k = k + 1, i = (k \bmod n) + 1$
 - 4: $\mathcal{X}^k = \arg \min_{\mathbf{x}_i \in \mathcal{X}_i} u_i(\mathbf{x}_i, \mathbf{x}^{k-1})$
 - 5: Set \mathbf{x}_i^k to be an arbitrary element in \mathcal{X}^k
 - 6: $\mathbf{x}_i^k \leftarrow \mathbf{x}_i^{k-1}, \forall k \neq i$
 - 7: **until** some convergence criterion is met
- Generalization of BCD

Convergence

- Surrogate function $u_i(\cdot, \cdot)$ satisfies the following assumptions

$$u_i(\mathbf{y}_i, \mathbf{y}) = f(\mathbf{y}), \quad \forall \mathbf{y} \in \mathcal{X}, \forall i \quad (\text{B1})$$

$$u_i(\mathbf{x}_i, \mathbf{y}) \geq f(\mathbf{y}_1, \dots, \mathbf{y}_{i-1}, \mathbf{x}_i, \mathbf{y}_{i+1}, \dots, \mathbf{y}_n),$$

$$\forall \mathbf{x}_i \in \mathcal{X}_i, \forall \mathbf{y} \in \mathcal{X}, \forall i \quad (\text{B2})$$

$$u'_i(\mathbf{x}_i, \mathbf{y}; \mathbf{d}_i) \big|_{\mathbf{x}_i = \mathbf{y}_i} = f'(\mathbf{y}; \mathbf{d}),$$

$$\forall \mathbf{d} = (\mathbf{0}, \dots, \mathbf{d}_i, \dots, \mathbf{0}) \text{ such that } \mathbf{y}_i + \mathbf{d}_i \in \mathcal{X}_i, \forall i \quad (\text{B3})$$

$$u_i(\mathbf{x}_i, \mathbf{y}) \text{ is continuous in } (\mathbf{x}_i, \mathbf{y}), \quad \forall i \quad (\text{B4})$$

- In short, $u_i(\mathbf{x}_i, \mathbf{x}^k)$ majorizes $f(\mathbf{x})$ on the i th block.

- Under assumptions B1-B4, for simplicity additionally assume that f is continuously differentiable,
 - $u_i(\mathbf{x}_i, \mathbf{y})$ is **quasi-convex** in \mathbf{x}_i ,
each subproblem $\min_{\mathbf{x}_i \in \mathcal{X}_i} u_i(\mathbf{x}_i, \mathbf{x}^{k-1})$ has a **unique** solution for any $\mathbf{x}^{k-1} \in \mathcal{X}$,
then every limit point of $\{\mathbf{x}^k\}$ is a stationary point.
 - level set $\mathcal{X}^0 = \{\mathbf{x} | f(\mathbf{x}) \leq f(\mathbf{x}^0)\}$ is **compact**,
each subproblem $\min_{\mathbf{x}_i \in \mathcal{X}_i} u_i(\mathbf{x}_i, \mathbf{x}^{k-1})$ has a **unique** solution for any $\mathbf{x}^{k-1} \in \mathcal{X}$ for at least $m-1$ blocks,
then $\lim_{k \rightarrow \infty} d(\mathbf{x}^k, \mathcal{X}^*) = 0$.
- More restrictive assumption than MM due to the cyclic update behavior.

- 1 The Majorization-Minimization Algorithm
 - Introduction
 - Construction Techniques
 - Example Algorithms
 - Applications
- 2 Block Successive Majorization-Minimization
 - Introduction
 - Block Coordinate Descent
 - Block Successive Majorization-Minimization
 - **Example Algorithms**
 - Applications
- 3 Distributed Algorithm for Nonlinear Programming
 - Exact Jacobi Successive Convex Approximation
 - Extensions

Alternating Proximal Minimization

- Consider the problem

$$\begin{array}{ll}\underset{\mathbf{x}}{\text{minimize}} & f(\mathbf{x}_1, \dots, \mathbf{x}_m) \\ \text{subject to} & \mathbf{x}_i \in \mathcal{X}_i,\end{array}$$

with $f(\cdot)$ being convex in each block.

- The convergence of BCD is not easy to establish since each subproblem may have multiple solutions.
- Alternating Proximal Minimization solves

$$\begin{array}{ll}\underset{\mathbf{x}_i}{\text{minimize}} & f(\mathbf{x}_1^k, \dots, \mathbf{x}_{i-1}^k, \mathbf{x}_i, \mathbf{x}_{i+1}^k, \dots, \mathbf{x}_m^k) + \frac{1}{2c} \|\mathbf{x}_i - \mathbf{x}_i^k\|^2 \\ \text{subject to} & \mathbf{x}_i \in \mathcal{X}_i\end{array}$$

- Strictly convex objective \rightarrow unique minimizer

Proximal Splitting Algorithm

- Consider the following problem

$$\begin{array}{ll}\underset{\mathbf{x}}{\text{minimize}} & \sum_{i=1}^m f_i(\mathbf{x}_i) + f_{m+1}(\mathbf{x}_1, \dots, \mathbf{x}_m) \\ \text{subject to} & \mathbf{x}_i \in \mathcal{X}_i, \quad i = 1, \dots, m\end{array}$$

with f_i convex and lower semicontinuous, f_{m+1} convex and

$$\|\nabla f_{m+1}(\mathbf{x}) - \nabla f_{m+1}(\mathbf{y})\| \leq \beta_i \|\mathbf{x}_i - \mathbf{y}_i\|$$

- Cyclically update:

$$\mathbf{x}_i^{k+1} = \text{prox}_{\gamma f_i} \left(\mathbf{x}_i^k - \gamma \nabla_{\mathbf{x}_i} f_{m+1}(\mathbf{x}^k) \right),$$

with the proximity operator defined as

$$\text{prox}_f(\mathbf{x}) = \arg \min_{\mathbf{y} \in \mathcal{X}} f(\mathbf{y}) + \frac{1}{2} \|\mathbf{x} - \mathbf{y}\|^2.$$

Proximal Splitting Algorithm

- BS-MM interpretation:

$$u_i(\mathbf{x}_i, \mathbf{x}^k) = f_i(\mathbf{x}_i) + \frac{1}{2\gamma} \|\mathbf{x}_i - \mathbf{x}_i^k\|^2 + \nabla_{\mathbf{x}_i} f_{m+1}(\mathbf{x}^k)^T (\mathbf{x}_i - \mathbf{x}_i^k) \\ + \sum_{j \neq i} f_j(\mathbf{x}_j^k) + f_{m+1}(\mathbf{x}_{-i}^k, \mathbf{x}_i).$$

- Check:

$$f_{m+1}(\mathbf{x}^k) + \frac{1}{2\gamma} \|\mathbf{x}_i - \mathbf{x}_i^k\|^2 + \nabla_{\mathbf{x}_i} f_{m+1}(\mathbf{x}^k)^T (\mathbf{x}_i - \mathbf{x}_i^k) \\ \geq f_{m+1}(\mathbf{x}^k) + \frac{\beta_i}{2} \|\mathbf{x}_i - \mathbf{x}_i^k\|^2 + \nabla_{\mathbf{x}_i} f_{m+1}(\mathbf{x}^k)^T (\mathbf{x}_i - \mathbf{x}_i^k) \\ \geq f_{m+1}(\mathbf{x}_{-i}^k, \mathbf{x}_i) \quad (\text{Descent lemma})$$

with $\gamma \in [\varepsilon_i, 2/\beta_i - \varepsilon_i]$ and $\varepsilon_i \in (0, \min\{1, 1/\beta_i\})$.

- 1 The Majorization-Minimization Algorithm
 - Introduction
 - Construction Techniques
 - Example Algorithms
 - Applications
- 2 Block Successive Majorization-Minimization
 - Introduction
 - Block Coordinate Descent
 - Block Successive Majorization-Minimization
 - Example Algorithms
 - Applications
- 3 Distributed Algorithm for Nonlinear Programming
 - Exact Jacobi Successive Convex Approximation
 - Extensions

Robust Estimation of Location and Scatter

- $\mathbf{x}_i \sim \text{elliptical}(\boldsymbol{\mu}, \mathbf{R})$
- [Sun-Bab-Pal'J15] Fitting \mathbf{x}_i to a Cauchy distribution with pdf

$$f(\mathbf{x}) \propto \det(\mathbf{R})^{-1/2} \left(1 + (\mathbf{x}_i - \boldsymbol{\mu})^T \mathbf{R}^{-1} (\mathbf{x}_i - \boldsymbol{\mu}) \right)^{-(K+1)/2}$$

- Shrinkage penalty

$$h(\mathbf{t}, \mathbf{T}) = K \log(\text{Tr}(\mathbf{R}^{-1} \mathbf{T})) + \log \det(\mathbf{R}) + \log \left(1 + (\mathbf{t} - \boldsymbol{\mu})^T \mathbf{R}^{-1} (\mathbf{t} - \boldsymbol{\mu}) \right)$$

- Solve the following problem:

$$\begin{aligned} \underset{\boldsymbol{\mu}, \mathbf{R} \succeq \mathbf{0}}{\text{minimize}} \quad & \log \det(\mathbf{R}) + \frac{K+1}{N} \sum_{i=1}^N \log \left(1 + (\mathbf{x}_i - \boldsymbol{\mu})^T \mathbf{R}^{-1} (\mathbf{x}_i - \boldsymbol{\mu}) \right) \\ & + \alpha h(\mathbf{t}, \mathbf{T}) \end{aligned}$$


- BS-MM Algorithm update:

$$\mu_{t+1} = \frac{(K+1) \sum_{i=1}^N w_i(\mu_t, \mathbf{R}_t) \mathbf{x}_i + N\alpha w_t(\mu_t, \mathbf{R}_t) \mathbf{t}}{(K+1) \sum_{i=1}^N w_i(\mu_t, \mathbf{R}_t) + N\alpha w_t(\mu_t, \mathbf{R}_t)}$$

$$\begin{aligned} \mathbf{R}_{t+1} = & \frac{K+1}{N+N\alpha} \sum_{i=1}^N w_i(\mu_{t+1}, \mathbf{R}_t) (\mathbf{x}_i - \mu_{t+1}) (\mathbf{x}_i - \mu_{t+1})^T \\ & + \frac{N\alpha}{N+N\alpha} w_t(\mu_{t+1}, \mathbf{R}_t) (\mu_{t+1} - \mathbf{t}) (\mu_{t+1} - \mathbf{t})^T \\ & + \frac{N\alpha K}{N+N\alpha} \frac{\mathbf{T}}{\text{Tr}(\mathbf{R}_t^{-1} \mathbf{T})} \end{aligned}$$

where

$$\begin{aligned} w_i(\mu, \mathbf{R}) &= \frac{1}{1 + (\mathbf{x}_i - \mu)^T \mathbf{R}^{-1} (\mathbf{x}_i - \mu)} \\ w_t(\mu, \mathbf{R}) &= \frac{1}{1 + (\mathbf{t} - \mu)^T \mathbf{R}^{-1} (\mathbf{t} - \mu)}. \end{aligned}$$



- 1 The Majorization-Minimization Algorithm
 - Introduction
 - Construction Techniques
 - Example Algorithms
 - Applications
- 2 Block Successive Majorization-Minimization
 - Introduction
 - Block Coordinate Descent
 - Block Successive Majorization-Minimization
 - Example Algorithms
 - Applications
- 3 Distributed Algorithm for Nonlinear Programming
 - Exact Jacobi Successive Convex Approximation
 - Extensions

Problem Statement

- Consider the following problem:

$$\begin{array}{ll} \underset{\mathbf{x}}{\text{minimize}} & f(\mathbf{x}) \\ \text{subject to} & \mathbf{x}_i \in \mathcal{X}_i \end{array}$$

where the \mathcal{X}_i 's are closed and convex sets,
 $f(\mathbf{x}) = \sum_{l=1}^L f_l(\mathbf{x}_1, \dots, \mathbf{x}_m)$.

- Conditional gradient update (Frank-Wolfe):

$$\mathbf{x}^{k+1} = \mathbf{x}^k + \gamma^k \mathbf{d}^k$$

- direction $\mathbf{d}^k \triangleq \bar{\mathbf{x}}^k - \mathbf{x}^k$ with

$$\bar{\mathbf{x}}_i^k = \arg \min_{\mathbf{x}_i \in \mathcal{X}_i} \nabla_{\mathbf{x}_i} f(\mathbf{x}^k)^T (\mathbf{x}_i - \mathbf{x}_i^k)$$

- step-size $\gamma^k \in (0, 1]$, chosen to guarantee convergence.

Exact Jacobi SCA Algorithm

- Idea:
 - Conditional gradient update linearize all the f_l 's at \mathbf{x}^k .
 - Each function f_l might be convex w.r.t. some block \mathbf{x}_i .
 - We want to preserve the convex property of $f_l(\mathbf{x}_i, \mathbf{x}_{-i}^k)$.
- Solution: keep the convex $f_l(\mathbf{x}_i, \mathbf{x}_{-i}^k)$'s and linearize the others.

- Define \mathcal{C}_i as the set of indices of l such that $f_l(\mathbf{x}_i, \mathbf{x}_{-i}^k)$ is convex.
- Approximate $f(\mathbf{x})$ on the i th block at point \mathbf{x}^k :

$$\begin{aligned} \tilde{f}_i(\mathbf{x}_i, \mathbf{x}^k) = & \underbrace{\sum_{l \in \mathcal{C}_i} f_l(\mathbf{x}_i, \mathbf{x}_{-i}^k)}_{\text{convex terms}} + \underbrace{\pi_i(\mathbf{x}^k)^T (\mathbf{x}_i - \mathbf{x}_i^k)}_{\substack{\text{linear approx.} \\ \text{non-convex terms}}} \\ & + \underbrace{\frac{\tau_i}{2} (\mathbf{x}_i - \mathbf{x}_i^k)^T \mathbf{H}_i(\mathbf{x}^k) (\mathbf{x}_i - \mathbf{x}_i^k)}_{\text{proximal term}}, \end{aligned}$$

with

$$\pi_i(\mathbf{x}^k) = \sum_{l \notin \mathcal{C}_i} \nabla_{\mathbf{x}_i} f_l(\mathbf{x}^k) \text{ and } \mathbf{H}_i(\mathbf{x}^k) \succ c_{H_i} \mathbf{I}.$$

- Exact Jacobi SCA update:

$$\hat{\mathbf{x}}_i(\mathbf{x}^k, \tau_i) = \arg \min_{\mathbf{x}_i \in \mathcal{X}_i} \tilde{f}_i(\mathbf{x}_i, \mathbf{x}^k)$$

$$\mathbf{x}^{k+1} = \mathbf{x}^k + \gamma^k (\hat{\mathbf{x}} - \mathbf{x}^k)$$

- Step-size rule
 - constant step-size that depends on the Lipschitz constant of ∇f
 - diminishing step-size
- Remark: update of the blocks can be done sequentially (Gauss-Seidel SCA Algorithm)

- 1 The Majorization-Minimization Algorithm
 - Introduction
 - Construction Techniques
 - Example Algorithms
 - Applications
- 2 Block Successive Majorization-Minimization
 - Introduction
 - Block Coordinate Descent
 - Block Successive Majorization-Minimization
 - Example Algorithms
 - Applications
- 3 Distributed Algorithm for Nonlinear Programming
 - Exact Jacobi Successive Convex Approximation
 - Extensions

Extensions

- FLEXA
 - non-smooth objective function
 - inexact update direction
 - flexible block update choice
- HyFLEXA

Comparison

	BS-MM	FLEXA
convergence	stationary point	stationary point
objective function	continuous may not be smooth	continuous may not be smooth
constraint set	Cartesian	Cartesian & convex
update rule	sequential	sequential or parallel
approx. function	global upper-bound unique minimizer can be non-convex	local approximation not required convex approx.

Summary

- We have studied
 - Majorization-Minimization algorithm
 - Block Coordinate Descent algorithm
 - Block Successive Majorization-Minimization algorithm
- We have briefly introduced
 - Distributed Successive Convex Approximation algorithm

References I



D. R. Hunter and K. Lange.

A tutorial on MM algorithms.

Amer. Statistician, volume 58, pp. 30–37, 2004.



M. Razaviyayn, M. Hong, and Z. Luo.

A unified convergence analysis of block successive minimization methods for nonsmooth optimization.

SIAM J. Optim., volume 23, no. 2, pp. 1126–1153, 2013.



G. Scutari, F. Facchinei, P. Song, D. Palomar, and J.-S. Pang.

Decomposition by partial linearization: Parallel optimization of multi-agent systems.

IEEE Trans. Signal Process., volume 62, no. 3, pp. 641–656, 2014.



Y. Sun, P. Babu, and D. Palomar.

Majorization-minimization algorithms in signal processing, communications, and machine learning.

IEEE Transactions on Signal Processing, volume 65, no. 3, pp. 794–816, 2017.



M. Chiang, C. W. Tan, D. Palomar, D. O'Neill, and D. Julian.

Power control by geometric programming.

IEEE Trans. Wireless Commun., volume 6, no. 7, pp. 2640–2651, 2007.

References II



E. J. Candes, M. Wakin, and S. Boyd.
Enhancing sparsity by reweighted l1 minimization.
J. Fourier Anal. Appl., volume 14, no. 5-6, pp. 877–905, 2008.



J. Song, P. Babu, and D. P. Palomar.
Sparse generalized eigenvalue problem via smooth optimization.
IEEE Trans. Signal Process., volume 63, no. 7, pp. 1627–1642, 2015.



—.
Optimization methods for designing sequences with low autocorrelation sidelobes.
IEEE Trans. Signal Process., volume 63, no. 15, pp. 3998–4009, 2015.



Y. Sun, P. Babu, and D. P. Palomar.
Regularized Tyler's scatter estimator: Existence, uniqueness, and algorithms.
IEEE Trans. Signal Processing, volume 62, no. 19, pp. 5143–5156, 2014.



D. P. Bertsekas.
Nonlinear Programming.
Athena Scientific, 1999.



L. Grippo and M. Sciandrone.
On the convergence of the block nonlinear gauss–seidel method under convex constraints.
Oper. Res. Lett., volume 26, no. 3, pp. 127–136, 2000.



Y. Sun, P. Babu, and D. P. Palomar.

Regularized robust estimation of mean and covariance matrix under heavy-tailed distributions.

IEEE Transactions on Signal Processing, volume 63, no. 12, pp. 3096–3109, 2015.

Thanks

For more information visit:

<http://www.danielppalomar.com>

