TYPE II FINITE TIME BLOW-UP FOR THE THREE DIMENSIONAL ENERGY CRITICAL HEAT EQUATION

MANUEL DEL PINO, MONICA MUSSO, JUNCHENG WEI, QIDI ZHANG, AND YIFU ZHOU

ABSTRACT. We consider the following Cauchy problem for three dimensional energy critical heat equation

$$\begin{cases} u_t = \Delta u + u^5, & \text{in } \mathbb{R}^3 \times (0, T), \\ u(x, 0) = u_0(x), & \text{in } \mathbb{R}^3. \end{cases}$$

We construct type II finite time blow-up solution u(x,t) with the blow-up rates $||u||_{L^{\infty}} \sim (T-t)^{-k}$, where k=1,2,... This gives a rigorous proof of the formal computations by Filippas, Herrero and Velazquez [14]. This is the first instance of type II finite time blow-up for three dimensional energy critical heat equation.

1. Introduction

The study of blow-up phenomena for the Fujita type nonlinear heat equation

$$\begin{cases} u_t = \Delta u + |u|^{p-1}u & \text{in } \mathbb{R}^n \times (0, T), \\ u(\cdot, 0) = u_0 & \text{in } \mathbb{R}^n \end{cases}$$
 (1.1)

is a classical topic with important applications in mathematical modelling and geometry. Here p > 1 and $n \ge 1$. A smooth solution of (1.1) blows-up at time T if

$$\lim_{t \to T} \|u(\cdot, t)\|_{L^{\infty}(\mathbb{R}^n)} = +\infty.$$

There are two types of blow-ups: the blow-up of a solution u(x,t) is of type I if it happens at most at the ODE rate:

$$\limsup_{t\to T} (T-t)^{\frac{1}{p-1}} \|u(\cdot,t)\|_{L^{\infty}(\mathbb{R}^n)} < +\infty$$

while the blow-up is said of type II if

$$\limsup_{t \to T} (T-t)^{\frac{1}{p-1}} \|u(\cdot,t)\|_{L^{\infty}(\mathbb{R}^n)} = +\infty.$$

It is known after a series of works, including [15–17], that type I is the only way possible if $p < p_S$ where p_S is the critical Sobolev exponent,

$$p_S := \begin{cases} \frac{n+2}{n-2} & \text{if } n \ge 3\\ +\infty & \text{if } n = 1, 2. \end{cases}$$

Stability and genericity of type I blow-up have been considered for instance in [4,30,32,34,35]. Solutions with type II blow-up are in fact much harder to detect. The first example was discovered in [20,21], for $p > p_{JL}$ where p_{JL} is the Joseph-Lundgreen exponent,

$$p_{JL} = \begin{cases} 1 + \frac{4}{n-4-2\sqrt{n-1}} & \text{if } n \ge 11\\ +\infty, & \text{if } n \le 10. \end{cases}$$

See the book [36] for a survey of related results. In fact, no type II blow-up is present for radial solutions if $p_S , while for radial positive solutions this is not possible if <math>p = \frac{n+2}{n-2}$ [14]. Examples of nonradial positive blow-up solutions for $p > p_{JL}$ have been found in [1,3]. Various different scenarios have been discovered or discarded in the supercritical case. See for instance [1,3,9,20,21,29,30,33] and the book [36].

In [14] Filippas, Herrero and Velazquez formally obtained sign-changing solution with type II blowup for $p = p_S$ in [14] in lower dimensions n = 3, 4, 5, 6. The blowup of radial blowup solutions found in [14] are given by

$$||u||_{L^{\infty}(\mathbb{R}^{n})} \sim \begin{cases} (T-t)^{-k}, & n=3, \\ (T-t)^{-k} |\log(T-t)|^{\frac{2k}{k-1}}, & n=4, \\ (T-t)^{-k}, & n=5, \\ (T-t)^{-\frac{1}{4}} |\log(T-t)|^{-\frac{15}{8}}, & n=6 \end{cases}$$
(1.2)

and this is corrected to

$$||u||_{L^{\infty}(\mathbb{R}^{n})} \sim \begin{cases} (T-t)^{-k}, & n=3, \\ (T-t)^{-k} |\log(T-t)|^{\frac{2k}{k-1}}, & n=4, \\ (T-t)^{-3k}, & n=5, \\ (T-t)^{-\frac{5}{2}} |\log(T-t)|^{-\frac{15}{4}}, & n=6 \end{cases}$$
(1.3)

by Harada [18].

The first rigorous proof of radial example was constructed by Schweyer in [40] for n = 4 and k = 1. (See [10] for multiple blow-ups and nonradial case in n = 4, k = 1.) There is a deep connection between dimension four energy critical and two-dimensional harmonic map flows ([6] [38]). Slower blowup rates for harmonic map flows are established by Raphael and Schweyer [39], which suggests that slower blowup rates are also possible for energy critical heat equation in the case $n = 4, k \ge 2$.

In dimension n=5 and critical case $p=p_S=\frac{7}{3}$, the first three authors gave a rigorous construction of type II blow-up in radial and nonradial cases (confirming the case of k=1, n=5). Higher speed blow-up solutions for $n=5, p=p_S=\frac{7}{3}, k\geq 2$ are carried out recently by Harada [18].

Very recently Harada succeeded in establishing the type II blow-up in dimension $n = 6, p = p_S = 2$ [19].

The purpose of this paper is to fill the gap in the last remaining dimension $n = 3, p = p_S = 5$.

Theorem 1. Let n = 3 and $k \in \mathbb{Z}_+$. For each T > 0 sufficiently small there exists an initial condition u_0 such that the solution of Problem (1.1) blows up at time T which looks like at main order

$$u(x,t) = \eta \left(\frac{x}{\sqrt{T-t}}\right) \left[\mu^{-\frac{1}{2}}(t)w\left(\frac{x}{\mu(t)}\right) + 2\mu'(t)\mu^{\frac{1}{2}}(t)J\left(\frac{x}{\mu(t)}\right)\right] + \left(1 - \eta\left(\frac{x}{\sqrt{T-t}}\right)\right) \eta\left(\frac{x}{\sqrt{T-t}}\right) (T-t)^k \frac{1}{|x|} C_k H_{2k}\left(\frac{|x|}{2\sqrt{T-t}}\right) + \theta(x,t)$$

where

$$w(y) = 3^{\frac{1}{4}} (1 + |y|^2)^{-\frac{1}{2}}, \quad C_k = \frac{(-1)^k k! \sqrt{3}}{(2k)!},$$

 H_{2k} is the Hermite polynomial defined in (2.3) and J is defined in (2.7). Moreover, the blow-up rate $\mu(t)$ satisfies

$$\mu(t) \sim \mu_0(t) = 3^{\frac{1}{2}} A (T - t)^{2k}, \quad k \in \mathbb{Z}_+$$

and $\|\theta\|_{L^{\infty}} < T^a$ for some a > 0.

The method of this paper is close in spirit to the analysis in the works [5–9], where the inner-outer gluing method is employed. This method reduces the original problem to solving a basically uncoupled system, which depends in subtle ways on the parameter choices (which are governed by relatively simple ODE systems). The main obstacle in proving finite time blow-up in dimension three is the very slow decaying behavior of the kernel $Z_0(y) \sim \frac{1}{|y|}$ which is not even in L^2 . To overcome this difficulty we follow the ideas in [7] in which the first three authors constructed infinite time blow-up for (1.1) with fast decaying initial condition. First we use the matched asymptotics expansion of [14] to construct a good inner and outer expansions. But this is not good enough as the error still carries slow decaying. As in [6] and [7], we use a global term to correct the slow decaying error. This global term carries all

the information needed to solve the scaling parameter. We then use the <u>inner-outer</u> gluing procedure to find a true solution. Interestingly the remainder of the scaling parameter solves a nonlocal ODE of the following form

$$\int_{t}^{T} \frac{\alpha(s)}{\sqrt{T-s}} ds = h(t), t < T, \alpha(T) = 0$$

which is Caputo derivative of $\frac{1}{2}$ for $\beta = \int_0^t \alpha(t)$. This is the <u>nonlocal feature</u> for three dimensional problem. As far as we know this seems to be the first construction of finite time blow-up for critical heat equation in \mathbb{R}^3 .

We remark that the phenomena of blow-up by bubbling (time dependent, energy invariant, asymptotically singular scalings of steady states) arises in various problems of parabolic, geometric and dispersive nature. This has been an intensively studied topic in the harmonic map flow [6, 27], critical heat equations [2, 8, 14, 29, 30, 36], critical Schrödinger maps [28, 31], critical wave maps and Yang-Mills [23, 37], and critical wave equations [12, 13, 22, 24–26].

The rest of this paper will be devoted to the proof of Theorem 1.

2. First approximation and matching

In this section, following the matched asymptotic expansions first developed in [14], we obtain an initial blow-up rate of finite blow-up solutions of the following nonlinear heat equation with critical exponent in \mathbb{R}^3 ,

$$u_t = \Delta u + u^5, \quad u(x,0) = u_0(x), \quad x \in \mathbb{R}^3, \quad t > 0.$$
 (2.1)

where the initial value u_0 will be determined later. Throughout the paper, we shall use the symbol " \lesssim " to denote " $\leq C$ " for a positive constant C independent of t and T, where C might be different from line to line.

2.1. Approximate solutions. In the self-similar variable

$$z = \frac{x}{\sqrt{T-t}}, \quad \tau = -\log(T-t), \quad \Phi(z,\tau) = (T-t)^{\frac{1}{4}}u(x,t),$$

problem (2.1) reads as

$$\Phi_{\tau} = \Delta \Phi - \frac{1}{2} (z \cdot \nabla \Phi) - \frac{\Phi}{4} + \Phi^{5}.$$

We next find good approximate solution for the above equation in both inner and outer regimes.

For the outer part, the first profile will be chosen as the solution to the linear problem

$$\partial_{\tau}\Phi_{out} = \Delta\Phi_{out} - \frac{1}{2}(z \cdot \nabla\Phi_{out}) - \frac{\Phi_{out}}{4}.$$
 (2.2)

Writing $\Phi_{out} = e^{\gamma \tau} m(z)$, we look for radially symmetric solutions to the following ODE

$$m'' + (\frac{2}{z} - \frac{1}{2}z)m' - (\gamma + \frac{1}{4})m = 0,$$

which turns out to be an eigenvalue problem ([14]). In order to get even solutions with polynomial growth rate in the outer regime, we take $\gamma = \frac{1}{4} - k$, $k \in \mathbb{Z}_+$, and then

$$m(z) = A^{\frac{1}{2}} C_k \frac{1}{z} H_{2k}(\frac{z}{2}),$$

where H_{2k} is Hermite polynomial

$$H_{2k}(\frac{z}{2}) = \frac{(-1)^k (2k)!}{k!} (1 - k\frac{z^2}{2} + \dots + a_k z^{2k}), \ C_k = \frac{(-1)^k k! \sqrt{3}}{(2k)!}, \tag{2.3}$$

 a_k is constant depending on k, and A is any positive constant. Therefore, we get special solutions of (2.2)

$$\Phi_{out}(z,\tau) = A^{\frac{1}{2}} e^{(\frac{1}{4}-k)\tau} C_k \frac{1}{z} H_{2k}(\frac{z}{2}),$$

which are sufficient for the matching to be carried out below, and we take the outer approximate solution of (2.1) to be

$$u_{out}(x,t) = (T-t)^{-\frac{1}{4}} \Phi_{out}(z,\tau) = A^{\frac{1}{2}} (T-t)^k \frac{1}{x} C_k H_{2k} (\frac{1}{2} \frac{x}{\sqrt{T-t}}). \tag{2.4}$$

It can be directly checked that in the original variable u_{out} satisfies

$$\partial_t u_{out} = \Delta u_{out}$$
.

For the inner part, we choose the approximate solution to be

$$\Phi_{in}(z,\tau) = \epsilon^{-1/2} (w(y) + \sigma(\tau)J(y)),$$

where

$$w(y) = 3^{\frac{1}{4}} (1 + |y|^2)^{-\frac{1}{2}}, \ y = \frac{z}{\epsilon}, \ \sigma(\tau) = 2\epsilon(\tau) \partial_{\tau} \epsilon(\tau) - \epsilon^2(\tau),$$

and $\epsilon(\tau)$ is a positive function to be determined later. Here J is the radial solution of

$$\Delta J + 5w^4 J + \frac{1}{2}y \cdot \nabla w + \frac{w}{4} = 0, \ J(0) = 0, \ J'(0) = 0.$$
 (2.5)

Set $\mu(t) := \sqrt{T - t} \epsilon(\tau)$ and $y = \frac{x}{\mu}$. Then one has

$$\bar{u}_{in}(x,t) = (T-t)^{-\frac{1}{4}} \Phi_{in}(z,\tau) = \mu^{-\frac{1}{2}} \left[w(\frac{x}{\mu}) + 2\mu \mu' J(\frac{x}{\mu}) \right]$$

since

$$\sigma = 2\epsilon \partial_{\tau} \epsilon - \epsilon^{2} = 2\epsilon \partial_{t} \epsilon (T - t) - \epsilon^{2}$$

$$= 2\mu (T - t)^{-\frac{1}{2}} \left[\mu' (T - t)^{-\frac{1}{2}} + \mu \frac{1}{2} (T - t)^{-\frac{3}{2}} \right] (T - t) - \mu^{2} (T - t)^{-1}$$

$$= 2\mu \mu'.$$

Denote

$$Z_0(y) := -\left[y \cdot \nabla w + \frac{w}{2}\right] = \frac{3^{\frac{1}{4}}}{2} \frac{|y|^2 - 1}{(1 + |y|^2)^{\frac{3}{2}}},\tag{2.6}$$

which is a radial kernel of the homogeneous part of (2.5). Then it is easy to see that

$$J(y) = Z_0(y) \int_0^y \frac{1}{Z_0^2(s)} s^{-2} \int_0^s Z_0(u) u^2 \frac{Z_0(u)}{2} du ds,$$
 (2.7)

and thus

$$J(y) \sim \frac{3^{\frac{1}{4}}}{8}y$$
 as $y \to \infty$.

2.2. Matching inner and outer solutions. In the region $\mu(t) \ll |x| \ll \sqrt{T-t}$, the inner and outer solutions have the following asymptotic behaviors respectively

$$\bar{u}_{in} \sim 3^{\frac{1}{4}} \mu^{\frac{1}{2}} |x|^{-1} + \frac{3^{\frac{1}{4}}}{4} \mu^{-\frac{1}{2}} \mu' |x|,$$
 (2.8)

$$u_{out} \sim A^{\frac{1}{2}} \sqrt{3} \frac{(T-t)^k}{|x|} - A^{\frac{1}{2}} \sqrt{3} \frac{k}{2} |x| (T-t)^{k-1}.$$
 (2.9)

Matching the inner and outer solutions, we get

$$\mu \sim 3^{\frac{1}{2}} A (T - t)^{2k}, \qquad k \in \mathbb{Z}_+$$

so it is natural to choose

$$\mu_0 := 3^{\frac{1}{2}} A (T - t)^{2k}$$

as the leading order of the scaling parameter $\mu(t)$.

We next choose

$$u_{in}(x,t) = \mu^{-\frac{1}{2}} w(\frac{x}{\mu}) + 2\mu'_0 \mu^{\frac{1}{2}} J(\frac{x}{\mu})$$
(2.10)

and take the first approximate solution as follows

$$U_1(x,t) := \eta \left(\frac{|x|}{r\sqrt{T-t}}\right) u_{in} + \left(1 - \eta \left(\frac{|x|}{r_1(T-t)^{\zeta_1}}\right)\right) \eta \left(\frac{|x|}{r_2(T-t)^{\zeta_2}}\right) u_{out}, \tag{2.11}$$

where η is the smooth cut-off function satisfying $\eta(t) = 1$ for $t \in [0, 1]$ and $\eta(t) = 0$ for $t \in [2, \infty)$, and $r, r_1, r_2(>0)$ are sufficiently small constants to be determined later. For simplicity, we write

$$\eta_1(x,t) := \eta(\frac{|x|}{r\sqrt{T-t}}),
\eta_{o1}(x,t) = \eta(\frac{|x|}{r_1(T-t)^{\zeta_1}}),
\eta_{o2}(x,t) = \eta(\frac{|x|}{r_2(T-t)^{\zeta_2}}).$$

2.3. Error of the first approximation. We define the error function

$$S(u) := -\partial_t u + \Delta_x u + u^5,$$

and compute

$$\begin{split} S(U_1) &= -\partial_t U_1 + \Delta_x U_1 + U_1^5 \\ &= -\partial_t u_{in} \eta_1 - u_{in} \partial_t \eta_1 + \Delta_x u_{in} \eta_1 + 2 \nabla_x u_{in} \cdot \nabla_x \eta_1 + u_{in} \Delta_x \eta_1 \\ &- \partial_t u_{out} (1 - \eta_{o1}) \eta_{o2} - u_{out} \partial_t ((1 - \eta_{o1}) \eta_{o2}) \\ &+ (1 - \eta_{o1}) \eta_{o2} \Delta_x u_{out} + 2 \nabla_x ((1 - \eta_{o1}) \eta_{o2}) \cdot \nabla_x u_{out} + u_{out} \Delta_x ((1 - \eta_{o1}) \eta_{o2}) \\ &+ \left[u_{in} \eta_1 + (1 - \eta_{o1}) \eta_{o2} u_{out} \right]^5 \\ &= \eta_1 (-\partial_t u_{in} + \Delta_x u_{in} + u_{in}^5) + (1 - \eta_{o1}) \eta_{o2} (-\partial_t u_{out} + \Delta_x u_{out} + u_{out}^5) \\ &- \partial_t \eta_1 u_{in} + \Delta_x \eta_1 u_{in} + 2 \nabla_x \eta_1 \nabla_x u_{in} \\ &- \partial_t [(1 - \eta_{o1}) \eta_{o2}] u_{out} + \Delta_x [(1 - \eta_{o1}) \eta_{o2}] u_{out} + 2 \nabla_x [(1 - \eta_{o1}) \eta_{o2}] \nabla_x u_{out} \\ &+ \left[\eta_1 u_{in} + (1 - \eta_{o1}) \eta_{o2} u_{out} \right]^5 - \eta_1 u_{in}^5 - (1 - \eta_{o1}) \eta_{o2} u_{out}^5. \end{split}$$

Let

$$S_{in} := -\partial_t u_{in} + \Delta_x u_{in} + u_{in}^5,$$

$$S_{out} := -\partial_t u_{out} + \Delta_x u_{out} + u_{out}^5,$$

where we compute

$$\begin{split} \partial_t u_{in} &= -\,\mu^{-\frac{3}{2}} \mu' \big(\frac{w}{2} + \nabla_y w \cdot y\big) + 2 \mu_0'' \mu^{\frac{1}{2}} J(\frac{x}{\mu}) \\ &+ \mu_0' \mu^{-\frac{1}{2}} \mu' J(\frac{x}{\mu}) - 2 \mu_0' \mu^{-\frac{1}{2}} \mu' \nabla_y J(\frac{x}{\mu}) \cdot \frac{x}{\mu}, \\ \partial_\mu \big(\mu^{\frac{1}{2}} J(\frac{x}{\mu})\big) &= \mu^{-\frac{1}{2}} \big[\frac{1}{2} J(\frac{x}{\mu}) - \nabla_y J(\frac{x}{\mu}) \cdot \frac{x}{\mu}\big]. \end{split}$$

Therefore, we have

$$S_{in} = \mu^{-\frac{3}{2}} \mu' \left(\frac{w}{2} + \nabla_{y} w \cdot \frac{x}{\mu}\right) - 2\mu''_{0} \mu^{\frac{1}{2}} J(\frac{x}{\mu})$$

$$- \mu'_{0} \mu^{-\frac{1}{2}} \mu' J(\frac{x}{\mu}) + 2\mu'_{0} \mu^{-\frac{1}{2}} \mu' \nabla_{y} J(\frac{x}{\mu}) \cdot \frac{x}{\mu}$$

$$+ \mu^{-\frac{5}{2}} \Delta_{y} w + 2\mu'_{0} \mu^{-\frac{3}{2}} \Delta_{y} J(\frac{x}{\mu})$$

$$+ \left[\mu^{-\frac{1}{2}} w(\frac{x}{\mu}) + 2\mu'_{0} \mu^{\frac{1}{2}} J(\frac{x}{\mu})\right]^{5}$$

$$= \mu^{-\frac{3}{2}} (\mu - \mu_{0})' \left(\frac{w}{2} + \nabla_{y} w \cdot \frac{x}{\mu}\right) - 2\mu''_{0} \mu^{\frac{1}{2}} J(\frac{x}{\mu})$$

$$- \mu'_{0} \mu^{-\frac{1}{2}} \mu' J(\frac{x}{\mu}) + 2\mu'_{0} \mu^{-\frac{1}{2}} \mu' \nabla_{y} J(\frac{x}{\mu}) \cdot \frac{x}{\mu}$$

$$+ \left[\mu^{-\frac{1}{2}} w(\frac{x}{\mu}) + 2\mu'_{0} \mu^{\frac{1}{2}} J(\frac{x}{\mu})\right]^{5} - \left(\mu^{-\frac{1}{2}} w(\frac{x}{\mu})\right)^{5} - 5\left(\mu^{-\frac{1}{2}} w(\frac{x}{\mu})\right)^{4} 2\mu'_{0} \mu^{\frac{1}{2}} J(\frac{x}{\mu}),$$

where the first term can be written as

$$\mu^{-\frac{3}{2}}(\mu - \mu_0)'(\frac{w}{2} + \nabla_y w \cdot \frac{x}{\mu}) = \mu^{-\frac{3}{2}}(\mu - \mu_0)'\frac{3^{\frac{1}{4}}}{2}\frac{1 - |y|^2}{(1 + |y|^2)^{\frac{3}{2}}}.$$

We notice that

$$(\mu' - \mu_0')\mu^{-\frac{1}{2}} = 2(\mu^{\frac{1}{2}} - \mu_0^{\frac{1}{2}})' + (\mu^{\frac{1}{2}} - \mu_0^{\frac{1}{2}})\mu_0^{-1}\mu_0' - (\mu^{\frac{1}{2}} - \mu_0^{\frac{1}{2}})^2\mu^{-\frac{1}{2}}\mu_0^{-1}\mu_0',$$

and

$$\frac{3^{\frac{1}{4}}}{2}\frac{1-|y|^2}{(1+|y|^2)^{\frac{3}{2}}} = \frac{3^{\frac{1}{4}}}{2}\mu\frac{\mu^2-|x|^2}{(\mu^2+|x|^2)^{\frac{3}{2}}} = 3^{\frac{1}{4}}\frac{\mu^3}{(\mu^2+|x|^2)^{\frac{3}{2}}} - \frac{3^{\frac{1}{4}}}{2}\frac{\mu}{(\mu^2+|x|^2)^{\frac{1}{2}}}.$$

So the leading term is

$$\eta_{1} \left[2(\mu^{\frac{1}{2}} - \mu_{0}^{\frac{1}{2}})' + (\mu^{\frac{1}{2}} - \mu_{0}^{\frac{1}{2}})\mu_{0}^{-1}\mu_{0}' \right] \mu^{-1} \left(-\frac{3^{\frac{1}{4}}}{2} \right) \frac{\mu}{(\mu^{2} + |x|^{2})^{\frac{1}{2}}} \\
= \eta_{1} \left[2(\mu_{0}^{\frac{1}{2}}\Lambda(t))' + \mu_{0}^{\frac{1}{2}}\Lambda(t)\mu_{0}^{-1}\mu_{0}' \right] \left(-\frac{3^{\frac{1}{4}}}{2} \right) \frac{1}{(\mu^{2} + |x|^{2})^{\frac{1}{2}}} \\
= \eta_{1} 2\mu_{0}^{-\frac{1}{2}} \left[\mu_{0}^{\frac{1}{2}}(\mu_{0}^{\frac{1}{2}}\Lambda(t))' + (\mu_{0}^{\frac{1}{2}}\Lambda(t))(\mu_{0}^{\frac{1}{2}})' \right] \left(-\frac{3^{\frac{1}{4}}}{2} \right) \frac{1}{(\mu^{2} + |x|^{2})^{\frac{1}{2}}} \\
= \eta_{1} \frac{\alpha(t)}{(\mu^{2} + |x|^{2})^{\frac{1}{2}}}.$$

Here we define

$$\mu(t) := \mu_0(t)(1 + \Lambda(t))^2,$$

$$\alpha(t) := (-3^{\frac{1}{4}})\mu_0^{-\frac{1}{2}}(t)(\mu_0(t)\Lambda(t))'.$$

For the last term in S_{in} , we compare the size of $\mu^{-\frac{1}{2}}w(\frac{x}{\mu})$ and $2\mu'_0\mu^{\frac{1}{2}}J(\frac{x}{\mu})$ in the regime $\frac{|x|}{r\sqrt{T-t}} \leq 2$ thanks to the cut-off η_1 : if $\frac{|x|}{\mu} \ll 1$, we have

$$\mu^{-\frac{1}{2}} \gg \mu_0' \mu^{\frac{1}{2}} \iff \mu^{-1} \gg \mu_0' \iff 3^{-\frac{1}{2}} A^{-1} (T-t)^{-2k} \gg 3^{\frac{1}{2}} A 2k (T-t)^{2k-1}.$$

Thus we get $\mu^{-\frac{1}{2}}w(\frac{x}{\mu})\gg 2\mu_0'\,\mu^{\frac{1}{2}}J(\frac{x}{\mu})$. If $\frac{|x|}{\mu}\gg 1$, we only need to check

$$\mu^{-\frac{1}{2}} \frac{\mu}{|x|} \gg \mu^{\frac{1}{2}} \mu'_0 \frac{|x|}{\mu} \iff \mu(\mu'_0)^{-1} \gg |x|^2 \iff (T-t) \gg |x|^2$$

since $\frac{|x|}{r\sqrt{T-t}} \le 2$ and $r \ll 1$. So $\mu^{-\frac{1}{2}}w(\frac{x}{\mu}) \gg 2\mu_0' \mu^{\frac{1}{2}}J(\frac{x}{\mu})$ is also satisfied. Therefore, the last term in S_{in} can be expanded as

$$\begin{split} & \left[\mu^{-\frac{1}{2}}w(\frac{x}{\mu}) + 2\mu_0'\,\mu^{\frac{1}{2}}J(\frac{x}{\mu})\right]^5 - \left(\mu^{-\frac{1}{2}}w(\frac{x}{\mu})\right)^5 - 5\left(\mu^{-\frac{1}{2}}w(\frac{x}{\mu})\right)^4 2\mu_0'\,\mu^{\frac{1}{2}}J(\frac{x}{\mu}) \\ &= 5\left[\mu^{-\frac{1}{2}}w(\frac{x}{\mu}) + \theta_1 2\mu_0'\,\mu^{\frac{1}{2}}J(\frac{x}{\mu})\right]^4 2\mu_0'\,\mu^{\frac{1}{2}}J(\frac{x}{\mu}) - 5\left(\mu^{-\frac{1}{2}}w(\frac{x}{\mu})\right)^4 2\mu_0'\,\mu^{\frac{1}{2}}J(\frac{x}{\mu}) \\ &= 20\left[\mu^{-\frac{1}{2}}w(\frac{x}{\mu}) + \theta_2\theta_1 2\mu_0'\,\mu^{\frac{1}{2}}J(\frac{x}{\mu})\right]^3\theta_1\left[2\mu_0'\,\mu^{\frac{1}{2}}J(\frac{x}{\mu})\right]^2 \\ &\lesssim \left[\mu^{-\frac{1}{2}}w(\frac{x}{\mu})\right]^3\,\left[2\mu_0'\,\mu^{\frac{1}{2}}J(\frac{x}{\mu})\right]^2, \end{split}$$

where $\theta_1, \theta_2 \in [0, 1]$ and

$$|J(y)| \le h(y) := \begin{cases} y^2 & \text{if } y \to 0; \\ y & \text{if } y \to \infty. \end{cases}$$
 (2.12)

We define that $\chi(x) = 1$ if $|x| \le 1$ and $\chi(x) = 0$ otherwise. Therefore, we have the following estimate

$$\eta_{1}S_{in} - \chi(\frac{x}{c_{0}(T-t)^{\frac{1}{2}}})\frac{\alpha(t)}{(\mu^{2}+|x|^{2})^{\frac{1}{2}}} \\
= \mu^{-\frac{3}{2}}(\mu-\mu_{0})'(\frac{w}{2}+\nabla_{y}w\cdot\frac{x}{\mu})\eta_{1} - \chi(\frac{x}{c_{0}(T-t)^{\frac{1}{2}}})\frac{\alpha(t)}{(\mu^{2}+|x|^{2})^{\frac{1}{2}}} \\
- 2\mu_{0}''\mu^{\frac{1}{2}}J(\frac{x}{\mu})\eta_{1} - \mu_{0}'\mu^{-\frac{1}{2}}\mu'J(\frac{x}{\mu})\eta_{1} + 2\mu_{0}'\mu^{-\frac{1}{2}}\mu'\nabla_{y}J(\frac{x}{\mu})\cdot\frac{x}{\mu}\eta_{1} \\
+ \left[\left[\mu^{-\frac{1}{2}}w(\frac{x}{\mu}) + 2\mu_{0}'\mu^{\frac{1}{2}}J(\frac{x}{\mu}) \right]^{5} - \left(\mu^{-\frac{1}{2}}w(\frac{x}{\mu}) \right)^{5} - 5\left(\mu^{-\frac{1}{2}}w(\frac{x}{\mu}) \right)^{4}2\mu_{0}'\mu^{\frac{1}{2}}J(\frac{x}{\mu}) \right] \eta_{1} \\
= -2\alpha(t)\frac{\mu^{2}}{(\mu^{2}+|x|^{2})^{\frac{3}{2}}}\eta_{1} + \frac{\alpha(t)}{(\mu^{2}+|x|^{2})^{\frac{1}{2}}}(\eta_{1} - \chi(\frac{x}{c_{0}(T-t)^{\frac{1}{2}}})) \\
- \Lambda^{2}\mu^{-\frac{1}{2}}\mu_{0}'(3^{\frac{1}{4}}\frac{\mu^{2}}{(\mu^{2}+|x|^{2})^{\frac{3}{2}}} - \frac{3^{\frac{1}{4}}}{2}\frac{1}{(\mu^{2}+|x|^{2})^{\frac{1}{2}}})\eta_{1} \\
- 2\mu_{0}''\mu^{\frac{1}{2}}J(\frac{x}{\mu})\eta_{1} - \mu_{0}'\mu^{-\frac{1}{2}}\mu'J(\frac{x}{\mu})\eta_{1} + 2\mu_{0}'\mu^{-\frac{1}{2}}\mu'\nabla_{y}J(\frac{x}{\mu})\cdot\frac{x}{\mu}\eta_{1} \\
+ \left[\left[\mu^{-\frac{1}{2}}w(\frac{x}{\mu}) + 2\mu_{0}'\mu^{\frac{1}{2}}J(\frac{x}{\mu}) \right]^{5} - \left(\mu^{-\frac{1}{2}}w(\frac{x}{\mu}) \right)^{5} - 5\left(\mu^{-\frac{1}{2}}w(\frac{x}{\mu}) \right)^{4}2\mu_{0}'\mu^{\frac{1}{2}}J(\frac{x}{\mu}) \right] \eta_{1} \\
\lesssim |\alpha(t)|\frac{\mu^{2}}{(\mu^{2}+|x|^{2})^{\frac{3}{2}}}\eta_{1} + \frac{|\alpha(t)|}{(\mu^{2}+|x|^{2})^{\frac{1}{2}}}\chi(r \leq \frac{|x|}{\sqrt{T-t}} \leq c_{0}) \\
+ \Lambda^{2}\mu^{-\frac{1}{2}}\mu_{0}\left| 3^{\frac{1}{4}}\frac{\mu^{2}}{(\mu^{2}+|x|^{2})^{\frac{3}{2}}} - \frac{3^{\frac{1}{4}}}{2}\frac{1}{(\mu^{2}+|x|^{2})^{\frac{1}{2}}} \right| \eta_{1} \\
+ 2\mu_{0}''\mu^{\frac{1}{2}}h(\frac{x}{\mu})\eta_{1} + \mu_{0}'\mu^{-\frac{1}{2}}\mu'h(\frac{x}{\mu})\eta_{1} + 2\mu_{0}'\mu^{-\frac{1}{2}}\mu'h(\frac{x}{\mu})\eta_{1} \\
+ \frac{\mu^{\frac{5}{2}}}{(\mu^{2}+|x|^{2})^{\frac{3}{2}}}(\mu'_{0})^{2}h^{2}(\frac{x}{\mu})\eta_{1}. \end{cases}$$

We decompose $u = U_1 + \Phi_1 + \Phi_2$ and compute

$$S(U_{1} + \Phi_{1} + \Phi_{2})$$

$$= -\partial_{t}U_{1} - \partial_{t}\Phi_{1} - \partial_{t}\Phi_{2} + \Delta_{x}U_{1} + \Delta_{x}\Phi_{1} + \Delta_{x}\Phi_{2} + U_{1}^{5} + (U_{1} + \Phi_{1} + \Phi_{2})^{5} - U_{1}^{5}$$

$$= S(U_{1}) - \partial_{t}\Phi_{1} - \partial_{t}\Phi_{2} + \Delta_{x}\Phi_{1} + \Delta_{x}\Phi_{2} + (U_{1} + \Phi_{1} + \Phi_{2})^{5} - U_{1}^{5}$$

$$= S(U_{1}) - \chi(\frac{x}{c_{0}(T - t)^{\frac{1}{2}}})\frac{\alpha(t)}{(\mu^{2} + |x|^{2})^{\frac{1}{2}}} + \chi(\frac{x}{c_{0}(T - t)^{\frac{1}{2}}})\frac{\alpha(t)}{(\mu^{2} + |x|^{2})^{\frac{1}{2}}} - \partial_{t}\Phi_{1}$$

$$- \partial_{t}\Phi_{2} + \Delta_{x}\Phi_{1} + \Delta_{x}\Phi_{2} + (U_{1} + \Phi_{1} + \Phi_{2})^{5} - U_{1}^{5}$$

where $\chi(x) = 1$ if $|x| \le 1$ and $\chi(x) = 0$ otherwise, c_0 is some constant, Φ_1 and Φ_2 are perturbations to be determined later.

2.4. Nonlocal correction: second approximation. Observe that in the error there is a slow decaying term $\frac{\alpha(t)}{(\mu^2(t)+|x|^2)^{1/2}}$. Following the idea in [7], we now introduce a nonlocal correction Φ_1 solving

$$\partial_t \Phi_1 = \Delta_x \Phi_1 + \chi(\frac{x}{c_0(T-t)^{\frac{1}{2}}}) \frac{\alpha(t)}{(\mu^2(t) + |x|^2)^{1/2}}.$$
 (2.14)

Choosing the initial data, we get from Duhamel's formula

$$\Phi_{1}(x,t) = \sum_{j=1}^{k} c_{j} \mathcal{B}^{(j)}(x,t) + \int_{0}^{t} \int_{\mathbb{R}^{3}} \left(\frac{1}{2\sqrt{\pi}}\right)^{3} (t-s)^{-\frac{3}{2}} e^{-\frac{|x-\xi|^{2}}{4(t-s)}} \chi\left(\frac{\xi}{c_{0}(T-s)^{\frac{1}{2}}}\right) \frac{\alpha(s)}{(\mu^{2}(s)+|\xi|^{2})^{1/2}} d\xi ds$$
(2.15)

where c_j are constants and $\mathcal{B}^{(j)}$ satisfy heat equation

$$\partial_t \mathcal{B}^{(j)}(x,t) = \Delta_x \mathcal{B}^{(j)}(x,t)$$

which will be determined later when solving $\alpha(t)$ from the reduced equation in Section 5.

$$S(U_1 + \Phi_1) = S(U_1) - \chi(\frac{x}{c_0(T - t)^{\frac{1}{2}}}) \frac{\alpha(t)}{(\mu^2 + |x|^2)^{\frac{1}{2}}} + (U_1 + \Phi_1)^5 - U_1^5$$
Let $\Phi = \Phi_1 + \Phi_2$. We have
$$S(U_1 + \Phi_1 + \Phi_2)$$

$$= S(U_1) - \chi(\frac{x}{c_0(T - t)^{\frac{1}{2}}}) \frac{\alpha(t)}{(\mu^2 + |x|^2)^{\frac{1}{2}}} - \partial_t \Phi_2 + \Delta_x \Phi_2 + (U_1 + \Phi_1 + \Phi_2)^5 - U_1^5$$

$$= S(U_1) - \chi(\frac{x}{c_0(T - t)^{\frac{1}{2}}}) \frac{\alpha(t)}{(\mu^2 + |x|^2)^{\frac{1}{2}}} + (U_1 + \Phi_1)^5 - U_1^5$$

$$- \partial_t \Phi_2 + \Delta_x \Phi_2 + (U_1 + \Phi_1 + \Phi_2)^5 - (U_1 + \Phi_1)^5$$

$$= S(U_1) - \chi \left(\frac{x}{c_0(T-t)^{\frac{1}{2}}}\right) \frac{\alpha(t)}{(\mu^2 + |x|^2)^{\frac{1}{2}}} + (U_1 + \Phi_1)^5 - U_1^5$$
$$- \partial_t \Phi_2 + \Delta_x \Phi_2 + 5(U_1 + \Phi_1)^4 \Phi_2 + (U_1 + \Phi_1 + \Phi_2)^5 - (U_1 + \Phi_1)^5 - 5(U_1 + \Phi_1)^4 \Phi_2.$$

3. Inner-outer gluing system

In this section, we set up the inner-outer gluing scheme for the nonlinear problem. We look for perturbation of the following form

$$\Phi_{2}(x,t) = \psi(x,t) + \phi^{in}(x,t) = \psi(x,t) + \eta(\frac{x}{R\mu_{0}})\hat{\phi}(x,t)$$

$$= \psi(x,t) + \eta(\frac{x}{R\mu_{0}})\mu_{0}^{-\frac{1}{2}}\phi(\frac{x}{\mu_{0}},t)$$

$$= \psi(x,t) + \eta_{R}\mu_{0}^{-\frac{1}{2}}\phi(\frac{x}{\mu_{0}},t)$$

where $\hat{\phi}(x,t) := \mu_0^{-\frac{1}{2}} \phi(\frac{x}{\mu_0},t), \ \eta_R := \eta(\frac{x}{R\mu_0}), \ \text{and} \ R(t) = \mu_0^{-\beta}(t) \ \text{with} \ \beta \in (0,1/2).$ We next compute $\partial_t \Phi_2(x,t) = \partial_t \psi(x,t) + \partial_t \eta_R \mu_0^{-\frac{1}{2}} \phi(\frac{x}{\mu_0},t) + \eta_R \partial_t (\mu_0^{-\frac{1}{2}} \phi(\frac{x}{\mu_0},t))$

where

$$\begin{split} \partial_t(\mu_0^{-\frac{1}{2}}\phi(\frac{x}{\mu_0},t)) &= -\frac{1}{2}\mu_0^{-\frac{3}{2}}\mu_0'\phi(\frac{x}{\mu_0},t) - \mu_0^{-\frac{3}{2}}\nabla_y\phi(\frac{x}{\mu_0},t) \cdot \frac{x}{\mu_0}\mu_0' + \mu_0^{-\frac{1}{2}}\partial_t\phi(\frac{x}{\mu_0},t) \\ &= -\mu_0^{-\frac{3}{2}}\mu_0'\left[\frac{1}{2}\phi(\frac{x}{\mu_0},t) + \nabla_y\phi(\frac{x}{\mu_0},t) \cdot \frac{x}{\mu_0}\right] + \mu_0^{-\frac{1}{2}}\partial_t\phi(\frac{x}{\mu_0},t) \end{split}$$

and

$$\Delta_x \Phi_2(x,t) = \Delta_x \psi(x,t) + \Delta_x \eta_R \mu_0^{-\frac{1}{2}} \phi(\frac{x}{\mu_0},t) + 2\nabla_x \eta_R \mu_0^{-\frac{3}{2}} \nabla_y \phi(\frac{x}{\mu_0},t) + \eta_R \mu_0^{-\frac{5}{2}} \Delta_y \phi(\frac{x}{\mu_0},t).$$

Therefore, we obtain

$$S(U_{1} + \Phi_{1} + \Phi_{2})$$

$$= S(U_{1}) - \chi(\frac{x}{c_{0}(T - t)^{\frac{1}{2}}}) \frac{\alpha(t)}{(\mu^{2} + |x|^{2})^{\frac{1}{2}}} + (U_{1} + \Phi_{1})^{5} - U_{1}^{5}$$

$$- \partial_{t}\psi(x, t) - \partial_{t}\eta_{R}\mu_{0}^{-\frac{1}{2}}\phi(\frac{x}{\mu_{0}}, t)$$

$$+ \eta_{R}\mu_{0}^{-\frac{3}{2}}\mu_{0}' \left[\frac{1}{2}\phi(\frac{x}{\mu_{0}}, t) + \nabla_{y}\phi(\frac{x}{\mu_{0}}, t) \cdot \frac{x}{\mu_{0}}\right] - \eta_{R}\mu_{0}^{-\frac{1}{2}}\partial_{t}\phi(\frac{x}{\mu_{0}}, t)$$

$$+ \Delta_{x}\psi(x, t) + \Delta_{x}\eta_{R}\mu_{0}^{-\frac{1}{2}}\phi(\frac{x}{\mu_{0}}, t) + 2\nabla_{x}\eta_{R} \cdot \mu_{0}^{-\frac{3}{2}}\nabla_{y}\phi(\frac{x}{\mu_{0}}, t) + \eta_{R}\mu_{0}^{-\frac{5}{2}}\Delta_{y}\phi(\frac{x}{\mu_{0}}, t)$$

$$+ 5(U_{1} + \Phi_{1})^{4}(\psi(x, t) + \eta_{R}\mu_{0}^{-\frac{1}{2}}\phi(\frac{x}{\mu_{0}}, t))$$

$$+ (U_{1} + \Phi_{1} + \Phi_{2})^{5} - (U_{1} + \Phi_{1})^{5} - 5(U_{1} + \Phi_{1})^{4}\Phi_{2}.$$

It can be directly checked that $S(U_1 + \Phi_1 + \Phi_2) = 0$ if ψ solves

$$-\partial_{t}\psi(x,t) + \Delta_{x}\psi(x,t) - \partial_{t}\eta_{R}\mu_{0}^{-\frac{1}{2}}\phi(\frac{x}{\mu_{0}},t) + \Delta_{x}\eta_{R}\mu_{0}^{-\frac{1}{2}}\phi(\frac{x}{\mu_{0}},t) + 2\nabla_{x}\eta_{R}\cdot\mu_{0}^{-\frac{3}{2}}\nabla_{y}\phi(\frac{x}{\mu_{0}},t) + 5(U_{1} + \Phi_{1})^{4}(1 - \eta_{R})\psi(x,t) + 5\left[(U_{1} + \Phi_{1})^{4} - \left(\mu^{-\frac{1}{2}}w(\frac{x}{\mu})\right)^{4}\right]\psi(x,t)\eta_{R} + (U_{1} + \Phi_{1} + \Phi_{2})^{5} - (U_{1} + \Phi_{1})^{5} - 5(U_{1} + \Phi_{1})^{4}\Phi_{2} + \left[S(U_{1}) - \chi(\frac{x}{c_{0}(T - t)^{\frac{1}{2}}})\frac{\alpha(t)}{(\mu^{2} + |x|^{2})^{\frac{1}{2}}}\right](1 - \eta_{R}) + \left[(U_{1} + \Phi_{1})^{5} - U_{1}^{5}\right](1 - \eta_{R}) = 0$$

$$(3.1)$$

and ϕ solves

$$\eta_{R}\mu_{0}^{-\frac{3}{2}}\mu_{0}'\left[\frac{1}{2}\phi(\frac{x}{\mu_{0}},t) + \nabla_{y}\phi(\frac{x}{\mu_{0}},t) \cdot \frac{x}{\mu_{0}}\right] - \eta_{R}\mu_{0}^{-\frac{1}{2}}\partial_{t}\phi(\frac{x}{\mu_{0}},t)
+ \eta_{R}\mu_{0}^{-\frac{5}{2}}\Delta_{y}\phi(\frac{x}{\mu_{0}},t) + 5\left(\mu^{-\frac{1}{2}}w(\frac{x}{\mu})\right)^{4}\psi(x,t)\eta_{R} + 5(U_{1} + \Phi_{1})^{4}\mu_{0}^{-\frac{1}{2}}\phi(\frac{x}{\mu_{0}},t)\eta_{R}
+ \eta_{R}\left[S(U_{1}) - \chi(\frac{x}{c_{0}(T-t)^{\frac{1}{2}}})\frac{\alpha(t)}{(\mu^{2}+|x|^{2})^{\frac{1}{2}}}\right] + \left[(U_{1} + \Phi_{1})^{5} - U_{1}^{5}\right]\eta_{R} = 0.$$
(3.2)

Recall that in the inner regime one has

$$U_1 = u_{in} = \mu^{-\frac{1}{2}} w(\frac{x}{\mu}) + 2\mu'_0 \mu^{\frac{1}{2}} J(\frac{x}{\mu}).$$

Writing $y = \frac{x}{\mu_0}$, we rearrange problems (3.1) and (3.2) and get

• The outer problem:

$$\partial_t \psi(x,t) = \Delta_x \psi(x,t) + \mathcal{G}(\phi,\psi,\alpha) \tag{3.3}$$

where

$$\mathcal{G}(\phi,\psi,\alpha) = -\partial_t \eta_R \mu_0^{-\frac{1}{2}} \phi(\frac{x}{\mu_0},t) + \Delta_x \eta_R \mu_0^{-\frac{1}{2}} \phi(\frac{x}{\mu_0},t) + 2\nabla_x \eta_R \cdot \mu_0^{-\frac{3}{2}} \nabla_y \phi(\frac{x}{\mu_0},t)
+ \left(5(U_1 + \Phi_1)^4 (1 - \eta_R) + 5 \left[(U_1 + \Phi_1)^4 - \left(\mu^{-\frac{1}{2}} w(\frac{x}{\mu}) \right)^4 \right] \eta_R \right) \psi
+ (U_1 + \Phi_1 + \Phi_2)^5 - (U_1 + \Phi_1)^5 - 5(U_1 + \Phi_1)^4 \Phi_2
+ \left[S(U_1) - \chi \left(\frac{x}{c_0 (T - t)^{\frac{1}{2}}} \right) \frac{\alpha(t)}{(\mu^2 + |x|^2)^{\frac{1}{2}}} \right] (1 - \eta_R) + \left[(U_1 + \Phi_1)^5 - U_1^5 \right] (1 - \eta_R)$$
(3.4)

• The inner problem:

$$\mu_0^2 \partial_t \phi(y, t) = \Delta_y \phi(y, t) + 5w^4(y)\phi(y, t) + \mathcal{H}(\phi, \psi, \alpha) \quad \text{in } B_{2R} \times (0, T),$$
(3.5)

where

$$\mathcal{H}(\phi,\psi,\alpha) = 5\left[(u_{in}(\mu_{0}y,t) + \Phi_{1}(\mu_{0}y,t))^{4} - \mu_{0}^{-2}w^{4}(y)\right] \mu_{0}^{2}\phi(y,t)$$

$$+ 5\mu_{0}^{\frac{1}{2}}(1+\Lambda)^{-4}w^{4}(\frac{y}{(1+\Lambda)^{2}})\psi(\mu_{0}y,t) + \mu_{0}\mu_{0}'\left[\frac{1}{2}\phi(y,t) + \nabla_{y}\phi(y,t) \cdot y\right]$$

$$+ \mu_{0}^{\frac{5}{2}}\left[S(U_{1}) - \chi(\frac{\mu_{0}y}{c_{0}(T-t)^{\frac{1}{2}}})\frac{\alpha(t)}{(\mu^{2} + |\mu_{0}y|^{2})^{\frac{1}{2}}}\right]$$

$$+ \mu_{0}^{\frac{5}{2}}\left[(u_{in}(\mu_{0}y,t) + \Phi_{1}(\mu_{0}y,t))^{5} - u_{in}^{5}(\mu_{0}y,t)\right]$$

$$(3.6)$$

We choose $\zeta_1 = \zeta_2 = \frac{1}{2}$ so that $\left(1 - \eta\left(\frac{|x|}{r_1(T-t)^{\zeta_1}}\right)\right)\eta\left(\frac{|x|}{r_2(T-t)^{\zeta_2}}\right)u_{out} \equiv 0$ in the inner region $\{x: |x| \leq 2R\mu_0\}$

$$r_1(T-t)^{\zeta_1} \gg 2R\mu_0$$

for $T \ll 1$ since $R(t) = \mu_0^{-\beta}$ with $\beta \in (0, 1/2)$.

Performing the change of variables

$$\frac{\mathrm{d}\tau}{\mathrm{d}t} = \mu_0^{-2}, \quad y = \frac{x}{\mu_0}$$

so that

$$\tau = \frac{1}{3A^2(4k-1)}(T-t)^{1-4k}, \quad \mu_0 = (3^{\frac{1}{2}}A)^{\frac{-1}{4k-1}}(4k-1)^{-\frac{2k}{4k-1}}\tau^{-\frac{2k}{4k-1}}$$

and writing $\phi(y,\tau) = \phi(y,t(\tau))$, we obtain

$$\begin{split} &-\partial_{\tau}\phi(y,\tau)+\Delta_{y}\phi(y,\tau)+5w^{4}(y)\phi(y,\tau)\\ &+5\left[\left(u_{in}(\mu_{0}y,t(\tau))+\Phi_{1}(\mu_{0}y,t(\tau))\right)^{4}-\mu_{0}^{-2}w^{4}(y)\right]\mu_{0}^{2}\phi(y,\tau)\\ &+5\mu_{0}^{\frac{1}{2}}(1+\Lambda)^{-4}w^{4}(\frac{y}{(1+\Lambda)^{2}})\psi(\mu_{0}y,t(\tau))+\mu_{0}\mu_{0}'\left[\frac{1}{2}\phi(y,\tau)+\nabla_{y}\phi(y,\tau)\cdot y\right]\\ &+\mu_{0}^{\frac{5}{2}}\left[S(U_{1})-\chi(\frac{\mu_{0}y}{c_{0}[3A^{2}(4k-1)\tau]^{\frac{1}{2(1-4k)}}})\frac{\alpha(t(\tau))}{(\mu^{2}+|\mu_{0}y|^{2})^{\frac{1}{2}}}\right]\\ &+\mu_{0}^{\frac{5}{2}}\left[\left(u_{in}(\mu_{0}y,t(\tau))+\Phi_{1}(\mu_{0}y,t(\tau))\right)^{5}-u_{in}^{5}(\mu_{0}y,t(\tau))\right]=0 \quad \text{in} \quad B_{2R}\times(\tau_{0},+\infty). \end{split}$$

In the sequel, we shall solve the inner-outer gluing system (3.3) and (3.5) by developing suitable linear theories and the fixed point argument.

4. Linear theories

4.1. Linear theory for the inner problem. By using the Fourier decomposition and delicate analysis, the authors in [7] developed the linear theory for the inner problem (3.5) in dimension 3. Denote Z_{-} by the positive radial bounded eigenfunction associated to the only negative eigenvalue λ_{-} to

$$\Delta \phi + 5w^4 \phi + \lambda_- \phi = 0, \quad \phi \in L^{\infty}(\mathbb{R}^3).$$

It is further known that λ_{-} is simple and Z_{-} satisfies

$$Z_{-}(y) \sim |y|^{-1} e^{-\sqrt{|\lambda_0|}|y|}$$
 as $|y| \to \infty$.

Consider an orthonormal basis $\{\Theta_i\}_{i=0}^{\infty}$ made up of spherical harmonics in $L^2(\mathbb{S}^2)$, i.e.

$$\Delta_{\mathbb{S}^2}\Theta_i + \lambda_i\Theta_i = 0$$
 in \mathbb{S}^2

with $0 = \lambda_0 < \lambda_1 = \lambda_2 = \lambda_3 = 2 < \lambda_4 \le \cdots$. More precisely, $\Theta_0(y) = a_0$, $\Theta_i(y) = a_1 y_i$, $i = 1, \dots, 3$ for two constants a_0 , a_1 . For $h \in L^2(B_{2R})$, we decompose

$$h(y,t) = \sum_{j=0}^{\infty} h_j(r,t)\Theta_j(y/r), \ r = |y|, \ h_j(r,t) = \int_{\mathbb{S}^2} h(r\theta,t)\Theta_j(\theta)d\theta$$

and write $h = h^0 + h^1 + h^{\perp}$ with

$$h^0 = h_0(r, t), \ h^1 = \sum_{j=1}^3 h_j(r, t)\Theta_j, \ h^{\perp} = \sum_{j=4}^\infty h_j(r, t)\Theta_j.$$

Also, we decompose $\phi = \phi^0 + \phi^1 + \phi^{\perp}$ in a similar form. Define

$$||h||_{\nu,2+\sigma} := \sup_{(y,t)\in B_{2R}\times(0,T)} \mu_0^{-\nu}(t)(1+|y|^{2+\sigma})|h(y,t)|. \tag{4.1}$$

Proposition 4.1 ([7]). Let ν , σ be given positive numbers with $0 < \sigma < 2$. Then, for all sufficiently large R > 0 and any h = h(y,t) with $||h||_{\nu,2+\sigma} < +\infty$ that satisfies for all $j = 0,1,\ldots 3$

$$\int_{B_{2R}} h(y,t)Z_j(y) \, \mathrm{d}y = 0 \quad \text{for all} \quad t \in (0,T)$$

there exist $\phi = \phi[h]$ and $e_0 = e_0[h]$ which solve

$$\mu_0^2 \phi_t = \Delta \phi + 5w^4 \phi + h(y,t) \text{ in } B_{2R} \times (0,T), \quad \phi(y,0) = e_0 Z_-(y) \text{ in } B_{2R}.$$

Moreover, they define linear operators of h that satisfies the estimates

$$|\phi(y,t)| \lesssim \mu_0^{\nu}(t) \left[\frac{R^{4-\sigma}}{1+|y|^3} \|h^0\|_{\nu,2+\sigma} + \frac{R^{4-\sigma}}{1+|y|^4} \|h^1\|_{\nu,2+\sigma} + \frac{\|h\|_{\nu,2+\sigma}}{1+|y|^{\sigma}} \right]$$

$$|\nabla_y \phi(y,t)| \lesssim \mu_0^{\nu}(t) \left[\frac{R^{4-\sigma}}{1+|y|^4} \|h^0\|_{\nu,2+\sigma} + \frac{R^{4-\sigma}}{1+|y|^5} \|h^1\|_{\nu,2+\sigma} + \frac{\|h\|_{\nu,2+\sigma}}{1+|y|^{\sigma+1}} \right]$$

and

$$|e_0[h]| \lesssim ||h||_{\nu,2+\sigma}.$$

Remark 4.1.

- (1) Since the blow-up profile constructed in the paper is radially symmetric, we only consider the case of mode 0 in the Fourier decomposition. The construction of nonradial solutions can be carried out in a similar manner as in [7, Section 10] with nonradial perturbation.
- (2) Using the method of supersolution as in [6, Lemma 7.3], one can improve the linear theory at mode 0 in the interior region:

$$|\phi^{0}(y,t)| + (1+|y|)|\nabla\phi^{0}(y,t)| \lesssim \frac{\mu_{0}^{\nu}(t)R^{\frac{4-\sigma}{3}}}{1+|y|} ||h^{0}||_{\nu,2+\sigma} \min\left\{1, (1+|y|)^{-2}R^{\frac{2(4-\sigma)}{3}}\right\}.$$

If we define

$$\|\phi\|_{0,\nu,\sigma} := \sup_{(y,t)\in B_{2R}\times(0,T)} \frac{1+|y|}{\mu_0^{\nu}(t)R^{\frac{4-\sigma}{3}}(t)} \left[|\phi(y,t)| + (1+|y|)|\nabla\phi(y,t)| \right],\tag{4.2}$$

then under the assumptions of Proposition 4.1, we have

$$\|\phi^0\|_{0,\nu,\sigma} \lesssim \|h^0\|_{\nu,2+\sigma}.$$

(3) Under the assumptions in Proposition 4.1, if the right hand side h(y,t) further satisfies

$$h(y,t) \in C^{2k-2+2\epsilon,k-1+\epsilon}_{y,t}(B_{4R} \times (0,T))$$

for some $0 < \epsilon < 1$, then

$$\|\phi\|_{C^{2k,k}_{y,t}(B_{2R}\times(0,T))} \lesssim T^{\epsilon}.$$

This is a consequence of scaling argument and the parabolic Schauder estimate.

4.2. Linear theory for the outer problem. In this section, we develop the linear theory for the outer problem. The model problem is

$$\begin{cases} \psi_t = \Delta \psi + f, & \text{in } \mathbb{R}^3 \times (0, T) \\ \psi(\cdot, 0) = \psi_0, & \text{in } \mathbb{R}^3 \end{cases}$$
(4.3)

where the non-homogeneous term f in (4.3) is assumed to be bounded with respect to the weights appearing in the outer problem (3.3). Define the weights

$$\begin{cases}
\varrho_{1} := \mu_{0}^{\nu - \frac{5}{2}}(t)R^{-2-a}(t)\chi_{\{|x| \leq 2\mu_{0}R\}} \\
\varrho_{2} := \frac{\mu_{0}^{\nu_{2}}}{|x|^{a_{2}}}\chi_{\{|x| \geq \mu_{0}R\}} \\
\varrho_{3} := 1
\end{cases}$$
(4.4)

where $\nu > 0$, 0 < a < 1, $1 \le a_2 \le 2$, $\nu_2 > 0$, and we choose $R(t) = \mu_0^{-\beta}(t)$ for $\beta \in (0, 1/2)$ throughout the paper. We define the norms

$$||f||_{**} := \sup_{(x,t)\in\mathbb{R}^3\times(0,T)} \left(\sum_{i=1}^3 \varrho_i(x,t)\right)^{-1} |f(x,t)| \tag{4.5}$$

$$\|\psi\|_{*} := \mu_{0}^{\frac{1}{2}-\nu}(0)R^{a}(0)\|\psi\|_{L^{\infty}(\mathbb{R}^{3}\times(0,T))} + \mu_{0}^{\frac{3}{2}-\nu}(0)R^{1+a}(0)\|\nabla\psi\|_{L^{\infty}(\mathbb{R}^{3}\times(0,T))}$$

$$+ \sup_{(x,t)\in\mathbb{R}^{3}\times(0,T)} \left[\mu_{0}^{\frac{1}{2}-\nu}(t)R^{a}(t)|\psi(x,t)-\psi(x,T)|\right]$$

$$+ \sup_{(x,t)\in\mathbb{R}^{3}\times(0,T)} \left[\mu_{0}^{\frac{3}{2}-\nu}(t)R^{1+a}(t)||\nabla\psi(x,t)-\nabla\psi(x,T)|\right]$$

$$+ \sup_{\mathbb{R}^{3}\times I_{T}} \frac{\mu_{0}^{2\gamma+\frac{1}{2}-\nu}(t_{2})R^{2\gamma+a}(t_{2})}{(t_{2}-t_{1})^{\gamma}}|\psi(x,t_{2})-\psi(x,t_{1})|,$$

$$(4.6)$$

where $\nu > 0$, $0 < a, \gamma < 1$, and the last supremum is taken over

$$\mathbb{R}^3 \times I_T = \left\{ (x, t_1, t_2) : x \in \mathbb{R}^3, \ 0 \le t_1 \le t_2 \le T, \ t_2 - t_1 \le \frac{1}{10} (T - t_2) \right\}.$$

For problem (4.3), we have the following estimates.

Proposition 4.2. Let ψ be the solution to problem (4.3) with $||f||_{**} < +\infty$. Then it holds that $\|\psi\|_* \lesssim \|f\|_{**}$.

Proposition 4.2 is established by the following three lemmas with different right hand sides.

Lemma 4.1. Let ψ solve problem (4.3) with right hand side

$$|f(x,t)| \lesssim \mu_0^{\nu - \frac{5}{2}}(t)R^{-2-a}(t)\chi_{\{|x| \le 2\mu_0 R\}}$$

with 0 < a < 1 and $\nu > 0$. Then it holds that

$$|\psi(x,t)| \lesssim \mu_0^{\nu - \frac{1}{2}}(0)R^{-a}(0),$$
 (4.7)

$$|\psi(x,t) - \psi(x,T)| \lesssim \mu_0^{\nu - \frac{1}{2}}(t)R^{-a}(t),$$
 (4.8)

$$|\nabla \psi(x,t)| \lesssim \mu_0^{\nu - \frac{3}{2}}(0)R^{-1-a}(0),$$
 (4.9)

$$|\nabla \psi(x,t) - \nabla \psi(x,T)| \lesssim \mu_0^{\nu - \frac{3}{2}}(t)R^{-1-a}(t),$$
 (4.10)

and

$$|\psi(x,t_2) - \psi(x,t_1)| \lesssim \mu_0^{\nu + \gamma_1 - \frac{5}{2}} (t_2) R^{\gamma_1 - 2 - a} (t_2) (t_2 - t_1)^{1 - \gamma_1/2},$$
 (4.11)

where $0 \le t_1 \le t_2 \le T$ with $t_2 - t_1 \le \frac{1}{10}(T - t_2)$ and $\gamma_1 \in (0, 1)$.

Lemma 4.2. Let ψ solve problem (4.3) with right hand side

$$|f(x,t)| \lesssim \frac{\mu_0^{\nu_2}}{|x|^{a_2}} \chi_{\{|x| \ge \mu_0 R\}},$$

where $\nu_2 > 0$ and $1 \le a_2 \le 2$. Then it holds that

$$|\psi(x,t)| \lesssim \mu_0^{\nu_2 + \frac{2-a_2}{4k}}(0),$$
 (4.12)

$$|\psi(x,t) - \psi(x,T)| \lesssim \mu_0^{\nu_2 + \frac{2-a_2}{4k}}(t), \tag{4.13}$$

$$|\nabla \psi(x,t)| \lesssim \mu_0^{\nu_2 + \frac{1-a_2}{4k}}(0), \tag{4.14}$$

$$|\nabla \psi(x,t)| \lesssim \mu_0^{\nu_2 + \frac{1-a_2}{4k}}(0),$$
 (4.14)

$$|\nabla \psi(x,t) - \nabla \psi(x,T)| \lesssim \mu_0^{\nu_2 + \frac{1-a_2}{4k}}(t),$$
 (4.15)

and

$$|\psi(x,t_2) - \psi(x,t_1)| \lesssim \frac{\mu_0^{\nu_2}(t)}{(\mu_0(t)R(t))^{2\gamma}} (t_2 - t_1)^{\gamma},$$
 (4.16)

where $0 \le t_1 \le t_2 \le T$ with $t_2 - t_1 \le \frac{1}{10}(T - t_2)$ and $\gamma \in (0, 1)$.

Lemma 4.3. Let ψ solve problem (4.3) with right hand side

$$|f(x,t)| \lesssim 1.$$

Then it holds that

$$|\psi(x,t)| \lesssim t,$$

$$|\psi(x,t) - \psi(x,T)| \lesssim (T-t),$$

$$|\nabla \psi(x,t)| \lesssim T^{1/2},$$

$$|\nabla \psi(x,t) - \nabla \psi(x,T)| \lesssim (T-t)^{1/2}.$$

and

$$|\psi(x,t_2) - \psi(x,t_1)| \lesssim (t_2 - t_1),$$

where $0 \le t_1 \le t_2 \le T$ with $t_2 - t_1 \le \frac{1}{10}(T - t_2)$.

Proposition 4.2 is a direct consequence of Lemma 4.1, Lemma 4.2 and Lemma 4.3, and the proofs of Lemma 4.1, Lemma 4.2 and Lemma 4.3 are achieved by using Duhamel's formula similarly as in [6]. Here we only give proof of Lemma 4.1 and Lemma 4.2 in the Appendix.

Remark 4.2.

(1) Let us point out the reason why we use the $\|\cdot\|_*$ -norm of ψ ((4.6)) only involving ν but not ν_2 appearing in Lemma 4.2. Lemma 4.2 is needed to deal with the right hand side of outer problem with cut-off $1 - \eta_R$ in front. For convenience, when we carry out the inner-outer gluing procedure to bound right hand sides in the chosen topology, we will adjust ν_2 such that the control of ψ in Lemma 4.2 is better than that of Lemma 4.1. This will result in a constraint for the parameters

$$\nu_2 + \frac{2 - a_2}{4k} > \nu - \frac{1}{2} + a\beta.$$

In fact, the above constraint will be satisfied by the choices of parameters in Section 6.4.

(2) Under the assumptions of Proposition 4.2, if we further assume that

$$f(x,t) \in C_{x,t}^{2k-2+2\epsilon,k-1+\epsilon}(\mathbb{R}^3 \times (0,T)),$$

then

$$\|\psi\|_{C^{2k,k}_{x,t}(B_{\mu_0(0)R(0)}\times(0,T))} \lesssim T^{\epsilon}$$

for some $0 < \epsilon < 1$.

5. The reduced equation for $\alpha(t)$

From the linear theory for the inner problem (3.5) in Section 4.1, orthogonality condition is required to guarantee the existence of solutions with sufficient space-time decay. In this section, we will adjust the scaling parameter $\mu(t)$ by such orthogonality condition.

Recall that the slow decaying kernel for the linearized operator

$$Z_0(y) = -\left[y \cdot \nabla w + \frac{w}{2}\right] = \frac{3^{\frac{1}{4}}}{2} \frac{|y|^2 - 1}{(1 + |y|^2)^{\frac{3}{2}}}.$$

and

$$\mu(t) = \mu_0(t)(1 + \Lambda(t))^2,$$

$$\alpha(t) = (-3^{\frac{1}{4}})\mu_0^{-\frac{1}{2}}(t)(\mu_0(t)\Lambda(t))'.$$

Since $\mu_0(t) \sim (T-t)^{2k}$, our aim is to look for

$$\alpha(t) \sim (T-t)^{k-1} \Lambda(t), \quad k \in \mathbb{Z}_+.$$

where $\Lambda(t) \to 0$ as $t \to T$.

From the linear theory in Section 4.1, the inner solution can be found in suitable topology if the following orthogonality condition is satisfied

$$\int_{B_{2R}} \mathcal{H}(\phi, \psi, \alpha) Z_0(y) dy = 0 \quad \text{for all} \quad t \in (0, T), \tag{5.1}$$

where $\mathcal{H}(\phi, \psi, \alpha)$ is defined in (3.6). Define

$$||h||_{\delta} := \sup_{t \in (0,T)} |(T-t)^{-\delta}h(t)|.$$
 (5.2)

It turns out that the reduced problem orthogonality is a problem involving the following nonlocal operator

$$\int_0^t \frac{\alpha(s)}{(t-s)^{1/2}} ds,$$

which is the nonlocal feature inherited from the slow decaying kernel $Z_0(y)$.

5.1. A linear theory for the reduced equation. Before we consider the reduced problem (5.1), we first develop a key linear theory for the following problem

$$\int_0^t \frac{\alpha(s)}{(t-s)^{1/2}} ds = \sum_{j=1}^k c_j \mathcal{B}^{(j)}(0,t) + h(t), \tag{5.3}$$

where $h(t) \in C_t^k(0,T)$, and for j = 1, ..., k, c_j are constants and $\mathcal{B}^{(j)}$ are smooth functions to be determined. We have the following lemma concerning the solvability of problem (5.3), which enables us to solve $\alpha(t)$ with sufficiently fast decay in problem (5.1).

Lemma 5.1. For problem (5.3), if $h(t) \in C_t^k(0,T)$, then there exist constants c_j and smooth functions $\mathcal{B}^{(j)}$ such that problem (5.3) has a solution satisfying

$$\alpha(t) \sim (T-t)^{k-1} \Lambda(t), \quad k \in \mathbb{Z}_+,$$

where $\Lambda(t) \to 0$ as $t \to T$.

Proof. In order to find $\alpha(t)$ with the above vanishing order, it suffices to show that

$$\alpha(T) = \alpha'(T) = \alpha''(T) = \dots = \alpha^{(k-1)}(T) = 0.$$

We shall choose c_i and $\mathcal{B}^{(j)}$ using solutions for heat equations as building blocks.

We first find solutions $B_j(x,t)$ to the following heat equation

$$\begin{cases} \partial_t B_j = \Delta B_j & \text{in } \mathbb{R}^3 \times (0, T) \\ B_j(\cdot, 0) = B_{0,j} & \text{in } \mathbb{R}^3 \end{cases}$$
 (5.4)

where the decaying initial data $B_{0,j}$ will be chosen. Using Duhamel's formula in problem (5.4), we write

$$B_j(0,t) = \int_{\mathbb{R}^3} e^{-\frac{|\xi|^2}{4t}} \frac{B_{0,j}(\xi)}{(4\pi t)^{3/2}} d\xi.$$

Let us choose the initial condition

$$B_{0,j}(|x|) = e^{-\kappa_j |x|^2}.$$

Then

$$B_{j}(0,t) = \frac{1}{(4\pi)^{3/2}} \int_{\mathbb{R}^{3}} t^{-3/2} e^{-\frac{|\xi|^{2}}{4t}} e^{-\kappa_{j}|\xi|^{2}} d\xi$$

$$= \frac{1}{(4\pi)^{3/2}} \int_{\mathbb{R}^{3}} t^{-3/2} e^{-|x|^{2}} \left(\kappa_{j} t + \frac{1}{4}\right)^{-3/2} dx$$

$$= \frac{1}{8} \left(\kappa_{j} t + \frac{1}{4}\right)^{-3/2}$$

16

and

$$\int_0^t \frac{B_j(0,s)}{(t-s)^{1/2}} ds = 2 \int_0^{t^{1/2}} B_j(0,t-u^2) du$$

$$= \frac{1}{4} \int_0^{t^{1/2}} \left(\kappa_j(t-u^2) + \frac{1}{4} \right)^{-3/2} du$$

$$= 2 \int_0^{t^{1/2}} \left(4\kappa_j(t-u^2) + 1 \right)^{-3/2} du$$

$$= \frac{2t^{1/2}}{4\kappa_j t + 1}.$$

We consider the linear combination of the initial data

$$\mathcal{B}_0^{(i)}(x) = \sum_{j=1}^k \ell_j^{(i)} B_{0,j}(|x|)$$

so that the corresponding solution at x = 0 is

$$\mathcal{B}^{(i)}(0,t) = \sum_{j=1}^{k} \ell_j^{(i)} \frac{1}{8} \left(\kappa_j t + \frac{1}{4} \right)^{-3/2}$$

and

$$\int_0^t \frac{\mathcal{B}^{(i)}(0,s)}{(t-s)^{1/2}} ds = \sum_{i=1}^k \ell_j^{(i)} \frac{2t^{1/2}}{4\kappa_j t + 1}.$$
 (5.5)

Rearranging the constants $\tilde{\ell}_j^{(i)} = \frac{\ell_j^{(i)}}{2a_j}$ and $\tilde{\kappa}_j = \frac{1}{4\kappa_j}$, we denote

$$\Upsilon_j(t) = \frac{t^{1/2}}{t + \tilde{\kappa}_j}, \quad \Upsilon^{(i)}(t) = \sum_{j=1}^k \tilde{\ell}_j^{(i)} \frac{t^{1/2}}{t + \tilde{\kappa}_j}.$$
(5.6)

By adjusting free parameters $\tilde{\ell}_{j}^{(i)}$ and $\tilde{\kappa}_{j}$, we can find solutions $\Upsilon^{(i)}(t)$ with vanishing order $(T-t)^{i}$ near T. Indeed, writing $\hat{\Upsilon}^{(i)}(t) = \sum_{j=1}^{k} \tilde{\ell}_{j}^{(i)} \frac{1}{t+\tilde{\kappa}_{j}}$, $\Upsilon^{(i)}(t) \sim (T-t)^{i}$ is equivalent to showing the invertibility of the following system

$$Mv_{\ell} = e_i, \tag{5.7}$$

where

$$M = \begin{pmatrix} \frac{1}{T + \tilde{\kappa}_1} & \frac{1}{T + \tilde{\kappa}_1} & \cdots & \frac{1}{T + \tilde{\kappa}_k} \\ -\frac{1}{(T + \tilde{\kappa}_1)^2} & -\frac{1}{(T + \tilde{\kappa}_2)^2} & \cdots & -\frac{1}{(T + \tilde{\kappa}_k)^2} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{(-1)^k}{(T + \tilde{\kappa}_1)^k} & \frac{(-1)^k}{(T + \tilde{\kappa}_2)^k} & \cdots & \frac{(-1)^k}{(T + \tilde{\kappa}_k)^k} \end{pmatrix},$$

$$v_{\ell} = (\tilde{\ell}_1^{(i)}, \tilde{\ell}_2^{(i)}, \dots, \tilde{\ell}_k^{(i)})^T, \quad e_i = (0, \dots, 0, \underbrace{1}_{i-\text{th entry}}, 0, \dots, 0)^T.$$

Letting $\tilde{\kappa}_1, \ldots, \tilde{\kappa}_k$ be different from each other, it is obvious that system (5.7) is invertible. So we construct the solutions $\Upsilon^{(i)}(t)$ with vanishing order $(T-t)^i$ near T. Choosing a linear combination of k such functions and plugging them into the reduced equation, we obtain

$$\int_0^t \frac{\alpha(s)}{(t-s)^{1/2}} ds = \sum_{j=1}^k c_j \mathcal{B}^{(j)}(0,t) + h(t),$$

where c_i are free parameters in the initial data to be adjusted below. Then from Lemma 5.11, we have

$$\int_0^t \alpha(s)ds = \frac{1}{(\Gamma(\frac{1}{2}))^2} \left(\sum_{j=1}^k c_j \int_0^t \frac{\mathcal{B}^{(j)}(0,s)}{(t-s)^{1/2}} ds + \int_0^t \frac{h(s)}{(t-s)^{1/2}} ds \right). \tag{5.8}$$

We write

$$\int_0^t \frac{h(s)}{(t-s)^{1/2}} ds = d_0(T) + \sum_{j=1}^k d_j (T-t)^j + \text{h.o.t.}$$

We can choose c_j such that problem (5.8) has solution $\alpha(t) = o((T-t)^{k-1})$. In other words, our aim is to show

$$\alpha(T) = \alpha'(T) = \alpha''(T) = \dots = \alpha^{(k-1)}(T) = 0.$$
 (5.9)

From (5.8), (5.5) and (5.6), we obtain

$$\int_0^t \alpha(s)ds = \frac{1}{(\Gamma(\frac{1}{2}))^2} \left(\sum_{j=1}^k c_j \Upsilon^{(j)}(t) + \sum_{j=1}^k d_j (T-t)^j \right) + \text{h.o.t.}$$

By the vanishing order of $\Upsilon^{(j)}(t)$ and choosing $c_j = d_j$, we conclude the validity of (5.9). The proof is complete.

5.2. Reduced equation for $\alpha(t)$. By the orthogonality condition (5.1), we directly compute

$$\int_{B_{2R}} 5 \left[\left(u_{in}(\mu_0 y, t(\tau)) + \Phi_1(\mu_0 y, t(\tau)) \right)^4 - \mu_0^{-2} w^4(y) \right] \mu_0^2 \phi(y, \tau) Z_0(y) \, \mathrm{d}y
+ \int_{B_{2R}} 5 \mu_0^{\frac{1}{2}} (1 + \Lambda)^{-4} w^4 \left(\frac{y}{(1 + \Lambda)^2} \right) \psi(\mu_0 y, t(\tau)) Z_0(y) \, \mathrm{d}y
+ \int_{B_{2R}} \mu_0 \mu_0' \left[\frac{1}{2} \phi(y, \tau) + \nabla_y \phi(y, \tau) \cdot y \right] Z_0(y) \, \mathrm{d}y
+ \int_{B_{2R}} \mu_0^{\frac{5}{2}} \left[S(U_1) - \chi \left(\frac{\mu_0 y}{c_0 (T - t)^{\frac{1}{2}}} \right) \frac{\alpha(t)}{(\mu^2 + |\mu_0 y|^2)^{\frac{1}{2}}} \right] Z_0(y) \, \mathrm{d}y
+ \int_{B_{2R}} \mu_0^{\frac{5}{2}} \left[\left(u_{in}(\mu_0 y, t(\tau)) + \Phi_1(\mu_0 y, t(\tau)) \right)^5 - u_{in}^5(\mu_0 y, t(\tau)) \right] Z_0(y) \, \mathrm{d}y = 0.$$

Since

$$u_{in}(x,t) = \mu^{-\frac{1}{2}}w(\frac{x}{\mu}) + 2\mu'_0\mu^{\frac{1}{2}}J(\frac{x}{\mu}) = \mu^{-\frac{1}{2}}w(\frac{y}{(1+\Lambda)^2}) + 2\mu'_0\mu^{\frac{1}{2}}J(\frac{y}{(1+\Lambda)^2}),$$

we obtain

$$\int_{B_{2R}} 5 \left[\left(\mu^{-\frac{1}{2}} w \left(\frac{y}{(1+\Lambda)^2} \right) + 2\mu'_0 \mu^{\frac{1}{2}} J \left(\frac{y}{(1+\Lambda)^2} \right) + \Phi_1(\mu_0 y, t(\tau)) \right)^4 - \mu_0^{-2} w^4(y) \right] \mu_0^2 \phi(y, \tau) Z_0(y) \, \mathrm{d}y \\
+ \int_{B_{2R}} 5 \mu_0^{\frac{1}{2}} (1+\Lambda)^{-4} w^4 \left(\frac{y}{(1+\Lambda)^2} \right) \psi(\mu_0 y, t(\tau)) Z_0(y) \, \mathrm{d}y \\
+ \int_{B_{2R}} \mu_0 \mu'_0 \left[\frac{1}{2} \phi(y, \tau) + \nabla_y \phi(y, \tau) \cdot y \right] Z_0(y) \, \mathrm{d}y \\
+ \int_{B_{2R}} \mu_0^{\frac{5}{2}} \left[S(U_1) - \chi \left(\frac{\mu_0 y}{c_0 (T-t)^{\frac{1}{2}}} \right) \frac{\alpha(t)}{(\mu^2 + |\mu_0 y|^2)^{\frac{1}{2}}} \right] Z_0(y) \, \mathrm{d}y \\
+ \int_{B_{2R}} \mu_0^{\frac{5}{2}} \left[\left(u_{in} + \Phi_1(\mu_0 y, t(\tau)) \right)^5 - u_{in}^5 \right] Z_0(y) \, \mathrm{d}y = 0,$$

where we expand

$$\begin{aligned} & \left(u_{in} + \Phi_{1}(\mu_{0}y, t(\tau))\right)^{5} - u_{in}^{5} \\ &= 5u_{in}^{4}\Phi_{1}(\mu_{0}y, t(\tau)) + \Phi_{1}^{2}(\mu_{0}y, t(\tau)) \int_{0}^{1} 20\left(u_{in} + s\Phi_{1}(\mu_{0}y, t(\tau))\right)^{3}(1 - s) \,\mathrm{d}s \\ &= 5(\mu_{0}^{-\frac{1}{2}}w(y))^{4}\Phi_{1}(\mu_{0}y, t(\tau)) + 5\left[u_{in}^{4} - (\mu_{0}^{-\frac{1}{2}}w(y))^{4}\right]\Phi_{1}(\mu_{0}y, t(\tau)) \\ & + \Phi_{1}^{2}(\mu_{0}y, t(\tau)) \int_{0}^{1} 20\left(u_{in} + s\Phi_{1}(\mu_{0}y, t(\tau))\right)^{3}(1 - s) \,\mathrm{d}s \\ &= 5(\mu_{0}^{-\frac{1}{2}}w(y))^{4}\Phi_{1}(0, t(\tau)) + 5(\mu_{0}^{-\frac{1}{2}}w(y))^{4}\left(\Phi_{1}(\mu_{0}y, t(\tau)) - \Phi_{1}(0, t(\tau))\right) \\ & + 5\left[u_{in}^{4} - (\mu_{0}^{-\frac{1}{2}}w(y))^{4}\right]\Phi_{1}(\mu_{0}y, t(\tau)) + \Phi_{1}^{2}(\mu_{0}y, t(\tau)) \int_{0}^{1} 20\left(u_{in} + s\Phi_{1}(\mu_{0}y, t(\tau))\right)^{3}(1 - s) \,\mathrm{d}s \end{aligned}$$

and we shall prove the leading term is

$$5(\mu_0^{-\frac{1}{2}}w(y))^4\Phi_1(0,t(\tau)),$$

and all other terms have sufficiently fast decay. Indeed, we simplify the above equation and evaluate

$$\int_{B_{2R}} 5 \left[\left(\mu^{-\frac{1}{2}} w \left(\frac{y}{(1+\Lambda)^2} \right) + 2\mu'_0 \mu^{\frac{1}{2}} J \left(\frac{y}{(1+\Lambda)^2} \right) + \Phi_1(\mu_0 y, t(\tau)) \right)^4 - \mu_0^{-2} w^4(y) \right] \mu_0^{\frac{3}{2}} \phi(y, \tau) Z_0(y) \, \mathrm{d}y \\
+ \int_{B_{2R}} 5 w^4(y) \psi(\mu_0 y, t(\tau)) Z_0(y) \, \mathrm{d}y + \int_{B_{2R}} 5 \left[(1+\Lambda)^{-4} w^4 \left(\frac{y}{(1+\Lambda)^2} \right) - w^4(y) \right] \psi(\mu_0 y, t(\tau)) Z_0(y) \, \mathrm{d}y \\
+ \int_{B_{2R}} \mu_0^{\frac{1}{2}} \mu'_0 \left[\frac{1}{2} \phi(y, \tau) + \nabla_y \phi(y, \tau) \cdot y \right] Z_0(y) \, \mathrm{d}y \\
+ \int_{B_{2R}} \mu_0^2 \left[S(U_1) - \chi \left(\frac{\mu_0 y}{c_0 (T-t)^{\frac{1}{2}}} \right) \frac{\alpha(t)}{(\mu^2 + |\mu_0 y|^2)^{\frac{1}{2}}} \right] Z_0(y) \, \mathrm{d}y \\
+ \int_{B_{2R}} 5 w^4(y) \Phi_1(0, t(\tau)) Z_0(y) \, \mathrm{d}y \\
+ \int_{B_{2R}} \mu_0^2 \left[\left(u_{in} + \Phi_1(\mu_0 y, t(\tau)) \right)^5 - u_{in}^5 - 5(\mu_0^{-\frac{1}{2}} w(y))^4 \Phi_1(0, t(\tau)) \right] Z_0(y) \, \mathrm{d}y = 0.$$

Recall that

$$Z_0(y) := -\left[y \cdot \nabla w + \frac{w}{2}\right] = \frac{3^{\frac{1}{4}}}{2} \frac{|y|^2 - 1}{(1 + |y|^2)^{\frac{3}{2}}},$$
$$w(y) = 3^{\frac{1}{4}} (1 + |y|^2)^{-\frac{1}{2}}.$$

So

$$\int_{B_{2R}} 5w^4(y)Z_0(y) \, dy = \frac{5}{2} 3^{\frac{5}{4}} \int_0^{2R} (1+r^2)^{-3.5} (r^2 - 1) 4\pi r^2 \, dr$$

$$= 10\pi 3^{\frac{5}{4}} \frac{r^3(r^2 - 5)}{15(r^2 + 1)^{\frac{5}{2}}} \Big|_0^{2R}$$

$$= 10\pi 3^{\frac{5}{4}} (\frac{1}{15} + O(R^{-2})).$$

Next we consider the nonlocal term

$$\Phi_{1}(0,t) = \sum_{j=1}^{k} c_{j} \mathcal{B}^{(j)}(0,t) + \int_{0}^{t} \int_{\mathbb{R}^{3}} (\frac{1}{2\sqrt{\pi}})^{3} (t-s)^{-\frac{3}{2}} e^{-\frac{|\xi|^{2}}{4(t-s)}} \chi(\frac{\xi}{c_{0}(T-s)^{\frac{1}{2}}}) \frac{\alpha(s)}{\mu(s) + |\xi|} d\xi ds$$

$$= \sum_{j=1}^{k} c_{j} \mathcal{B}^{(j)}(0,t) + \int_{0}^{t} \int_{\mathbb{R}^{3}} (\frac{1}{2\sqrt{\pi}})^{3} (t-s)^{-\frac{3}{2}} e^{-\frac{|\xi|^{2}}{4(t-s)}} \chi(\frac{\xi}{c_{0}(T-s)^{\frac{1}{2}}}) \frac{\alpha(s)}{|\xi|} d\xi ds$$

$$+ \int_{0}^{t} \int_{\mathbb{R}^{3}} (\frac{1}{2\sqrt{\pi}})^{3} (t-s)^{-\frac{3}{2}} e^{-\frac{|\xi|^{2}}{4(t-s)}} \chi(\frac{\xi}{c_{0}(T-s)^{\frac{1}{2}}}) \alpha(s) (\frac{1}{\mu(s) + |\xi|} - \frac{1}{|\xi|}) d\xi ds.$$

Notice

$$\int_{0}^{t} \int_{\mathbb{R}^{3}} \left(\frac{1}{2\sqrt{\pi}}\right)^{3} (t-s)^{-\frac{3}{2}} e^{-\frac{|\xi|^{2}}{4(t-s)}} \chi\left(\frac{\xi}{c_{0}(T-s)^{\frac{1}{2}}}\right) \frac{\alpha(s)}{|\xi|} d\xi ds$$

$$= \int_{0}^{t} \int_{0}^{c_{0}(T-s)^{\frac{1}{2}}} \left(\frac{1}{2\sqrt{\pi}}\right)^{3} (t-s)^{-\frac{3}{2}} e^{-\frac{r^{2}}{4(t-s)}} \frac{\alpha(s)}{r} 4\pi r^{2} dr ds$$

$$= 4^{-1} \pi^{-\frac{1}{2}} \int_{0}^{t} \int_{0}^{c_{0}(T-s)^{\frac{1}{2}}} (t-s)^{-\frac{3}{2}} e^{-\frac{r^{2}}{4(t-s)}} \alpha(s) d(r^{2}) ds$$

$$= 4^{-1} \pi^{-\frac{1}{2}} \int_{0}^{t} (t-s)^{-\frac{3}{2}} \alpha(s) (-4) (t-s) e^{-\frac{r}{4(t-s)}} \Big|_{r=0}^{c_{0}^{2}(T-s)} ds$$

$$= \pi^{-\frac{1}{2}} \int_{0}^{t} (t-s)^{-\frac{1}{2}} \alpha(s) [1 - e^{-\frac{c_{0}^{2}(T-s)}{4(t-s)}}] ds.$$

Then

$$\Phi_{1}(0,t) = \sum_{j=1}^{k} c_{j} \mathcal{B}^{(j)}(0,t) + \pi^{-\frac{1}{2}} \int_{0}^{t} (t-s)^{-\frac{1}{2}} \alpha(s) [1 - e^{-\frac{c_{0}^{2}(T-s)}{4(t-s)}}] ds$$

$$+ \int_{0}^{t} \int_{\mathbb{R}^{3}} (\frac{1}{2\sqrt{\pi}})^{3} (t-s)^{-\frac{3}{2}} e^{-\frac{|\xi|^{2}}{4(t-s)}} \chi(\frac{\xi}{c_{0}(T-s)^{\frac{1}{2}}}) \alpha(s) (\frac{1}{\mu(s) + |\xi|} - \frac{1}{|\xi|}) d\xi ds$$

and thus

$$\begin{split} &\int\limits_{B_{2R}} 5 \left[\left(\mu^{-\frac{1}{2}} w (\frac{y}{(1+\Lambda)^2}) + 2 \mu_0' \, \mu^{\frac{1}{2}} J (\frac{y}{(1+\Lambda)^2}) + \Phi_1(\mu_0 y, t(\tau)) \right)^4 - \mu_0^{-2} w^4(y) \right] \mu_0^{\frac{3}{2}} \phi(y,\tau) Z_0(y) \, \mathrm{d}y \\ &+ \int\limits_{B_{2R}} 5 w^4(y) \psi(\mu_0 y, t(\tau)) Z_0(y) \, \mathrm{d}y + \int\limits_{B_{2R}} 5 \left[(1+\Lambda)^{-4} w^4 (\frac{y}{(1+\Lambda)^2}) - w^4(y) \right] \psi(\mu_0 y, t(\tau)) Z_0(y) \, \mathrm{d}y \\ &+ \int\limits_{B_{2R}} \mu_0^{\frac{1}{2}} \left[\frac{1}{2} \phi(y,\tau) + \nabla_y \phi(y,\tau) \cdot y \right] Z_0(y) \, \mathrm{d}y \\ &+ \int\limits_{B_{2R}} \mu_0^2 \left[S(U_1) - \chi (\frac{\mu_0 y}{c_0(T-t)^{\frac{1}{2}}}) \frac{\alpha(t)}{(\mu^2 + |\mu_0 y|^2)^{\frac{1}{2}}} \right] Z_0(y) \, \mathrm{d}y \\ &+ 10\pi 3^{\frac{5}{4}} (\frac{1}{15} + O(R^{-2})) (\sum_{j=1}^k c_j \mathcal{B}^{(j)}(0,t)) \\ &+ 10\pi 3^{\frac{5}{4}} (\frac{1}{15} + O(R^{-2})) \pi^{-\frac{1}{2}} \int_0^t (t-s)^{-\frac{1}{2}} \alpha(s) [1 - e^{-\frac{c_0^2(T-s)}{4(t-s)}}] \, \mathrm{d}s \\ &+ 10\pi 3^{\frac{5}{4}} (\frac{1}{15} + O(R^{-2})) \int_0^t \int\limits_{\mathbb{R}^3} (\frac{1}{2\sqrt{\pi}})^3 (t-s)^{-\frac{3}{2}} e^{-\frac{|\xi|^2}{4(t-s)}} \chi (\frac{\xi}{c_0(T-s)^{\frac{1}{2}}}) (\frac{1}{\mu(s) + |\xi|} - \frac{1}{|\xi|}) \, \mathrm{d}\xi \, \mathrm{d}s \\ &+ \int\limits_{B_{2R}} \mu_0^2 \left[\left(u_{in} + \Phi_1(\mu_0 y, t(\tau)) \right)^5 - u_{in}^5 - 5(\mu_0^{-\frac{1}{2}} w(y))^4 \Phi_1(0, t(\tau)) \right] Z_0(y) \, \mathrm{d}y = 0. \end{split}$$

The ansatz for the parameter function is

$$\alpha(t) := (-3^{\frac{1}{4}})\mu_0^{-\frac{1}{2}}(\mu_0\Lambda(t))' \to 0 \text{ as } t \to T$$

which is possible provided

$$\alpha(t) \sim (T-t)^{-k} (T-t)^{2k} \Lambda(t) (T-t)^{-1} \sim (T-t)^{k-1} \Lambda(t)$$

Next we compute

$$\int_{0}^{t} (t-s)^{-\frac{1}{2}} \alpha(s) \, ds$$

$$= \pi^{\frac{1}{2}} \sum_{j=1}^{k} c_{j} \mathcal{B}^{(j)}(0,t) + O(R^{-2}) \pi^{\frac{1}{2}} \sum_{j=1}^{k} c_{j} \mathcal{B}^{(j)}(0,t) + \int_{0}^{t} (t-s)^{-\frac{1}{2}} \alpha(s) e^{-\frac{c_{0}^{2}(T-s)}{4(t-s)}} \, ds$$

$$- \int_{B_{2R}} \frac{5}{2\pi^{\frac{1}{2}} 3^{\frac{1}{4}}} \left[\left(\mu^{-\frac{1}{2}} w \left(\frac{y}{(1+\Lambda)^{2}} \right) + 2\mu'_{0} \mu^{\frac{1}{2}} J \left(\frac{y}{(1+\Lambda)^{2}} \right) + \Phi_{1}(\mu_{0}y, t(\tau)) \right)^{4} - \mu_{0}^{-2} w^{4}(y) \right]$$

$$\times \mu_{0}^{\frac{3}{2}} \phi(y, \tau) Z_{0}(y) \, dy - \int_{B_{2R}} \frac{5}{2\pi^{\frac{1}{2}} 3^{\frac{1}{4}}} w^{4}(y) \psi(\mu_{0}y, t(\tau)) Z_{0}(y) \, dy$$

$$- \int_{B_{2R}} \frac{5}{2\pi^{\frac{1}{2}} 3^{\frac{1}{4}}} \left[(1+\Lambda)^{-4} w^{4} \left(\frac{y}{(1+\Lambda)^{2}} \right) - w^{4}(y) \right] \psi(\mu_{0}y, t(\tau)) Z_{0}(y) \, dy$$

$$- \int_{B_{2R}} \frac{1}{2\pi^{\frac{1}{2}} 3^{\frac{1}{4}}} \mu_{0}^{\frac{1}{2}} \mu_{0}^{\frac{1}{2}} \left[\frac{1}{2} \phi(y, \tau) + \nabla_{y} \phi(y, \tau) \cdot y \right] Z_{0}(y) \, dy$$

$$+ \int_{B_{2R}} \frac{1}{2\pi^{\frac{1}{2}} 3^{\frac{1}{4}}} \mu_{0}^{2} \left[S(U_{1}) - \chi \left(\frac{\mu_{0}y}{c_{0}(T-t)^{\frac{1}{2}}} \right) \frac{\alpha(t)}{(\mu^{2} + |\mu_{0}y|^{2})^{\frac{1}{2}}} \right] Z_{0}(y) \, dy$$

$$- O(R^{-2}) \int_{0}^{t} (t-s)^{-\frac{1}{2}} \alpha(s) \left[1 - e^{-\frac{c_{0}^{2}(T-s)}{4(t-s)}} \right] ds$$

$$- (1 + O(R^{-2})) \pi^{\frac{1}{2}} \int_{0}^{t} \int_{\mathbb{R}^{3}} \left(\frac{1}{2\sqrt{\pi}} \right)^{3} (t-s)^{-\frac{3}{2}} e^{-\frac{|\xi|^{2}}{4(t-s)}} \chi \left(\frac{\xi}{c_{0}(T-s)^{\frac{1}{2}}} \right) \alpha(s) \left(\frac{1}{\mu(s) + |\xi|} - \frac{1}{|\xi|} \right) d\xi \, ds$$

$$- \int_{B_{2R}} \frac{1}{2\pi^{\frac{1}{2}} 3^{\frac{1}{4}}} \mu_{0}^{2} \left[\left(u_{in} + \Phi_{1}(\mu_{0}y, t(\tau)) \right)^{5} - u_{in}^{5} - 5(\mu_{0}^{-\frac{1}{2}} w(y))^{4} \Phi_{1}(0, t(\tau)) \right] Z_{0}(y) \, dy.$$
(5.10)

From the theory on Riemann-Liouville fractional differential operator (see [11] for instance), we have

$$\frac{1}{\Gamma(\frac{1}{2})} \int_{0}^{t} (t-y)^{-\frac{1}{2}} \frac{1}{\Gamma(\frac{1}{2})} \int_{0}^{y} (y-s)^{-\frac{1}{2}} \alpha(s) \, ds \, dy$$

$$= \frac{1}{\Gamma(\frac{1}{2})} \int_{0}^{t} (t-y)^{-\frac{1}{2}} \frac{1}{\Gamma(\frac{1}{2})} \int_{0}^{t} \chi[s \le y] (y-s)^{-\frac{1}{2}} \alpha(s) \, ds \, dy$$

$$= \frac{1}{(\Gamma(\frac{1}{2}))^{2}} \int_{0}^{t} \int_{s}^{t} (t-s)^{-1} (\frac{t-y}{t-s})^{-\frac{1}{2}} (\frac{y-s}{t-s})^{-\frac{1}{2}} \alpha(s) \, dy \, ds$$

$$= \int_{0}^{t} \alpha(s) \, ds. \tag{5.11}$$

Therefore, we obtain

Changing the variable $(t-s)^{1/2} = u$, we have

$$\begin{split} &\int_{0}^{t} \alpha(s) \, \mathrm{d}s \\ &= \frac{1}{\Gamma(\frac{1}{2})} \sum_{j=1}^{k} c_{j} \int_{0}^{t} (t-a)^{-\frac{1}{2}} \mathcal{B}^{(j)}(0,a) \, \mathrm{d}a + \frac{2}{\Gamma(\frac{1}{2})} \sum_{j=1}^{k} c_{j} \int_{0}^{t^{\frac{1}{2}}} O(R^{-2}(t-u^{2})) \mathcal{B}^{(j)}(0,t-u^{2}) \, \mathrm{d}u \\ &+ \frac{2}{\Gamma^{2}(\frac{1}{2})} \int_{0}^{t^{\frac{1}{2}}} \int_{0}^{t^{\frac{1}{2}}} \int_{0}^{t^{\frac{1}{2}}} \int_{B_{2R(t-u^{2})}} \left[(\mu^{-\frac{1}{2}}(t-u^{2})w(\frac{y}{(1+\Lambda(t-u^{2}))^{2}}) + \Phi_{1}(\mu_{0}(t-u^{2})y,t-u^{2}))^{d} - \mu_{0}^{-2}(t-u^{2})w^{4}(y) \right] \\ &+ 2\mu_{0}(t-u^{2})\mu^{\frac{1}{2}}(t-u^{2})J(\frac{y}{(1+\Lambda(t-u^{2}))^{2}}) + \Phi_{1}(\mu_{0}(t-u^{2})y,t-u^{2}))^{d} - \mu_{0}^{-2}(t-u^{2})w^{4}(y) \right] \\ &\times \mu_{0}^{\frac{1}{2}}(t-u^{2})\phi(y,t-u^{2})Z_{0}(y) \, \mathrm{d}y \, \mathrm{d}u \\ &- \frac{5}{\pi^{\frac{1}{2}}3^{\frac{1}{2}}} \int_{0}^{t^{\frac{1}{2}}} \int_{B_{2R(t-u^{2})}} w^{4}(y)\psi(\mu_{0}(t-u^{2})y,t-u^{2})Z_{0}(y) \, \mathrm{d}y \, \mathrm{d}u \\ &- \frac{5}{\pi^{\frac{1}{2}}3^{\frac{1}{2}}} \int_{0}^{t^{\frac{1}{2}}} \int_{B_{2R(t-u^{2})}} \left[(1+\Lambda(t-u^{2}))^{-4}w^{4}(\frac{y}{(1+\Lambda(t-u^{2}))^{2}}) - w^{4}(y) \right] \\ &\times \psi(\mu_{0}(t-u^{2})y,t-u^{2})Z_{0}(y) \, \mathrm{d}y \, \mathrm{d}u \\ &- \frac{1}{\pi^{\frac{1}{2}}3^{\frac{1}{2}}} \int_{0}^{t^{\frac{1}{2}}} \int_{B_{2R(t-u^{2})}} \mu_{0}^{2}(t-u^{2})\mu'_{0}(t-u^{2})\left[\frac{1}{2}\phi(y,t-u^{2}) + \nabla_{y}\phi(y,t-u^{2}) \cdot y\right]Z_{0}(y) \, \mathrm{d}y \, \mathrm{d}u \\ &+ \frac{1}{\pi^{\frac{3}{2}}3^{\frac{1}{2}}} \int_{0}^{t^{\frac{1}{2}}} \int_{B_{2R(t-u^{2})}} \mu_{0}^{2}(t-u^{2}) \times \left[S(U_{1})(\mu_{0}(t-u^{2})y,t-u^{2}) - u^{2}\right] \\ &- \chi(\frac{\mu_{0}(t-u^{2})y}{c_{0}(T-t+u^{2})^{\frac{1}{2}}}) \frac{\alpha(t-u^{2})}{(\mu^{2}(t-u^{2}) + |\mu_{0}(t-u^{2})y|^{\frac{1}{2}}}\right] Z_{0}(y) \, \mathrm{d}y \, \mathrm{d}u \\ &- \frac{2}{\pi^{\frac{1}{2}}} \int_{0}^{t^{\frac{1}{2}}} O(R^{-2}(t-u^{2})) \int_{0}^{t-u^{2}} (t-u^{2}-s)^{-\frac{3}{2}} e^{-\frac{|\mu|^{2}}{4(t-u^{2}-s)^{\frac{1}{2}}}} \chi(\frac{\xi}{c_{0}(T-s)^{\frac{1}{2}}})\alpha(s) \left(\frac{1}{\mu(s)+|\xi|} - \frac{1}{|\xi|}\right) \, \mathrm{d}\xi \, \mathrm{d}s \, \mathrm{d}u \\ &- \frac{1}{\pi^{\frac{3}{2}}3^{\frac{1}{2}}} \int_{0}^{t^{\frac{1}{2}}} \int_{B_{2R(t-u^{2})}} \mu_{0}^{2}(t-u^{2}) \times \left[\left(u_{in}(y,t-u^{2}) + \Phi_{1}(\mu_{0}(t-u^{2})y,t-u^{2})\right)^{5} \\ &- u_{in}^{5}(y,t-u^{2}) - 5(\mu_{0}^{\frac{1}{2}}(t-u^{2})w(y))^{4} \Phi_{1}(0,t-u^{2}) \right] Z_{0}(y) \, \mathrm{d}y \, \mathrm{d}u. \end{split}$$

Now we check the differentiability of right hand side in (5.12) term by term. First, we consider

$$\int_0^{t^{\frac{1}{2}}} O(R^{-2}(t-u^2)) \mathcal{B}^{(j)}(0,t-u^2) \, \mathrm{d}u.$$

This term is smooth about t near T and we have the following estimate:

$$\partial_t^{(i)} \int_0^{t^{\frac{1}{2}}} O(R^{-2}(t-u^2)) \mathcal{B}^{(j)}(0,t-u^2) \, \mathrm{d}u \Big|_{t=T} \sim O(T^{\frac{1}{2}-i}).$$

Next, we have

$$\partial_t \left(\int_0^{t^{\frac{1}{2}}} \int_0^{t-u^2} (t - u^2 - s)^{-\frac{1}{2}} \alpha(s) e^{-\frac{c_0^2(T-s)}{4(t-u^2-s)}} \, \mathrm{d}s \, \mathrm{d}u \right)$$

$$= \int_0^{t^{\frac{1}{2}}} \int_0^{t-u^2} \partial_t \left((t - u^2 - s)^{-\frac{1}{2}} \alpha(s) e^{-\frac{c_0^2(T-s)}{4(t-u^2-s)}} \right) \, \mathrm{d}s \, \mathrm{d}u.$$

Similarly, we can check that this term is smooth about t near T and have the following estimate:

$$\partial_t^{(i)} \left(\int_0^{t^{\frac{1}{2}}} \int_0^{t-u^2} (t-u^2-s)^{-\frac{1}{2}} \alpha(s) e^{-\frac{c_0^2(T-s)}{4(t-u^2-s)}} \, \mathrm{d}s \, \mathrm{d}u \right) \Big|_{t=T} \sim o(T^{\frac{1}{2}-i}).$$

Then, the term

$$\int_{0}^{t^{\frac{1}{2}}} \int_{B_{2R(t-u^2)}} \left[\left(\mu^{-\frac{1}{2}}(t-u^2)w(\frac{y}{(1+\Lambda(t-u^2))^2}) + 2\mu'_0(t-u^2)\mu^{\frac{1}{2}}(t-u^2)J(\frac{y}{(1+\Lambda(t-u^2))^2}) \right. \\ \left. + \Phi_1(\mu_0(t-u^2)\ y,t-u^2) \right)^4 - \mu_0^{-2}(t-u^2)w^4(y) \right] \times \mu_0^{\frac{3}{2}}(t-u^2)\phi(y,t-u^2)Z_0(y)\,\mathrm{d}y\,\mathrm{d}u$$

is C_t^k for t near T if we have $\phi \in C_{x,t}^{2k,k}$, and we have the following estimate

$$\partial_{t}^{(i)} \int_{0}^{t^{\frac{1}{2}}} \int_{B_{2R(t-u^{2})}} \left[\left(\mu^{-\frac{1}{2}}(t-u^{2})w(\frac{y}{(1+\Lambda(t-u^{2}))^{2}}) + 2\mu'_{0}(t-u^{2})\mu^{\frac{1}{2}}(t-u^{2})J(\frac{y}{(1+\Lambda(t-u^{2}))^{2}}) \right. \\ \left. + \Phi_{1}(\mu_{0}(t-u^{2})\ y,t-u^{2})\right)^{4} - \mu_{0}^{-2}(t-u^{2})w^{4}(y) \right] \\ \left. \times \mu_{0}^{\frac{3}{2}}(t-u^{2})\phi(y,t-u^{2})Z_{0}(y) \,\mathrm{d}y \,\mathrm{d}u \right|_{t-T} = o(T^{\frac{1}{2}-i}).$$

For the term

$$\int_0^{t^{\frac{1}{2}}} \int_{B_{2R(t-u^2)}} w^4(y)\psi(\mu_0(t-u^2)y, t-u^2) Z_0(y) \, \mathrm{d}y \, \mathrm{d}u,$$

it is C^k for t near T if we have $\psi \in C^{2k,k}_{x,t}$. Further, we have the estimate

$$\partial_t^{(i)} \int_0^{t^{\frac{1}{2}}} \int_{B_{2R(t-u^2)}} w^4(y) \psi(\mu_0(t-u^2)y, t-u^2) Z_0(y) \, \mathrm{d}y \, \mathrm{d}u \Big|_{t=T} = o(T^{\frac{1}{2}-i}).$$

Next, we consider

$$\int_0^{t^{\frac{1}{2}}} \int_{B_{2R(t-u^2)}} \left[(1 + \Lambda(t-u^2))^{-4} w^4 (\frac{y}{(1 + \Lambda(t-u^2))^2}) - w^4(y) \right] \psi(\mu_0(t-u^2)y, t-u^2) Z_0(y) \, dy \, du,$$

which is C_t^k if we assume $\psi \in C_{x,t}^{2k,k}$, and

$$\partial_t^{(i)} \int_0^{t^{\frac{1}{2}}} \int_{B_{2R(t-u^2)}} \left[(1 + \Lambda(t-u^2))^{-4} w^4 (\frac{y}{(1 + \Lambda(t-u^2))^2}) - w^4(y) \right] \\ \psi(\mu_0(t-u^2)y, t-u^2) Z_0(y) \, \mathrm{d}y \, \mathrm{d}u \Big|_{t=T} \sim O(T^{\frac{1}{2}-i}).$$

Next we consider

$$\int_0^{t^{\frac{1}{2}}} \int_{B_{2R(t-u^2)}} \mu_0^{\frac{1}{2}}(t-u^2) \mu_0'(t-u^2) \left[\frac{1}{2} \phi(y,t-u^2) + \nabla_y \phi(y,t-u^2) \cdot y \right] Z_0(y) \, \mathrm{d}y \, \mathrm{d}u$$

It is C^k_t about t near T if we have $\phi \in C^{2k,k}_{x,t}$ and we have the following estimate

$$\partial_t^{(i)} \int_0^{t^{\frac{1}{2}}} \int_{B_{2R(t-u^2)}} \mu_0^{\frac{1}{2}}(t-u^2) \mu_0'(t-u^2) \left[\frac{1}{2} \phi(y,t-u^2) + \nabla_y \phi(y,t-u^2) \cdot y \right] Z_0(y) \, \mathrm{d}y \, \mathrm{d}u \Big|_{t=T} \sim o(T^{\frac{1}{2}-i}).$$

Next, for the term

$$\int_0^{t^{\frac{1}{2}}} \int_{B_{2R(t-u^2)}} \mu_0^2(t-u^2) \times \left[S(U_1)(\mu_0(t-u^2) \ y, t-u^2) - \chi(\frac{\mu_0(t-u^2)y}{c_0(T-t+u^2)^{\frac{1}{2}}}) \frac{\alpha(t-u^2)}{(\mu^2(t-u^2) + |\mu_0(t-u^2)y|^2)^{\frac{1}{2}}} \right] Z_0(y) \, \mathrm{d}y \, \mathrm{d}u,$$

it is C_t^k if we assume $\Lambda \in C_t^{k+1}$, and we have

$$\partial_t^{(i)} \int_0^{t^{\frac{1}{2}}} \int_{B_{2R(t-u^2)}} \mu_0^2(t-u^2) \times \left[S(U_1)(\mu_0(t-u^2) \ y, t-u^2) \right.$$
$$\left. - \chi(\frac{\mu_0(t-u^2)y}{c_0(T-t+u^2)^{\frac{1}{2}}}) \frac{\alpha(t-u^2)}{(\mu^2(t-u^2) + |\mu_0(t-u^2)y|^2)^{\frac{1}{2}}} \right] Z_0(y) \, \mathrm{d}y \, \mathrm{d}u \Big|_{t=T} \sim o(T^{\frac{1}{2}-i}).$$

We next consider

$$\int_0^{t^{\frac{1}{2}}} O(R^{-2}(t-u^2)) \int_0^{t-u^2} (t-u^2-s)^{-\frac{1}{2}} \alpha(s) [1 - e^{-\frac{c_0^2(T-s)}{4(t-u^2-s)}}] \, \mathrm{d}s \, \mathrm{d}u.$$

It is smooth for t near T and we have the following estimate

$$\partial_t^{(i)} \int_0^{t^{\frac{1}{2}}} O(R^{-2}(t-u^2)) \int_0^{t-u^2} (t-u^2-s)^{-\frac{1}{2}} \alpha(s) \left[1 - e^{-\frac{c_0^2(T-s)}{4(t-u^2-s)}}\right] ds du \Big|_{t=T} \sim o(T^{\frac{1}{2}-i}).$$

Next,

$$\int_{0}^{t^{\frac{1}{2}}} (1 + O(R^{-2}(t - u^{2}))) \times \int_{0}^{t-u^{2}} \int_{\mathbb{R}^{3}} (\frac{1}{2\sqrt{\pi}})^{3} (t - u^{2} - s)^{-\frac{3}{2}} e^{-\frac{|\xi|^{2}}{4(t - u^{2} - s)}} \chi(\frac{\xi}{c_{0}(T - s)^{\frac{1}{2}}}) \alpha(s) (\frac{1}{\mu(s) + |\xi|} - \frac{1}{|\xi|}) d\xi ds du$$

It is C_t^k about t near T if we assume $\Lambda \in C_t^{k+1}$ and the following estimate holds

$$\partial_t^{(i)} \int_0^{t^{\frac{1}{2}}} (1 + O(R^{-2}(t - u^2))) \int_0^{t - u^2} \int_{\mathbb{R}^3} (\frac{1}{2\sqrt{\pi}})^3 (t - u^2 - s)^{-\frac{3}{2}} e^{-\frac{|\xi|^2}{4(t - u^2 - s)}} \chi(\frac{\xi}{c_0(T - s)^{\frac{1}{2}}}) \times \alpha(s) (\frac{1}{\mu(s) + |\xi|} - \frac{1}{|\xi|}) d\xi ds du \Big|_{t = T} \sim o(T^{\frac{1}{2} - i}).$$

Next,

$$\int_{0}^{t^{\frac{1}{2}}} \int_{B_{2R(t-u^2)}} \mu_0^2(t-u^2) \times \left[\left(u_{in}(y,t-u^2) + \Phi_1(\mu_0(t-u^2)y,t-u^2) \right)^5 - u_{in}^5(y,t-u^2) - 5(\mu_0^{-\frac{1}{2}}(t-u^2)w(y))^4 \Phi_1(0,t-u^2) \right] Z_0(y) \, \mathrm{d}y \, \mathrm{d}y$$

It is C_t^k about t near T if $\Lambda \in C_t^k$, and

$$\partial_t^{(i)} \int_0^{t^{\frac{1}{2}}} \int_{B_{2R(t-u^2)}} \mu_0^2(t-u^2) \times \left[\left(u_{in}(y,t-u^2) + \Phi_1(\mu_0(t-u^2)y,t-u^2) \right)^5 - u_{in}^5(y,t-u^2) - 5(\mu_0^{-\frac{1}{2}}(t-u^2)w(y))^4 \Phi_1(0,t-u^2) \right] Z_0(y) \, \mathrm{d}y \, \mathrm{d}u \Big|_{t=T} \sim o(T^{\frac{1}{2}-i}).$$

Then, we have the following equation

$$\alpha(t) = \partial_t \left[\frac{1}{\Gamma(\frac{1}{2})} \sum_{j=1}^k c_j \int_0^t (t-a)^{-\frac{1}{2}} \mathcal{B}^{(j)}(0,a) \, \mathrm{d}a + \frac{2}{\Gamma^2(\frac{1}{2})} \int_0^{t^{\frac{1}{2}}} h[\vec{c}, \Lambda, \psi, \phi](t-u^2) \, \mathrm{d}u \right]$$
(5.13)

and thus

$$\Lambda(t) = \mu_0^{-1}(t) \int_t^T 3^{-\frac{1}{4}} \mu_0^{\frac{1}{2}}(b) \partial_b \left[\frac{1}{\Gamma(\frac{1}{2})} \sum_{j=1}^k c_j \int_0^b (b-a)^{-\frac{1}{2}} \mathcal{B}^{(j)}(0,a) da + \frac{2}{\Gamma^2(\frac{1}{2})} \int_0^{b^{\frac{1}{2}}} h[\vec{c}, \Lambda, \psi, \phi](b-u^2) du \right] db.$$
(5.14)

Define the space

$$\mathcal{X}_{\Lambda} := \left\{ \Lambda(t) : \|\Lambda(t)(T-t)^{-1+\varepsilon}\|_{L^{\infty}} \le C_{0\Lambda}, \|\Lambda^{(1)}(t)(T-t)^{\varepsilon}\|_{L^{\infty}} \le C_{1\Lambda}, \dots, \|\Lambda^{(k)}(t)(T-t)^{k-1+\varepsilon}\|_{L^{\infty}} \le C_{k\Lambda}, \Lambda(t) \in C^{k+1,\rho}(0,T-\delta), \text{ for all } \delta \in (0,T) \right\}$$
(5.15)

where $C_{0\Lambda}, C_{1\Lambda}, \dots, C_{k\Lambda}$ are some fixed small constants and ε is a small positive constant.

$$\mathcal{X}_c := \left\{ \vec{c} = (c_1, c_2, \dots, c_k) : |c_j| \le C_c T^{\frac{1}{2} - j - \varepsilon}, \ j = 1, 2, \dots, k \right\}$$
 (5.16)

where C_c is a fixed constant.

We aim to solve (5.14) for $\Lambda \in \mathcal{X}_{\Lambda}$, $\vec{c} \in \mathcal{X}_{c}$ by Schauder fixed point theorem.

For all $\Lambda_1 \in \mathcal{X}_{\Lambda}$, $\vec{c}_1 \in \mathcal{X}_c$, we want to find the unique \vec{c}_2 to get $\Lambda_2 \in \mathcal{X}_{\Lambda}$ satisfying the following equation:

$$\Lambda_{2}(t) = \mu_{0}^{-1}(t) \int_{t}^{T} 3^{-\frac{1}{4}} \mu_{0}^{\frac{1}{2}}(b) \partial_{b} \left[\frac{1}{\Gamma(\frac{1}{2})} \sum_{j=1}^{k} c_{2j} \int_{0}^{b} (b-a)^{-\frac{1}{2}} \mathcal{B}^{(j)}(0,a) da + \frac{2}{\Gamma^{2}(\frac{1}{2})} \int_{0}^{b^{\frac{1}{2}}} h[\vec{c}_{1}, \Lambda_{1}, \psi[\Lambda_{1}], \phi[\Lambda_{1}]](b-u^{2}) du \right] db.$$
(5.17)

Since we expect $\Lambda_2 \in \mathcal{X}_{\lambda}$, we have to choose suitable c_{2j} to cancel the lower power of T-t on the right hand side:

$$c_{2j} = -\frac{2}{\Gamma(\frac{1}{2})j!} \partial_b^j \left(\int_0^{b^{\frac{1}{2}}} h[\vec{c}_1, \Lambda_1, \psi[\Lambda_1], \phi[\Lambda_1]](b - u^2) \, \mathrm{d}u \right) \Big|_{b=T} = O(T^{\frac{1}{2} - j}).$$

The higher derivatives of h are well defined here since $\Lambda \in \mathcal{X}_{\Lambda}$ and $\alpha(t) \in C^k_t(0, T - \delta)$. We can use Schauder estimate to improve the regularity of ϕ , ψ to $C^{2k+2+2\rho,k+1+\rho}_{x,t}$ for $t \in (0, T - \delta)$ for any small δ . So h also has higher derivatives up to order k.

Taking higher derivatives for (5.17) and choosing T small enough, we have $\Lambda_2 \in \mathcal{X}_{\Lambda}$. Next, we show the Lipschitz continuity of Λ_2 . For any $0 \le t_1 < t_2 \le T - \delta$,

$$\begin{aligned} & \left| \Lambda_{2}(t_{1}) - \Lambda_{2}(t_{2}) \right| \\ &= \left| \mu_{0}^{-1}(t_{1}) \int_{t_{1}}^{t_{2}} 3^{-\frac{1}{4}} \mu_{0}^{\frac{1}{2}}(b) \partial_{b} \left[\frac{1}{\Gamma(\frac{1}{2})} \sum_{j=1}^{k} c_{j2} \int_{0}^{b} (b-a)^{-\frac{1}{2}} \mathcal{B}^{(j)}(0,a) \, \mathrm{d}a \right. \\ &\quad + \frac{2}{\Gamma^{2}(\frac{1}{2})} \int_{0}^{b^{\frac{1}{2}}} h[\vec{c}_{1}, \Lambda_{1}, \psi[\Lambda_{1}], \phi[\Lambda_{1}]](b-u^{2}) \, \mathrm{d}u \right] \, \mathrm{d}b \\ &\quad + (\mu_{0}^{-1}(t_{1}) - \mu^{-1}(t_{2})) \times \int_{t_{2}}^{T} 3^{-\frac{1}{4}} \mu_{0}^{\frac{1}{2}}(b) \partial_{b} \left[\frac{1}{\Gamma(\frac{1}{2})} \sum_{j=1}^{k} c_{j2} \int_{0}^{b} (b-a)^{-\frac{1}{2}} \mathcal{B}^{(j)}(0,a) \, \mathrm{d}a \right. \\ &\quad + \frac{2}{\Gamma^{2}(\frac{1}{2})} \int_{0}^{b^{\frac{1}{2}}} h[\vec{c}_{1}, \Lambda_{1}, \psi[\Lambda_{1}], \phi[\Lambda_{1}]](b-u^{2}) \, \mathrm{d}u \right] \, \mathrm{d}b \\ &\lesssim (T-t_{1})^{-2k} \int_{t_{1}}^{t_{2}} (T-b)^{2k} \, \mathrm{d}b + (T-\theta t_{1} - (1-\theta)t_{2})^{-2k-1} (t_{2}-t_{1}) \int_{t_{2}}^{T} (T-b)^{2k} \, \mathrm{d}b \\ &\leq t_{2}-t_{1}. \end{aligned}$$

Similarly, we can take k-th derivative of (5.17) to prove $\Lambda_2^{(k)}(t) \in \text{Lip}(0, T - \delta)$, $\forall \delta \in (0, T)$. We shall solve the full inner–outer gluing system together with the reduced problem (5.1) in Section

6. Solving the inner-outer gluing system

In this section, we will solve the inner outer gluing system (3.3) and (3.5) by using the linear theories developed in Section 4 together with the fixed point argument. We first estimate the right hand sides of problems (3.3) and (3.5) under the topology chosen in Section 4.

6.1. The outer problem: estimate of \mathcal{G} . Recall from (3.4) that

6.

$$\mathcal{G}(\phi,\psi,\alpha) = -\partial_t \eta_R \mu_0^{-\frac{1}{2}} \phi(\frac{x}{\mu_0},t) + \Delta_x \eta_R \mu_0^{-\frac{1}{2}} \phi(\frac{x}{\mu_0},t) + 2\nabla_x \eta_R \cdot \mu_0^{-\frac{3}{2}} \nabla_y \phi(\frac{x}{\mu_0},t)$$

$$+ \left(5(U_1 + \Phi_1)^4 (1 - \eta_R) + 5 \left[(U_1 + \Phi_1)^4 - \left(\mu^{-\frac{1}{2}} w(\frac{x}{\mu})\right)^4 \right] \eta_R \right) \psi$$

$$+ (U_1 + \Phi_1 + \Phi_2)^5 - (U_1 + \Phi_1)^5 - 5(U_1 + \Phi_1)^4 \Phi_2$$

$$+ \left[S(U_1) - \chi \left(\frac{x}{c_0 (T - t)^{\frac{1}{2}}}\right) \frac{\alpha(t)}{(\mu^2 + |x|^2)^{\frac{1}{2}}} \right] (1 - \eta_R) + \left[(U_1 + \Phi_1)^5 - U_1^5 \right] (1 - \eta_R).$$

We write

$$g_{1} := -\partial_{t}\eta_{R}\mu_{0}^{-\frac{1}{2}}\phi(\frac{x}{\mu_{0}},t) + \Delta_{x}\eta_{R}\mu_{0}^{-\frac{1}{2}}\phi(\frac{x}{\mu_{0}},t) + 2\nabla_{x}\eta_{R}\cdot\mu_{0}^{-\frac{3}{2}}\nabla_{y}\phi(\frac{x}{\mu_{0}},t),$$

$$g_{2} := \left(5(U_{1}+\Phi_{1})^{4}(1-\eta_{R}) + 5\left[(U_{1}+\Phi_{1})^{4} - \left(\mu^{-\frac{1}{2}}w(\frac{x}{\mu})\right)^{4}\right]\eta_{R}\right)\psi,$$

$$g_{3} := (U_{1}+\Phi_{1}+\Phi_{2})^{5} - (U_{1}+\Phi_{1})^{5} - 5(U_{1}+\Phi_{1})^{4}\Phi_{2},$$

$$g_{4} := \left[S(U_{1}) - \chi(\frac{x}{c_{0}(T-t)^{\frac{1}{2}}})\frac{\alpha(t)}{(\mu^{2}+|x|^{2})^{\frac{1}{2}}}\right](1-\eta_{R}) + \left[(U_{1}+\Phi_{1})^{5} - U_{1}^{5}\right](1-\eta_{R}).$$

$$(6.1)$$

Estimate of g_1

Since $\|\phi\|_{0,\nu,\sigma} < +\infty$, we have

$$|g_1| \lesssim \mu_0^{\nu - \frac{5}{2}}(t)R^{-1-\sigma}(t)\chi_{\{|x| \sim \mu_0 R\}}$$

and thus for some $\epsilon > 0$

$$||g_1||_{**} \lesssim T^{\epsilon} (1 + ||\phi||_{0,\nu,\sigma})$$
 (6.2)

provided

$$1 + a - \sigma < 0. \tag{6.3}$$

Estimate of g_2

Since $\|\psi\|_* < +\infty$ and we choose the initial data such that $\psi(0,T) = 0$, we have

$$\begin{aligned} |\psi(x,t)| &= |\psi(x,t) - \psi(0,T)| \\ &\leq |\psi(x,t) - \psi(x,T)| + |\psi(x,T) - \psi(0,T)| \\ &\lesssim \mu_0^{\nu - \frac{1}{2}}(t) R^{-a}(t) \|\psi\|_*. \end{aligned}$$

Then thanks to the cut-off η_R , we have

$$|g_{2}| \lesssim (1 - \eta_{R}) \frac{1}{1 + |y|^{4}} |\psi|$$

$$\lesssim \frac{\mu_{0}^{\nu - \frac{1}{2}}(t) R^{-a}(t)}{1 + |y|^{4}} ||\psi||_{*} \chi_{\{|x| \geq \mu_{0} R\}}$$

$$\lesssim \frac{\mu_{0}^{\nu - \frac{1}{2} + a_{2}}(t) R^{a_{2} - a - 4}(t)}{|x|^{a_{2}}} ||\psi||_{*} \chi_{\{|x| \geq \mu_{0} R\}}.$$

So we have that for some $\epsilon > 0$

$$||g_2||_{**} \lesssim T^{\epsilon} ||\psi||_* \tag{6.4}$$

provided

$$\nu - \nu_2 - \frac{1}{2} + a_2 - \beta(a_2 - a - 4) > 0. \tag{6.5}$$

Estimate of g_3

We evaluate

$$\begin{split} |g_3| &\lesssim |(U_1 + \Phi_1)^3 \Phi_2^2| \\ &\lesssim \frac{\mu_0^{-\frac{3}{2}}(t)}{1 + |y|^3} \left(|\psi|^2 \chi_{\{|x| \lesssim \sqrt{T - t}\}} + \mu_0^{-1} |\phi|^2 \chi_{\{|x| \lesssim \mu_0 R\}} \right) \\ &\lesssim \frac{\mu_0^{-\frac{3}{2}}(t)}{1 + |y|^3} \left(\mu_0^{2\nu - 1}(t) R^{-2a}(t) \|\psi\|_*^2 \chi_{\{|x| \lesssim \sqrt{T - t}\}} + \frac{\mu_0^{2\nu - 1}(t) R^{\frac{8 - 2\sigma}{3}}(t)}{1 + |y|^2} \|\phi\|_{0, \nu, \sigma}^2 \chi_{\{|x| \lesssim \mu_0 R\}} \right) \\ &\lesssim \mu_0^{2\nu - \frac{5}{2}}(t) R^{-2a}(t) \|\psi\|_*^2 \chi_{\{|x| \lesssim \mu_0 R\}} + \mu_0^{2\nu - \frac{5}{2}}(t) R^{\frac{8 - 2\sigma}{3}}(t) \|\phi\|_{0, \nu, \sigma}^2 \chi_{\{|x| \lesssim \mu_0 R\}} \\ &+ \mu_0^{2\nu - \frac{5}{2} + a_2}(t) R^{a_2 - 2a - 3}(t) \frac{1}{|x|^{a_2}} \|\psi\|_*^2 \chi_{\{\mu_0 R \lesssim |x| \lesssim \sqrt{T - t}\}}. \end{split}$$

Then for some $\epsilon > 0$

$$||g_3||_{**} \lesssim T^{\epsilon} (1 + ||\psi||_*^2 + ||\phi||_{0,\nu,\sigma}^2)$$
 (6.6)

provided

$$\begin{cases}
\nu - \beta(2 - a) > 0 \\
\nu - \beta(\frac{14 - 2\sigma}{3} + a) > 0 \\
2\nu - \frac{5}{2} + a_2 - \beta(a_2 - 2a - 3) - \nu_2 > 0
\end{cases}$$
(6.7)

Estimate of g_4

To estimate g_4 , we first estimate

$$\left| \left[S(U_{1}) - \chi(\frac{x}{c_{0}(T-t)^{\frac{1}{2}}}) \frac{\alpha(t)}{(\mu^{2} + |x|^{2})^{\frac{1}{2}}} \right] (1 - \eta_{R}) \right| \\
\lesssim \left| \left[\eta_{1} S_{in} - \chi(\frac{x}{c_{0}(T-t)^{\frac{1}{2}}}) \frac{\alpha(t)}{(\mu^{2} + |x|^{2})^{\frac{1}{2}}} \right] (1 - \eta_{R}) \right| \\
+ \left| (1 - \eta_{o1}) \eta_{o2} (-\partial_{t} u_{out} + \Delta_{x} u_{out} + u_{out}^{5}) \right| \\
+ \left| -\partial_{t} \eta_{1} u_{in} + \Delta_{x} \eta_{1} u_{in} + 2 \nabla_{x} \eta_{1} \nabla_{x} u_{in} - \partial_{t} [(1 - \eta_{o1}) \eta_{o2}] u_{out} \right.$$

$$\left. + \Delta_{x} [(1 - \eta_{o1}) \eta_{o2}] u_{out} + 2 \nabla_{x} [(1 - \eta_{o1}) \eta_{o2}] \nabla_{x} u_{out} \right| \\
+ \left| \left[\eta_{1} u_{in} + (1 - \eta_{o1}) \eta_{o2} u_{out} \right]^{5} - \eta_{1} u_{in}^{5} - (1 - \eta_{o1}) \eta_{o2} u_{out}^{5} \right| .$$
(6.8)

From (2.13), we have

$$\left| \left[\eta_{1} S_{in} - \chi \left(\frac{x}{c_{0} (T - t)^{\frac{1}{2}}} \right) \frac{\alpha(t)}{(\mu^{2} + |x|^{2})^{\frac{1}{2}}} \right] (1 - \eta_{R}) \right| \\
\lesssim |\alpha(t)| \frac{\mu^{2}}{(\mu^{2} + |x|^{2})^{\frac{3}{2}}} \eta_{1} (1 - \eta_{R}) + \frac{|\alpha(t)|}{(\mu^{2} + |x|^{2})^{\frac{1}{2}}} \chi(r \leq \frac{|x|}{\sqrt{T - t}} \leq c_{0}) (1 - \eta_{R}) \\
+ \Lambda^{2} \mu^{-\frac{1}{2}} \mu'_{0} \left| 3^{\frac{1}{4}} \frac{\mu^{2}}{(\mu^{2} + |x|^{2})^{\frac{3}{2}}} - \frac{3^{\frac{1}{4}}}{2} \frac{1}{(\mu^{2} + |x|^{2})^{\frac{1}{2}}} \right| \eta_{1} (1 - \eta_{R}) \\
+ \left| 2\mu''_{0} \mu^{\frac{1}{2}} h(\frac{x}{\mu}) \eta_{1} + \mu'_{0} \mu^{-\frac{1}{2}} \mu' h(\frac{x}{\mu}) \eta_{1} + 2\mu'_{0} \mu^{-\frac{1}{2}} \mu' h(\frac{x}{\mu}) \eta_{1} \right| (1 - \eta_{R}) \\
+ \frac{\mu^{\frac{5}{2}}}{(\mu^{2} + |x|^{2})^{\frac{3}{2}}} (\mu'_{0})^{2} \left| h^{2} \left(\frac{x}{\mu_{0}} \right) \right| \eta_{1} (1 - \eta_{R}) \\
\lesssim \frac{\mu_{0}^{-\frac{3}{2} + a_{2}} \mu'_{0} R^{a_{2} - 3}}{|x|^{a_{2}}} \chi_{\{|x| \geq \mu_{0}R\}} \|\Lambda\|_{\infty} + \frac{\mu_{0}^{-\frac{1}{2}} \mu'_{0}}{|x|^{a_{2}}} \chi_{\{|x| \geq \mu_{0}R\}} \|\Lambda\|_{\infty} \\
+ \frac{\mu_{0}^{\frac{1}{2} + \frac{a_{2} - 3}{4k}}}{|x|^{a_{2}}} \chi_{\{|x| \geq \mu_{0}R\}} + \frac{\mu_{0}^{\frac{5}{2} + \frac{a_{2} - 5}{4k}}}{|x|^{a_{2}}} \chi_{\{|x| \geq \mu_{0}R\}}$$

where the function h is defined in (2.12). Similarly, we have the following estimates for the rest terms. We evaluate the term

$$|S_{out}| = |-\partial_t u_{out} + \Delta u_{out} + u_{out}^5| \lesssim (T-t)^{5(k-\zeta_1)}.$$
(6.10)

If we choose $\zeta_1 = \zeta_2 = \frac{1}{2}$, $r = r_1$ and $r_2 > 3r$, then we have

$$\left| -\partial_{t}\eta_{1}u_{in} + \Delta_{x}\eta_{1}u_{in} + 2\nabla_{x}\eta_{1}\nabla_{x}u_{in} - \partial_{t}[(1 - \eta_{o1})\eta_{o2}]u_{out} \right|$$

$$+ \Delta_{x}[(1 - \eta_{o1})\eta_{o2}]u_{out} + 2\nabla_{x}[(1 - \eta_{o1})\eta_{o2}]\nabla_{x}u_{out}$$

$$\lesssim |\partial_{t}\eta_{1}(u_{in} - u_{out})| + |\nabla_{x}\eta_{1} \cdot \nabla_{x}(u_{in} - u_{out})| + |\Delta_{x}\eta_{1}(u_{in} - u_{out})|$$

$$\lesssim (T - t)^{k - \frac{1}{2}},$$

$$(6.11)$$

where we have used the cancellation in the matching (2.8)–(2.9). Since $\zeta_1 = \zeta_2 = \frac{1}{2}$, we have

$$\begin{aligned} & |[\eta_1 u_{in} + (1 - \eta_{o1})\eta_{o2} u_{out}]^5 - \eta_1 u_{in}^5 - (1 - \eta_{o1})\eta_{o2} u_{out}^5| \\ & \lesssim |u_{out}|^5 \chi_{\{|x| \sim \sqrt{T - t}\}} \lesssim (T - t)^{5(k - \frac{1}{2})}. \end{aligned}$$

$$(6.12)$$

We choose initial condition of Φ_1 such that $\Phi_1(0,T)=0$. Then by Duhamel's formula, we can show that

$$|\Phi_1(x,t)| \lesssim \alpha(t)(T-t)^{\frac{3}{2}}.$$

Therefore, we obtain

$$|[(U_1 + \Phi_1)^5 - U_1^5](1 - \eta_R)| \lesssim (1 - \eta_R)|U_1^4 \Phi_1|$$

$$\lesssim (1 - \eta_R) \frac{\mu_0^{-2}(t)}{1 + |y|^4} \alpha(t) (T - t)^{\frac{3}{2}}.$$
(6.13)

Collecting estimates (6.8)–(6.13), we conclude that

$$||g_4||_{**} \lesssim T^{\epsilon} \left(1 + ||\Lambda||_{\infty}\right) \tag{6.14}$$

provided

$$\begin{cases}
a_{2} - \frac{1}{2} - \frac{1}{2k} + \beta(3 - a_{2}) - \nu_{2} > 0 \\
\frac{1}{2} - \frac{1}{2k} - \nu_{2} > 0 \\
\frac{1}{2} + \frac{a_{2} - 3}{4k} - \nu_{2} > 0 \\
5(\frac{1}{2} - \frac{1}{4k}) - \nu + \frac{5}{2} - \beta(2 + a) > 0 \\
2 - \nu - \frac{1}{4k} - \beta(2 + a) > 0 \\
a_{2} + \frac{1}{4k} - \frac{3}{2} + \beta(4 - a_{2}) - \nu_{2} > 0
\end{cases}$$
(6.15)

In conclusion, from (6.2), (6.3), (6.4), (6.5), (6.6), (6.7), (6.14) and (6.15), we obtain that for some $\epsilon > 0$

$$\|\mathcal{G}\|_{**} \lesssim T^{\epsilon} \left(1 + \|\psi\|_{*} + \|\phi\|_{0,\nu,\sigma} + \|\Lambda\|_{\infty}\right) \tag{6.16}$$

provided

$$\begin{cases}
1 + a - \sigma < 0 \\
\nu - \nu_2 - \frac{1}{2} + a_2 - \beta(a_2 - a - 4) > 0 \\
\nu - \beta(2 - a) > 0 \\
\nu - \beta(\frac{14 - 2\sigma}{3} + a) > 0 \\
2\nu - \frac{5}{2} + a_2 - \beta(a_2 - 2a - 3) - \nu_2 > 0 \\
a_2 - \frac{1}{2} - \frac{1}{2k} + \beta(3 - a_2) - \nu_2 > 0 \\
\frac{1}{2} - \frac{1}{2k} - \nu_2 > 0 \\
\frac{1}{2} + \frac{a_2 - 3}{4k} - \nu_2 > 0 \\
5(\frac{1}{2} - \frac{1}{4k}) - \nu + \frac{5}{2} - \beta(2 + a) > 0 \\
2 - \nu - \frac{1}{4k} - \beta(2 + a) > 0 \\
a_2 + \frac{1}{4k} - \frac{3}{2} + \beta(4 - a_2) - \nu_2 > 0
\end{cases} (6.17)$$

6.2. The inner problem: estimate of \mathcal{H} . Recall from (3.6) that

$$\mathcal{H}(\phi, \psi, \alpha) = 5 \left[(u_{in}(\mu_0 y, t) + \Phi_1(\mu_0 y, t))^4 - \mu_0^{-2} w^4(y) \right] \mu_0^2 \phi(y, t)$$

$$+ 5\mu_0^{\frac{1}{2}} (1 + \Lambda)^{-4} w^4 \left(\frac{y}{(1 + \Lambda)^2} \right) \psi(\mu_0 y, t) + \mu_0 \mu_0' \left[\frac{1}{2} \phi(y, t) + \nabla_y \phi(y, t) \cdot y \right]$$

$$+ \mu_0^{\frac{5}{2}} \left[S(U_1) - \chi \left(\frac{\mu_0 y}{c_0 (T - t)^{\frac{1}{2}}} \right) \frac{\alpha(t)}{(\mu^2 + |\mu_0 y|^2)^{\frac{1}{2}}} \right]$$

$$+ \mu_0^{\frac{5}{2}} \left[(u_{in}(\mu_0 y, t) + \Phi_1(\mu_0 y, t))^5 - u_{in}^5(\mu_0 y, t) \right].$$

We evaluate

$$\left| 5 \left[(u_{in}(\mu_0 y, t) + \Phi_1(\mu_0 y, t))^4 - \mu_0^{-2} w^4(y) \right] \mu_0^2 \phi(y, t) \right|
\lesssim \frac{\mu_0' \mu_0^{\frac{3}{4k}}}{1 + |y|^3} \frac{\mu_0' R^{\frac{4-\sigma}{3}}}{1 + |y|} \|\phi\|_{0, \nu, \sigma}, \tag{6.18}$$

$$\left| 5\mu_0^{\frac{1}{2}} (1+\Lambda)^{-4} w^4 \left(\frac{y}{(1+\Lambda)^2} \right) \psi(\mu_0 y, t) + \mu_0 \mu_0' \left[\frac{1}{2} \phi(y, t) + \nabla_y \phi(y, t) \cdot y \right] \right| \\
\lesssim \frac{\mu_0' R^{-a}}{1 + |y|^4} \|\psi\|_* + \frac{\mu_0 \mu_0' \mu_0' R^{\frac{4-\sigma}{3}}}{1 + |y|} \|\phi\|_{0, \nu, \sigma}. \tag{6.19}$$

By (2.13), we have

$$\left| \mu_0^{\frac{5}{2}} \left[S(U_1) - \chi(\frac{\mu_0 y}{c_0 (T - t)^{\frac{1}{2}}}) \frac{\alpha(t)}{(\mu^2 + |\mu_0 y|^2)^{\frac{1}{2}}} \right] \right|$$

$$\lesssim \frac{\mu_0 \mu_0' \|\Lambda\|_{\infty}}{1 + |y|^3} + \mu_0^{4 - \frac{1}{k}} (1 + |y|) + \frac{(\mu_0')^2 \mu_0^2}{1 + |y|}$$
(6.20)

and

$$\left| \mu_0^{\frac{5}{2}} \left[(u_{in}(\mu_0 y, t) + \Phi_1(\mu_0 y, t))^5 - u_{in}^5(\mu_0 y, t) \right] \right| \lesssim \frac{\mu_0' \mu_0^{\frac{3}{4k}}}{1 + |y|^4} \|\Lambda\|_{\infty}.$$
 (6.21)

From estimates (6.18)–(6.21), we obtain that for some $\epsilon > 0$

$$\|\mathcal{H}\|_{\nu,2+\sigma} \lesssim T^{\epsilon} \left(1 + \|\phi\|_{0,\nu,\sigma} + \|\psi\|_* + \|\Lambda\|_{\infty}\right) \tag{6.22}$$

provided

$$\begin{cases}
1 + \frac{1}{4k} - \frac{\beta(4-\sigma)}{3} > 0 \\
0 < \sigma < 2 \\
a > 0 \\
2 - \frac{1}{2k} - \frac{\beta(2\sigma+7)}{3} > 0 \\
2 - \frac{1}{2k} - \beta(\sigma-1) > 0 \\
4 - \frac{1}{k} - \beta(3+\sigma) > 0 \\
1 + \frac{1}{4k} - \nu > 0
\end{cases}$$
(6.23)

6.3. The fixed point formulation. The inner-outer gluing system (3.3) and (3.5) can be formulated as a fixed point problem for operators we shall describe below.

We first define the following function spaces

$$\mathcal{X}_{\phi} := \left\{ \phi \in L^{\infty}(B_{2R} \times (0,T)) \cap C_{y,t}^{2k+2\rho,k+\rho}(B_{2R} \times (0,T-\delta)) : \|\phi\|_{0,\nu,\sigma} < +\infty \right\}
\mathcal{X}_{\psi} := \left\{ \psi \in L^{\infty}(\mathbb{R}^{3} \times (0,T)) \cap C_{x,t}^{2k+2\rho,k+\rho}(\mathbb{R}^{3} \times (0,T-\delta)) : \|\psi\|_{*} < +\infty \right\}
\mathcal{X}_{\Lambda} := \left\{ \Lambda(t) : \|\Lambda(t)(T-t)^{-1+\varepsilon}\|_{L^{\infty}} \le C_{0\Lambda}, \|\Lambda^{(1)}(t)(T-t)^{\varepsilon}\|_{L^{\infty}} \le C_{1\Lambda}, \cdots, \right.
\|\Lambda^{(k)}(t)(T-t)^{k-1+\varepsilon}\|_{L^{\infty}} \le C_{k\Lambda}, \Lambda(t) \in C^{k+1,\rho}(0,T-\delta), \text{ for all } \delta \in (0,T) \right\}
\mathcal{X}_{\vec{c}} := \left\{ \vec{c} = (c_{1}, c_{2}, \dots, c_{k}) : |c_{j}| \le C_{c} T^{\frac{1}{2} - j - \varepsilon}, \ j = 1, 2, \dots, k \right\}$$

Define

$$\mathcal{X} = \mathcal{X}_{\phi} \times \mathcal{X}_{\psi} \times \mathcal{X}_{\Lambda} \times \mathcal{X}_{\vec{c}}. \tag{6.25}$$

We shall solve the inner-outer gluing system in a closed ball \mathcal{B} in $(\phi, \psi, \Lambda, \vec{c}) \in \mathcal{X}$.

The inner–outer gluing system (3.3) and (3.5) can be formulated as a fixed point problem, where we define an operator \mathcal{F} which returns the solution from \mathcal{B} to \mathcal{X}

$$\mathcal{F}: \mathcal{B} \subset \mathcal{X} \to \mathcal{X}$$

$$v \mapsto \mathcal{F}(v) = (\mathcal{F}_{\phi}(v), \mathcal{F}_{\psi}(v), \mathcal{F}_{\Lambda}(v), \mathcal{F}_{\vec{c}}(v))$$

with

$$\mathcal{F}_{\phi}(\phi, \psi, \Lambda, \vec{c}) = \mathcal{T}_{\phi}(\mathcal{H}[\phi, \psi, \Lambda])$$

$$\mathcal{F}_{\psi}(\phi, \psi, \Lambda, \vec{c}) = \mathcal{T}_{\psi}(\mathcal{G}(\phi, \psi, \Lambda))$$

$$\mathcal{F}_{\Lambda}(\phi, \psi, \Lambda, \vec{c}) = \mathcal{T}_{\Lambda}(\phi, \psi, \Lambda, \vec{c})$$

$$\mathcal{F}_{\vec{c}}(\phi, \psi, \Lambda, \vec{c}) = \mathcal{T}_{\vec{c}}(\phi, \psi, \Lambda, \vec{c})$$
(6.26)

Here \mathcal{T}_{ϕ} is the operators given from Proposition 4.1 which solves the inner problem (3.5). The operator \mathcal{T}_{ψ} defined by Proposition 4.2 deals with the outer problem (3.3). Operators \mathcal{T}_{Λ} and $\mathcal{T}_{\vec{c}}$ handle the reduced equation (5.1).

6.4. Choice of constants. In this section, we list all the constraints of the parameters which are sufficient for the inner–outer gluing scheme to work.

We first indicate all the parameters used in different norms.

- $R(t) = \mu_0^{-\beta}(t)$ with $\beta \in (0, 1/2)$.
- The norm for ϕ solving the inner problem (3.5) is $\|\cdot\|_{0,\nu,\sigma}$ which is defined in (4.2), where we require that $\nu > 0$, $0 < \sigma < 2$.
- The norm for ψ solving the outer problem (3.3) is $\|\cdot\|_*$ which is defined in (4.6), while the $\|\cdot\|_{**}$ norm for the right hand side of the outer problem (3.3) is defined in (4.5). Here we require that $\nu, \nu_2 > 0$ and $a, \gamma \in (0, 1)$. Also, as mentioned in Remark 4.2, we require $\nu_2 + \frac{2-a_2}{4k} > \nu \frac{1}{2} + a\beta$ such that the $\|\cdot\|_*$ -norm is well defined.

In order to get the desired estimates for the outer problem (3.3), by (6.17), we need the following restrictions

$$\begin{cases} 1+a-\sigma<0\\ \nu-\nu_2-\frac{1}{2}+a_2-\beta(a_2-a-4)>0\\ \nu-\beta(2-a)>0\\ \nu-\beta(\frac{14-2\sigma}{3}+a)>0\\ 2\nu-\frac{5}{2}+a_2-\beta(a_2-2a-3)-\nu_2>0\\ a_2-\frac{1}{2}-\frac{1}{2k}+\beta(3-a_2)-\nu_2>0\\ \frac{1}{2}-\frac{1}{2k}-\nu_2>0\\ \frac{1}{2}+\frac{a_2-3}{4k}-\nu_2>0\\ 5(\frac{1}{2}-\frac{1}{4k})-\nu+\frac{5}{2}-\beta(2+a)>0\\ 2-\nu-\frac{1}{4k}-\beta(2+a)>0\\ a_2+\frac{1}{4k}-\frac{3}{2}+\beta(4-a_2)-\nu_2>0 \end{cases}$$

In order to get the desired estimates for the inner problem (3.5), by (6.23), we need

$$\begin{cases} 1 + \frac{1}{4k} - \frac{\beta(4-\sigma)}{3} > 0 \\ 0 < \sigma < 2 \\ a > 0 \\ 2 - \frac{1}{2k} - \frac{\beta(2\sigma+7)}{3} > 0 \\ 2 - \frac{1}{2k} - \beta(\sigma - 1) > 0 \\ 4 - \frac{1}{k} - \beta(3+\sigma) > 0 \\ 1 + \frac{1}{4\ell} - \nu > 0 \end{cases}$$

Elementary computations show that suitable choices of the parameters satisfying all the restrictions in this section can be found, which ensures the implementation of the gluing procedure.

6.5. **Proof of Theorem 1.** Consider the operator

$$\mathcal{F} = (\mathcal{F}_{\phi}, \mathcal{F}_{\psi}, \mathcal{F}_{\Lambda}, \mathcal{F}_{\vec{c}}) \tag{6.27}$$

given in (6.26). To prove Theorem 1, our strategy is to show the existence of a fixed point for the operator \mathcal{F} in \mathcal{B} by the Schauder fixed point theorem. By collecting the estimates (6.16), (6.22), and using Proposition 4.2, Proposition 4.1 and discussions in Section 5, we conclude that for $(\phi, \psi, \Lambda, \vec{c}) \in \mathcal{B}$

$$\begin{cases}
\|\mathcal{F}_{\phi}(\phi, \psi, \Lambda, \vec{c})\|_{0,\nu,\sigma} \leq CT^{\epsilon} \\
\|\mathcal{F}_{\psi}(\phi, \psi, \Lambda, \vec{c})\|_{*} \leq CT^{\epsilon} \\
\|\mathcal{F}_{\Lambda}(\phi, \psi, \Lambda, \vec{c})\|_{\Lambda} \leq CT^{\epsilon} \\
\|\mathcal{F}_{\vec{c}}(\phi, \psi, \Lambda, \vec{c})\|_{\vec{c}} \leq CT^{\epsilon}
\end{cases}$$
(6.28)

where C>0 is a constant independent of T, and $\epsilon>0$ is a small fixed number. On the other hand, compactness of the operator \mathcal{F} defined in (6.27) can be proved by proper variants of (6.28). Indeed, if we vary the parameters slightly such that all the restrictions in Section 6.4 are still satisfied, then we get (6.28) with the norms in the left hand side defined by the new parameters, while the closed ball \mathcal{B} remains the same. To be more specific, for fixed ν', a' which are close to ν, a , one can show that if $(\phi, \psi, \Lambda) \in \mathcal{B}$, then

$$\|\mathcal{F}_{\phi}(\phi, \psi, \Lambda, \vec{c})\|_{0, \nu', \sigma'} \leq CT^{\epsilon'}.$$

Furthermore, one can show that for $\nu' > \nu$ and $\nu' - \frac{\beta(4-\sigma')}{3} > \nu - \frac{\beta(4-\sigma)}{3}$, one has a compact embedding in the sense that if a sequence $\{\phi_n^0\}$ is bounded in the $\|\cdot\|_{0,\nu',\sigma'}$ -norm, then there exists a subsequence which converges in the $\|\cdot\|_{0,\nu,\sigma}$ -norm. Thus, the compactness follows directly from a standard diagonal argument by Arzelà–Ascoli's theorem. Arguing in a similar manner, the compactness for the rest operators can be proved. Therefore, the existence of the desired blow-up solution is concluded from the Schauder fixed point theorem.

7. Nonradial case: blow-up at multiple points

As a by-product, the inner-outer gluing method carried out in this paper can be applied to construct non-radial type II blow-up at multiple N points for the first blow-up rate k = 1. To be more precise, we take the first approximation to be

$$U_N = \sum_{j=1}^{N} \mu_j^{-\frac{1}{2}}(t) w \left(\frac{x - \xi_j(t)}{\mu_j(t)} \right),$$

where we expect that the scaling and translation parameters satisfy

$$\mu_i(t) \to 0$$
, $\xi_i(t) \to q_i$ as $t \to T$

for $j=1,\ldots,N$, where q_j are given points in \mathbb{R}^3 with $\max_{k,l=1,\ldots,N}|q_k-q_l|>\delta$ for uniform $\delta>0$. Formally, the error of U_N behaves like

$$S(U_N) \sim \sum_{j=1}^{N} \left(\mu_j^{-\frac{3}{2}}(t) \dot{\mu}_j(t) Z_0(y_j) + \mu_j^{-\frac{3}{2}}(t) \dot{\xi}_j \cdot \nabla w(y_j) \right)$$

$$:= \sum_{j=1}^{N} \left(\mathcal{E}_{0,j} + \mathcal{E}_{1,j} \right)$$

where $y_j = \frac{x - \xi_j(t)}{\mu_j(t)}$. To cancel out the slow decay error at mode 0 near each point q_j , we introduce the correction $\Phi_{(j)}$ solving

$$\partial_t \Phi_{(j)} = \Delta \Phi_{(j)} + \mathcal{E}_{0,j} \quad \text{for } j = 1 \dots, N$$

so that the corrected approximation is

$$u_* = \sum_{j=1}^{N} \mu_j^{-\frac{1}{2}}(t) w \left(\frac{x - \xi_j(t)}{\mu_j(t)} \right) + \Phi_{(j)}.$$

We then look for the solution

$$u = u_* + \sum_{j=1}^{N} \mu_j^{-\frac{1}{2}}(t) \eta_{R_j} \phi(y_j, t) + \psi(x, t).$$

Let us emphasize that in the non-radial case, the blow-up rate for k=1 will be obtained by <u>orthogonality condition</u> instead of the matching in the general case $k \geq 2$. The orthogonality condition at scaling mode 0 is basically

$$\int_{B_{2R}} 5w^4 \Phi_{(j)} Z_0(y) dy + \int_{B_{2R}} 5w^4 \psi Z_0(y) dy \approx 0$$

which turns out to be a nonlocal equation like before, and using the method in Section 5, we have

$$\mu_j(t) \sim (T-t)^2 \text{ for } j=1,\cdots,N.$$

Indeed, the orthogonality condition at mode 0 gives a nonlocal reduced equation of the following form:

$$\int_0^t \frac{\mu_j^{-1/2}(s)\dot{\mu}_j(s)}{(t-s)^{1/2}} ds = c_{*,j},$$

where $c_{*,j} < 0$ is some constant coming from the initial data. We rewrite the above integro-differential equation as

$$\int_0^t \frac{\dot{v}_j(s)}{(t-s)^{1/2}} ds = c_{*,j},$$

where $v_j(t) = 2\mu_j^{1/2}(t)$. Imposing $v_j(T) = 0$ and using (5.11), we obtain that for some c > 0

$$v_i(t) - v_i(T) = v_i(t) = c(T^{1/2} - t^{1/2}) \sim T - t$$

and thus

$$\mu_j(t) \sim (T-t)^2$$

which is precisely the first rate (k = 1) predicted in [14].

On the other hand, the orthogonality condition at translation mode 1

$$\int_{B_{2R}} \mathcal{E}_{1,j} Z_{\ell}(y) dy \approx 0$$

simply implies

$$\xi_j(t) \sim q_j$$
 for $j = 1, \dots, N$,

where

$$Z_{\ell}(y) = \partial_{y_{\ell}} w \quad \ell = 1, 2, 3.$$

We will not elaborate on the details.

APPENDIX A. PROOFS OF TECHNICAL LEMMAS

In this appendix, we prove the technical lemmas in Section 4.2.

Proof of lemma 4.1. Duhamel's formula gives

$$|\psi| \lesssim \int_0^t \int_{|w| \le 2\mu_0(s)R(s)} \frac{e^{-\frac{|x-w|^2}{4(t-s)}}}{(4\pi(t-s))^{3/2}} \mu_0^{\nu-\frac{5}{2}}(s) R^{-2-a}(s) dw ds$$

We decompose

$$\int_{0}^{t} \mu_{0}^{\nu - \frac{5}{2}}(s) R^{-2-a}(s) \int_{|w| \le 2\mu_{0}(s)R(s)} \frac{e^{-\frac{|x-w|^{2}}{4(t-s)}}}{(4\pi(t-s))^{3/2}} dw ds
= \left(\int_{0}^{t-(T-t)} + \int_{t-(T-t)}^{t-\mu_{0}^{\delta_{1}}(t)} + \int_{t-\mu_{0}^{\delta_{1}}(t)}^{t} \right) \mu_{0}^{\nu - \frac{5}{2}}(s) R^{-2-a}(s) \int_{|w| \le 2\mu_{0}(s)R(s)} \frac{e^{-\frac{|x-w|^{2}}{4(t-s)}}}{(4\pi(t-s))^{3/2}} dw ds
:= I_{11} + I_{12} + I_{13}$$

for some $\delta_1 \geq 1$ to be found. Directly integrating, we obtain

$$I_{11} = \int_{0}^{t-(T-t)} \mu_{0}^{\nu-\frac{5}{2}}(s) R^{-2-a}(s) \int_{|w| \leq 2\mu_{0}(s)R(s)} \frac{e^{-\frac{|x-w|^{2}}{4(t-s)}}}{(4\pi(t-s))^{3/2}} dwds$$

$$\lesssim \int_{0}^{t-(T-t)} \mu_{0}^{\nu-\frac{5}{2}}(s) R^{-2-a}(s) \int_{|\tilde{w}| \leq \frac{2\mu_{0}(s)R(s)}{\sqrt{t-s}}} e^{-\frac{|\tilde{x}-\tilde{w}|^{2}}{4}} d\tilde{w}ds$$

$$\lesssim \int_{0}^{t-(T-t)} \mu_{0}^{\nu-\frac{5}{2}}(s) R^{-2-a}(s) \frac{(\mu_{0}(s)R(s))^{3}}{(t-s)^{3/2}} ds$$

$$\lesssim \int_{0}^{t-(T-t)} \frac{\mu_{0}^{\nu+\frac{1}{2}}(s) R^{1-a}(s)}{(T-s)^{3/2}} ds$$

$$\lesssim \mu_{0}^{\nu+\frac{1}{2}-\frac{1}{4k}}(0) R^{1-a}(0),$$

Similarly we compute

$$I_{12} \lesssim \int_{t-(T-t)}^{t-\mu_0^{\delta_1}(t)} \mu_0^{\nu-\frac{5}{2}}(s) R^{-2-a}(s) \frac{(\mu_0(s)R(s))^3}{(t-s)^{3/2}} ds$$

$$\lesssim \mu_0^{\nu+\frac{1-\delta_1}{2}}(t) R^{1-a}(t)$$
(A.1)

and

$$I_{13} \lesssim \int_{t-\mu_0^{\delta_1}(t)}^{t} \mu_0^{\nu - \frac{5}{2}}(s) R^{-2-a}(s) ds$$

$$\lesssim \mu_0^{\nu - \frac{5}{2} + \delta_1}(t) R^{-2-a}(t). \tag{A.2}$$

Since $\beta \in (0, 1/2)$, we can choose $\delta_1 = 2 - 2\beta$. Therefore, we get

$$I_{11} + I_{12} + I_{13} \lesssim \mu_0^{\nu - \frac{1}{2}}(0)R^{-a}(0)$$

as desired.

Similarly, to prove (4.8), we decompose

$$|\psi(x,t) - \psi(x,T)| \le I_{21} + I_{22} + I_{23}$$

with

$$I_{21} = \int_{0}^{t-(T-t)} \int_{\mathbb{R}^{3}} |G(x-w,t-s) - G(x-w,T-s)| |f(w,s)| dwds$$

$$I_{22} = \int_{t-(T-t)}^{t} \int_{\mathbb{R}^{3}} |G(x-w,t-s) - G(x-w,T-s)| |f(w,s)| dwds$$

$$I_{23} = \int_{t}^{T} \int_{\mathbb{R}^{3}} |G(x-w,T-s)| |f(w,s)| dwds,$$

where G(x,t) is the heat kernel

$$G(x,t) = \frac{e^{-\frac{|x|^2}{4t}}}{(4\pi t)^{3/2}}.$$
(A.3)

For the first integral I_{21} , we have

$$I_{21} \leq (T-t) \int_0^1 \int_0^{t-(T-t)} \int_{|w| \leq 2\mu_0(s)R(s)} |\partial_t G(x-w,t_v-s)| \mu_0^{\nu-\frac{5}{2}}(s) R^{-2-a}(s) dw ds dv,$$

where $t_v = vT + (1 - v)t$. Changing variables, we evaluate

$$\int_{|w| \le 2\mu_0(s)R(s)} |\partial_t G(x - w, t_v - s)| dw$$

$$\lesssim \int_{|w| \le 2\mu_0(s)R(s)} e^{-\frac{|x - w|^2}{4(t_v - s)}} \left(\frac{|x - w|^2}{(t_v - s)^{\frac{7}{2}}} + \frac{1}{(t_v - s)^{\frac{5}{2}}} \right) dw$$

$$= \int_{|w_v| \le \frac{2\mu_0(s)R(s)}{\sqrt{t_v - s}}} e^{-\frac{|x_v - w_v|^2}{4}} \left(1 + |x_v - w_v|^2 \right) \frac{1}{t_v - s} dw_v$$

and thus

$$\begin{split} & \int_0^{t-(T-t)} \int_{|w| \leq 2\mu_0(s)R(s)} |\partial_t G(x-w,t_v-s)| \mu_0^{\nu-\frac{5}{2}}(s) R^{-2-a}(s) dw ds \\ & \lesssim \int_0^{t-(T-t)} \mu_0^{\nu-\frac{5}{2}}(s) R^{-2-a}(s) \frac{(\mu_0(s)R(s))^3}{(t_v-s)^{5/2}} ds \\ & \lesssim \int_0^{t-(T-t)} \mu_0^{\nu+\frac{1}{2}}(s) R^{1-a}(s) (T-s)^{-5/2} ds \\ & \lesssim \mu_0^{\nu+\frac{1}{2}-\frac{3}{4k}}(t) R^{1-a}(t), \end{split}$$

from which we conclude that

$$I_{21} \lesssim \mu_0^{\nu + \frac{1}{2} - \frac{1}{4k}}(t)R^{1-a}(t).$$
 (A.4)

For I_{22} , we have

$$I_{22} \leq \int_{t-(T-t)}^{t} \int_{|w| \leq 2\mu_0(s)R(s)} |G(x-w,t-s)| \mu_0^{\nu-\frac{5}{2}}(s) R^{-2-a}(s) dw ds$$
$$+ \int_{t-(T-t)}^{t} \int_{|w| \leq 2\mu_0(s)R(s)} |G(x-w,T-s)| \mu_0^{\nu-\frac{5}{2}}(s) R^{-2-a}(s) dw ds.$$

The first integral above can be estimated as

$$\begin{split} &\int_{t-(T-t)}^t \int_{|w| \le 2\mu_0(s)R(s)} |G(x-w,t-s)| \mu_0^{\nu-\frac{5}{2}}(s) R^{-2-a}(s) dw ds \\ &= \left(\int_{t-(T-t)}^{t-\mu_0^{\delta_1}(t)} + \int_{t-\mu_0^{\delta_1}(t)}^t \right) |G(x-w,t-s)| \mu_0^{\nu-\frac{5}{2}}(s) R^{-2-a}(s) dw ds. \end{split}$$

Notice that we already estimate the above integral in (A.1) and (A.2). So the choice $\delta_1 = 2 - 2\beta$, one has

$$\int_{t-(T-t)}^{t} \int_{|w| \le 2\mu_0(s)R(s)} |G(x-w,t-s)| \mu_0^{\nu-\frac{5}{2}}(s) R^{-2-a}(s) dw ds$$

$$\lesssim \mu_0^{\nu-\frac{1}{2}}(t) R^{-a}(t).$$

Similarly, it holds that

$$\int_{t-(T-t)}^{t} \int_{|(w_r,w_z)-\xi(s)| \le 2\mu_0(s)R(s)} |G(x-w,T-s)| \mu_0^{\nu-3}(s) R^{-2-\alpha}(s) dw ds$$

$$\lesssim \mu_0^{\nu-\frac{1}{2}}(t) R^{-a}(t).$$

Therefore, we obtain

$$I_{22} \lesssim \mu_0^{\nu - \frac{1}{2}}(t)R^{-a}(t).$$
 (A.5)

For I_{23} , changing variables, one has

$$I_{23} \lesssim \int_{t}^{T} \int_{|w| \leq 2\mu_{0}(s)R(s)} \frac{e^{-\frac{|x-w|^{2}}{4(T-s)}}}{(T-s)^{3/2}} \mu_{0}^{\nu-\frac{5}{2}}(s) R^{-2-a}(s) dw ds$$

$$\lesssim \int_{t}^{T} \int_{|\tilde{w}| \leq \frac{2\mu_{0}(s)R(s)}{\sqrt{T-s}}} e^{-\frac{|\tilde{x}-\tilde{w}|^{2}}{4}} \mu_{0}^{\nu-\frac{5}{2}}(s) R^{-2-a}(s) d\tilde{w} ds$$

$$\lesssim \int_{t}^{T} \mu_{0}^{\nu-\frac{5}{2}}(s) R^{-2-a}(s) \frac{(\mu_{0}(s)R(s))^{3}}{(T-s)^{3/2}} ds$$

$$\lesssim \mu_{0}^{\nu+\frac{1}{2}-\frac{1}{2k}}(t) R^{1-a}(t)$$
(A.6)

Collecting (A.4), (A.5) and (A.6), we conclude the validity of (4.8).

Then we prove the gradient estimate (4.9). By the heat kernel, we get

$$|\nabla \psi(x,t)| \lesssim \int_0^t \frac{\mu_0^{\nu - \frac{5}{2}}(s)R^{-2-a}(s)}{(t-s)^{\frac{5}{2}}} \int_{|w| \leq 2\mu_0(s)R(s)} e^{-\frac{|x-w|^2}{4(t-s)}} |x-w| dw ds$$

$$\lesssim \int_0^t \frac{\mu_0^{\nu - \frac{5}{2}}(s)R^{-2-a}(s)}{(t-s)^{1/2}} \int_{|\tilde{w}| \leq \frac{2\mu_0(s)R(s)}{\sqrt{t-s}}} e^{-\frac{|\tilde{w}|^2}{4}} (1+|\tilde{w}|) d\tilde{w} ds,$$

where $\tilde{x} = x(t-s)^{-1/2}$. First, we compute

$$\int_{0}^{t-(T-t)} \frac{\mu_{0}^{\nu-\frac{5}{2}}(s)R^{-2-a}(s)}{(t-s)^{1/2}} \int_{|w| \leq \frac{2\mu_{0}(s)R(s)}{\sqrt{t-s}}} e^{-\frac{|\tilde{w}|^{2}}{4}} (1+|\tilde{w}|) d\tilde{w} ds$$

$$\lesssim \int_{0}^{t-(T-t)} \frac{\mu_{0}^{\nu-\frac{5}{2}}(s)R^{-2-a}(s)}{(t-s)^{1/2}} \frac{(\mu_{0}(s)R(s))^{3}}{(t-s)^{3/2}} ds$$

$$\lesssim \int_{0}^{t-(T-t)} \frac{\mu_{0}^{\nu+\frac{1}{2}}(s)R^{1-a}(s)}{(t-s)^{2}} ds$$

$$\lesssim \int_{0}^{t-(T-t)} \mu_{0}^{\nu+\frac{1}{2}}(s)R^{1-a}(s)(T-s)^{-2} ds$$

$$\lesssim \mu_{0}^{\nu+\frac{1}{2}-\frac{1}{2k}}(0)R^{1-a}(0).$$
(A.7)

Then we compute

$$\int_{t-(T-t)}^{t-\mu_0^{\delta_2}(t)} \frac{\mu_0^{\nu-\frac{5}{2}}(s)R^{-2-a}(s)}{(t-s)^{1/2}} \int_{|w| \leq \frac{2\mu_0(s)R(s)}{\sqrt{t-s}}} e^{-\frac{|\tilde{w}|^2}{4}} (1+|\tilde{w}|) d\tilde{w} ds
\lesssim \int_{t-(T-t)}^{t-\mu_0^{\delta_2}(t)} \frac{\mu_0^{\nu+\frac{1}{2}}(s)R^{1-a}(s)}{(t-s)^2} ds
\lesssim \mu_0^{\nu+\frac{1}{2}-\delta_2}(t)R^{1-a}(t),$$
(A.8)

where $\delta_2 \geq 1$ is a constant to be determined. On the other hand, we have

$$\int_{t-\mu_0^{\delta_2}(t)}^{t} \frac{\mu_0^{\nu-\frac{5}{2}}(s)R^{-2-a}(s)}{(t-s)^{\frac{5}{2}}} \int_{|w| \le 2\mu_0(s)R(s)} e^{-\frac{|x-w|^2}{4(t-s)}} |x-w| dw ds$$

$$\lesssim \int_{t-\mu_0^{\delta_2}(t)}^{t} \frac{\mu_0^{\nu-\frac{5}{2}}(s)R^{-2-a}(s)}{(t-s)^{1/2}} ds$$

$$\lesssim \mu_0^{\nu-\frac{5}{2}+\frac{\delta_2}{2}}(t)R^{-2-a}(t).$$
(A.9)

By choosing $\delta_2 = 2 - 2\beta$ and combining (A.7)–(A.9), we prove the validity of the gradient estimate (4.9). The proof of (4.10) is similar to that of (4.8). We omit the details.

To prove the Hölder estimate (4.11), we decompose

$$|\psi(x,t_2) - \psi(x,t_1)| \le J_{11} + J_{12} + J_{13}$$

with

$$J_{11} = \int_0^{t_1 - (t_2 - t_1)} \int_{\mathbb{R}^3} |G(x - w, t_2 - s) - G(x - w, t_1 - s)| f(w, s) dw ds,$$

$$J_{12} = \int_{t_1 - (t_2 - t_1)}^{t_1} \int_{\mathbb{R}^3} |G(x - w, t_2 - s) - G(x - w, t_1 - s)| f(w, s) dw ds,$$

and

$$J_{13} = \int_{t_1}^{t_2} \int_{\mathbb{R}^3} G(x - w, t_2 - s) f(w, s) dw ds,$$

where G(x,t) is the heat kernel (A.3). Here we assume that $0 < t_1 < t_2 < T$ with $t_2 < 2t_1$. For J_{11} , by letting $t_v = vt_2 + (1-v)t_1$, we have

$$J_{11} \leq (t_2 - t_1) \int_0^1 \int_0^{t_1 - (t_2 - t_1)} \int_{\mathbb{R}^3} |\partial_t G(x - w, t_v - s)| f(w, s) dw ds dv$$

$$\lesssim (t_2 - t_1) \int_0^1 \int_0^{t_1 - (t_2 - t_1)} \int_{|w| \leq 2\mu_0(s)R(s)} e^{-\frac{|x - w|^2}{4(t_v - s)}} \left(\frac{|x - w|^2}{(t_v - s)^{\frac{7}{2}}}\right) ds ds dv$$

$$+ \frac{1}{(t_v - s)^{\frac{5}{2}}} \mu_0^{\nu - \frac{5}{2}} (s) R^{-2-a}(s) dw ds dv,$$

and

$$\int_{|w| \le 2\mu_0(s)R(s)} e^{-\frac{|x-w|^2}{4(t_v-s)}} \left(\frac{|x-w|^2}{(t_v-s)^{\frac{7}{2}}} + \frac{1}{(t_v-s)^{\frac{5}{2}}} \right) \mu_0^{\nu-\frac{5}{2}}(s) R^{-2-a}(s) dw$$

$$= \int_{|w_v| \le \frac{2\mu_0(s)R(s)}{\sqrt{t_v-s}}} e^{-\frac{|x_v-w_v|^2}{4}} \left(1 + |x_v-w_v|^2 \right) \frac{\mu_0^{\nu-\frac{5}{2}}(s) R^{-2-a}(s)}{t_v-s} dw_v.$$

Observing that for any $\gamma_1 \in (0,1)$, we have

$$\int_{|w_v| \le \frac{2\mu_0(s)R(s)}{\sqrt{t_v - s}}} e^{-\frac{|x_v - w_v|^2}{4}} \left(1 + |x_v - w_v|^2\right) dw_v \lesssim \left(\frac{2\mu_0(s)R(s)}{\sqrt{t_v - s}}\right)^{\gamma_1}.$$

Thus, one has

$$J_{11} \lesssim (t_2 - t_1) \int_0^{t_1 - (t_2 - t_1)} \frac{\mu_0^{\nu - \frac{5}{2} + \gamma_1}(s) R^{-2 - a + \gamma_1}(s)}{(t_2 - s)^{1 + \frac{\gamma_1}{2}}} ds.$$

Recalling that $R(t) = \mu_0^{-\beta}(t)$ for $\beta \in (0, 1/2)$, we have the following two cases

• If $\nu - \frac{5}{2} + \gamma_1 + \beta(2 + a - \gamma_1) < 0$, then we have

$$\int_{0}^{t_{1}-(t_{2}-t_{1})} \frac{\mu_{0}^{\nu-\frac{5}{2}+\gamma_{1}}(s)R^{-2-a+\gamma_{1}}(s)}{(t_{2}-s)^{1+\frac{\gamma_{1}}{2}}} ds$$

$$\lesssim \mu_{0}^{\nu-\frac{5}{2}+\gamma_{1}}(t_{1})R^{-2-a+\gamma_{1}}(t_{1}) \int_{0}^{t_{1}-(t_{2}-t_{1})} \frac{1}{(t_{2}-s)^{1+\frac{\gamma_{1}}{2}}} ds$$

$$\lesssim \mu_{0}^{\nu-\frac{5}{2}+\gamma_{1}}(t_{1})R^{-2-a+\gamma_{1}}(t_{1})(t_{2}-t_{1})^{-\gamma_{1}/2}.$$

• If $\nu - \frac{5}{2} + \gamma_1 + \beta(2 + a - \gamma_1) \ge 0$, then we decompose

$$\begin{split} & \int_0^{t_1-(t_2-t_1)} \frac{\mu_0^{\nu-\frac{5}{2}+\gamma_1}(s)R^{-2-a+\gamma_1}(s)}{(t_2-s)^{1+\frac{\gamma_1}{2}}} ds \\ & = \left(\int_0^{t_1-(T-t_1)} + \int_{t_1-(T-t_1)}^{t_1-(t_2-t_1)} \right) \frac{\mu_0^{\nu-\frac{5}{2}+\gamma_1}(s)R^{-2-a+\gamma_1}(s)}{(t_2-s)^{1+\frac{\gamma_1}{2}}} ds. \end{split}$$

Assuming

$$2k[\nu - \frac{5}{2} + \gamma_1 + \beta(2 + a - \gamma_1)] - \frac{\gamma_1}{2} < 0,$$

we obtain that

$$\int_{0}^{t_{1}-(T-t_{1})} \frac{\mu_{0}^{\nu-\frac{5}{2}+\gamma_{1}}(s)R^{-2-a+\gamma_{1}}(s)}{(t_{2}-s)^{1+\frac{\gamma_{1}}{2}}} ds$$

$$\lesssim \int_{0}^{t_{1}-(T-t_{1})} \frac{\mu_{0}^{\nu-\frac{5}{2}+\gamma_{1}}(s)R^{-2-a+\gamma_{1}}(s)}{(T-s)^{1+\frac{\gamma_{1}}{2}}} ds$$

$$= \int_{0}^{t_{1}-(T-t_{1})} (T-s)^{2k[\nu-\frac{5}{2}+\gamma_{1}+\beta(2+a-\gamma_{1})]-1-\frac{\gamma_{1}}{2}} ds$$

$$\lesssim \mu_{0}^{\nu-\frac{5}{2}+\gamma_{1}}(t_{2})R^{-2-a+\gamma_{1}}(t_{2})(t_{2}-t_{1})^{-\gamma_{1}/2}$$

and similarly

$$\int_{t_1-(T-t_1)}^{t_1-(t_2-t_1)} \frac{\mu_0^{\nu-\frac{5}{2}+\gamma_1}(s)R^{-2-a+\gamma_1}(s)}{(t_2-s)^{1+\frac{\gamma_1}{2}}} ds \lesssim \mu_0^{\nu-\frac{5}{2}+\gamma_1}(t_2)R^{-2-a+\gamma_1}(t_2)(t_2-t_1)^{-\gamma_1/2}.$$

In both cases, we have

$$J_{11} \lesssim \mu_0^{\nu - \frac{5}{2} + \gamma_1} (t_2) R^{-2 - a + \gamma_1} (t_2) (t_2 - t_1)^{1 - \gamma_1/2}.$$

For J_{12} , we evaluate

$$\int_{t_1 - (t_2 - t_1)}^{t_1} \int_{\mathbb{R}^3} |G(x - w, t_1 - s)| f(w, s) dw ds$$

$$\lesssim \int_{t_1 - (t_2 - t_1)}^{t_1} \mu_0^{\nu - \frac{5}{2}}(s) R^{-2 - a}(s) \int_{|\tilde{w}| \le \frac{2\mu_0(s)R(s)}{\sqrt{t_1 - s}}} e^{-\frac{|\tilde{x} - \tilde{w}|^2}{4}} d\tilde{w} ds$$

$$\lesssim \int_{t_1 - (t_2 - t_1)}^{t_1} \mu_0^{\nu - \frac{5}{2}}(s) R^{-2 - a}(s) \left(\frac{2\mu_0(s)R(s)}{\sqrt{t_1 - s}}\right)^{\gamma_1} ds$$

$$\lesssim \mu_0^{\nu - \frac{5}{2} + \gamma_1}(t_2) R^{-2 - a + \gamma_1}(t_2)(t_2 - t_1)^{1 - \gamma_1/2},$$

where $\gamma_1 \in (0,1)$. Similarly, we have

$$\int_{t_1-(t_2-t_1)}^{t_1} \int_{\mathbb{R}^3} |G(x-w,t_2-s)| f(w,s) dw ds \lesssim \mu_0^{\nu-\frac{5}{2}+\gamma_1}(t_2) R^{-2-a+\gamma_1}(t_2) (t_2-t_1)^{1-\gamma_1/2}.$$

Thus we conclude that

$$J_{12} \lesssim \mu_0^{\nu - \frac{5}{2} + \gamma_1} (t_2) R^{-2 - a + \gamma_1} (t_2) (t_2 - t_1)^{1 - \gamma_1/2}.$$

Finally, for J_{13} ,

$$J_{13} = \int_{t_1}^{t_2} \int_{\mathbb{R}^3} G(x - w, t_2 - s) f(w, s) dw ds$$

$$\lesssim \int_{t_1}^{t_2} \mu_0^{\nu - \frac{5}{2}}(s) R^{-2 - a}(s) \int_{\tilde{w} \leq \frac{2\mu_0(s)R(s)}{\sqrt{t_2 - s}}} e^{-\frac{|\tilde{x} - \tilde{w}|^2}{4}} d\tilde{w} ds$$

$$\lesssim \mu_0^{\nu - \frac{5}{2} + \gamma_1}(t_2) R^{-2 - a + \gamma_1}(t_2) (t_2 - t_1)^{1 - \gamma_1/2}$$

follows from the same argument as before. This completes the proof of (4.11).

Proof of Lemma 4.2. We first prove (4.12). Similar to the proof of Lemma 4.1, Duhamel's formula gives

$$|\psi(x,t)| \lesssim \int_0^t \frac{\mu_0^{\nu_2}(s)}{(t-s)^{3/2}} \int_{\mu_0(s)R(s) \le |w|} \frac{e^{-\frac{|x-w|^2}{4(t-s)}}}{|w|^a} dw ds$$
$$\lesssim \int_0^t \frac{\mu_0^{\nu_2}(s)}{(t-s)^{a/2}} \int_{\frac{\mu_0(s)R(s)}{4(t-s)} \le |\tilde{w}|} \frac{e^{-\frac{|\tilde{x}-\tilde{w}|^2}{4}}}{|\tilde{w}|^a} d\tilde{w} ds,$$

where $\tilde{x} = x(t-s)^{-1/2}$. Notice that for A > 0 we have

$$\int_{A<|\tilde{w}|} \frac{e^{-\frac{|\tilde{x}-\tilde{w}|^2}{4}}}{|\tilde{w}|^a} d\tilde{w} \lesssim \frac{1}{A^a} \text{ for } 0 < a \leq 2.$$

So

$$\int_{0}^{t-\mu_{0}^{2}(t)R^{2}(t)} \frac{\mu_{0}^{\nu_{2}}(s)}{(t-s)^{a/2}} \int_{\frac{\mu_{0}(s)R(s)}{\sqrt{t-s}} \leq |\tilde{w}|} \frac{e^{-\frac{|\tilde{x}-\tilde{w}|^{2}}{4}}}{|\tilde{w}|^{a}} d\tilde{w} ds$$

$$\lesssim \int_{0}^{t-\mu_{0}^{2}(t)R^{2}(t)} \frac{\mu_{0}^{\nu_{2}}(s)}{(t-s)^{a/2}} ds$$

$$\lesssim \mu_{0}^{\nu_{2}+\frac{2-a}{4k}}(0)$$
(A.10)

and

$$\int_{t-\mu_0^2(t)R^2(t)}^{t} \frac{\mu_0^{\nu_2}(s)}{(t-s)^{a/2}} \int_{\frac{\mu_0(s)R(s)}{\sqrt{t-s}} \le |\tilde{w}|} \frac{e^{-\frac{|\tilde{x}-\tilde{w}|^2}{4}}}{|\tilde{w}|^a} d\tilde{w} ds$$

$$\lesssim \int_{t-\mu_0^2(t)R^2(t)}^{t} \frac{\mu_0^{\nu_2}(s)}{(\mu_0(s)R(s))^a} ds$$

$$\lesssim \mu_0^{\nu_2+2-a}(0)R^{2-a}(0). \tag{A.11}$$

By (A.10)–(A.11), we conclude the validity of (4.12).

To prove (4.14), we have

$$\begin{split} |\nabla \psi(x,t)| \lesssim & \int_0^t \frac{\mu_0^{\nu_2}(s)}{(t-s)^{5/2}} \int_{\mu_0(s)R(s) \leq |w|} \frac{e^{-\frac{|x-w|^2}{4(t-s)}}|x-w|}{|w|^a} dw ds \\ \lesssim & \int_0^t \frac{\mu_0^{\nu_2}(s)}{(t-s)^{(a+1)/2}} \int_{\frac{\mu_0(s)R(s)}{\sqrt{t-s}} \leq |\tilde{w}|} \frac{e^{-\frac{|\tilde{x}-\tilde{w}|^2}{4}}|\tilde{x}-\tilde{w}|}{|\tilde{w}|^a} d\tilde{w} ds \\ \lesssim & \int_0^{t-\mu_0^2(t)R^2(t)} \frac{\mu_0^{\nu_2}(s)}{(t-s)^{(a+1)/2}} ds + \int_{t-\mu_0^2(t)R^2(t)}^t \frac{\mu_0^{\nu_2}(s)}{(t-s)^{(a+1)/2}} \frac{(t-s)^{a/2}}{(\mu_0(s)R(s))^a} ds \\ \lesssim & \mu_0^{\nu_2 + \frac{1-a_2}{4k}}(0). \end{split}$$

All the rest estimates can be proved similarly.

Acknowledgements: M. del Pino has been supported by a UK Royal Society Research Professorship. M. Musso is partly supported by EPSRC of UK. The research of J. Wei is partially supported by NSERC of Canada.

References

- [1] C. Collot, Non radial type II blow up for the energy supercritical semilinear heat equation. Anal. PDE 10 (2017) 127–252.
- [2] C. Collot, F. Merle, P. Raphael, Dynamics near the ground state for the energy critical nonlinear heat equation in large dimensions. Comm. Math. Phys. 352 (2017), 103-157.
- [3] C. Collot, F. Merle, P. Raphael, On strongly anisotropic type II blow up, To appear in J. Amer. Math. Soc.
- [4] C. Collot, P. Raphael, J. Szeftel On the stability of type I blow up for the energy super critical heat equation. Preprint ArXiv:1605.07337
- [5] C. Cortázar, M. del Pino, M. Musso, Green's function and infinite-time bubbling in the critical nonlinear heat equation. Preprint arXiv:1604.07117. To appear in J. Eur. Math. Soc. (JEMS).
- [6] J. Dávila, M. del Pino, J. Wei, Singularity formation for the two-dimensional harmonic map flow into S². Inventione Mathematicae 219 (2020), no.2, 345466.
- [7] M. del Pino, M. Musso, J. Wei, Infinite time blow-up for the 3-dimensional energy critical heat equation. To appear in Anal. PDE. Preprint ArXiv:1705.01672.
- [8] M. del Pino, M. Musso, J. Wei, Geometry driven Type II higher dimensional blow-up for the critical heat equation. Preprint Arxiv:1710.11461.
- [9] M. del Pino, M. Musso, J. Wei, Type II blow-up in the 5-dimensional energy critical heat equation. To appear in Acta Mathematica Sinica (Special issue in honor of Carlos Kenig).
- [10] Manuel del Pino, Monica Musso, J. Wei and Y. Zhou. Type II finite time blow-up for the energy critical heat equation in R⁴. To appear in DCDS-A (Special volume for W.-M. Ni's 70th birthday)
- [11] Kai Diethelm. The analysis of fractional differential equations: An application-oriented exposition using differential operators of Caputo type. Springer Science & Business Media, 2010.
- [12] R. Donninger, J. Krieger, Nonscattering solutions and blowup at infinity for the critical wave equation, <u>Math. Ann.</u> 357 (2013), no. 1, 89-163
- [13] T.Duyckaerts, C.E. Kenig, F. Merle, Classification of radial solutions of the focusing, energy-critical wave equation, Cambridge Journal of Mathematics 1 (2013), no. 1, 75–144.
- [14] S. Filippas, M.A. Herrero, J.J.L. Velázquez, Fast blow-up mechanisms for sign-changing solutions of a semilinear parabolic equation with critical nonlinearity. <u>R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci.</u> 456 (2000), no. 2004, 2957–2982.
- [15] Giga, Yoshikazu; Kohn, Robert V. Asymptotically self-similar blow-up of semilinear heat equations. <u>Comm. Pure Appl. Math.</u> 38 (1985), no. 3, 297-319.
- [16] Giga, Yoshikazu; Kohn, Robert V. Characterizing blowup using similarity variables. <u>Indiana Univ. Math. J.</u> 36 (1987), no. 1, 1-40.
- [17] Giga, Yoshikazu; Kohn, Robert V. Nondegeneracy of blowup for semilinear heat equations. <u>Comm. Pure Appl. Math.</u> 42 (1989), no. 6, 845-884.
- [18] J. Harada, A higher speed type II blowup for the five dimensional energy critical heat equation, to appear in Ann. Inst. H. Poincare Anal. Non Lineaire.
- [19] J. Harada, A type II blowup for the six dimensional energy critical heat equation, arXiv:2002.00528.
- [20] M.A. Herrero, J.J.L. Velázquez. Explosion de solutions d'equations paraboliques semilineaires supercritiques. [Blowup of solutions of supercritical semilinear parabolic equations] <u>C. R. Acad. Sci. Paris Ser. I Math.</u> 319 (1994), no. 2, 141-145.
- [21] M.A. Herrero, J.J.L. Velázquez. A blow up result for semilinear heat equations in the supercritical case. Unpublished.
- [22] C. E. Kenig and F. Merle, Global well-posedness, scattering and blow-up for the energy-critical focusing non-linear wave equation. Acta Math. 201, 147–212 (2008)
- [23] J. Krieger, W. Schlag, Concentration compactness for critical wave maps, Monographs of the European Mathematical Society (2012).
- [24] J. Krieger, W. Schlag, Full range of blow up exponents for the quintic wave equation in three dimension, <u>J. Math. Pures Appl.</u> (9) 101 (2014), no.6, 873–900
- [25] J. Krieger, K. Nakanishi, W. Schlag. Center-stable manifold of the ground state in the energy space for the critical wave equation. Math. Ann. 361 (2015), no. 1-2, 150.
- [26] J. Krieger, W. Schlag, D. Tataru, Slow blow-up solutions for the $H^1(\mathbb{R}^3)$ critical focusing semilinear wave equation. <u>Duke Math. J.</u> 147 (2009), no. 1, 1-53.
- [27] F.H. Lin, C.Y. Wang, The analysis of harmonic maps and their heat flows. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2008.
- [28] F. Merle and P. Raphael. The blow-up dynamic and upper bound on the blow-up rate for critical nonlinear Schrödinger equation, Ann. of Math. 161 (2005), no. 1, 157-222.
- [29] H. Matano, F. Merle. On nonexistence of type II blowup for a supercritical nonlinear heat equation, <u>Comm. Pure Appl. Math.</u> 57 (2004) 1494-1541.
- [30] H. Matano, F. Merle. Threshold and generic type I behaviors for a supercritical nonlinear heat equation. <u>J. Funct.</u> Anal. 261 (2011), no. 3, 716–748.

- [31] F. Merle, P. Raphael, I. Rodnianski, Blow-up dynamics for smooth data equivariant solutions to the critical Schrödinger map problem. Invent. Math. 193 (2013), no. 2, 249-365.
- [32] F. Merle and H. Zaag. Stability of the blow-up profile for equations of the type $u_t = \Delta u + |u|^{p-1}u$. Duke Math. J. 86(1) (1997) 143–195.
- [33] Mizoguchi, N. and Souplet, P. Optimal condition for blow-up of the critical L^q norm for the semilinear heat equation, arXiv:1812.11352
- [34] P. Polacik, E. Yanagida, On bounded and unbounded global solutions of a supercritical semilinear heat equation, Math. Annalen 327 (2003), 745-771.
- [35] P. Polacik, E. Yanagida, Global unbounded solutions of the Fujita equation in the intermediate range. Math. Ann. 360 (2014), no. 1-2, 255-266.
- [36] P. Quittner, Ph. Souplet. Superlinear parabolic problems. Blow- up, global existence and steady states. Birkhauser Advanced Texts. Birkhauser, Basel, 2007.
- [37] P. Raphael, I. Rodnianksi, Stable blow up dynamics for the critical co-rotational wave maps and equivariant YangMills problems. Publ. Math. Inst. Hautes Etudes Sci. 115 (2012), 1-122.
- [38] P. Raphaël, R. Schweyer, Stable blowup dynamics for the 1-corotational energy critical harmonic heat flow. Comm. Pure Appl. Math. 66 (2013), no. 3, 414–480.
- [39] P. Raphael and R. Schweyer, Quantized slow blow-up dynamics for the corotational energy-critical harmonic heat flow. Anal. PDE 7 (2014), no. 8, 17131805.
- [40] R. Schweyer. Type II blow-up for the four dimensional energy critical semi linear heat equation. <u>J. Funct. Anal.</u> 263 (2012), no. 12, 3922-3983.

Department of Mathematical Sciences, University of Bath, Bath BA2 7AY, United Kingdom $E\text{-}mail\ address:\ m.delpino@bath.ac.uk}$

Department of Mathematical Sciences, University of Bath, Bath BA2 7AY, United Kingdom $E\text{-}mail\ address:\ m.musso@bath.ac.uk}$

Department of Mathematics, University of British Columbia, Vancouver, B.C., V6T 1Z2, Canada E-mail address: jcwei@math.ubc.ca

Department of Mathematics, University of British Columbia, Vancouver, B.C., V6T 1Z2, Canada $E\text{-}mail\ address:\ \mathtt{qidi@math.ubc.ca}$

Department of Mathematics, University of British Columbia, Vancouver, B.C., V6T 1Z2, Canada E-mail address: yfzhou@math.ubc.ca