



Infection Prevention and Control Strategies for Seasonal Influenza in Healthcare Settings



For Health Care Providers

APR. 28, 2025

WHAT TO KNOW

This guidance emphasizes the importance of a comprehensive influenza prevention and control strategy that can be applied across the entire spectrum of healthcare settings. CDC will continue to evaluate new information as it becomes available and will update or expand this guidance as needed.

For Awareness

Updates: Recommendations for duration of work exclusion for healthcare personnel with Seasonal Influenza have been reviewed as part of updates to the **Guideline for Infection Control in Healthcare Personnel, 1998**. The draft has been approved by the Healthcare Infection Control Practices Advisory Committee (HICPAC) for posting in the Federal Register for a public comment period before being returned to HICPAC for additional review. Information about how to access the guideline in the Federal Register and further updates about the guideline development are available at: [HICPAC Guideline Updates](#) | [HICPAC](#) | [CDC](#)

Definition of Healthcare Settings

For the purposes of this guidance, healthcare settings include, but are not limited to, acute-care hospitals; long-term care facilities, such as nursing homes and skilled nursing facilities; physicians' offices; urgent-care centers, outpatient clinics; and home healthcare. This guidance is not intended to apply to other settings whose primary purpose is not healthcare, such as schools or worksites, because many of the aspects of the populations and feasible countermeasures will differ substantially across settings. However, elements of this guidance may be applicable to specific sites within non-healthcare settings where care is routinely delivered (e.g., a medical clinic embedded within a workplace or school).

Note

As the signs and symptoms of influenza can be similar to SARS-CoV-2 infection and other respiratory infections, please also refer to

[Preventing Transmission of Viral Respiratory Pathogens in Healthcare Settings](#)

Specific guidance for SARS-CoV-2, including diagnostic considerations and recommendations on the administration of SARS-CoV-2 immunization can be found on the [COVID-19 website](#).

Definition of Healthcare Personnel

HCP refers to all persons, paid and unpaid, working in healthcare settings who have the potential for exposure to patients and/or to infectious materials, including body substances, contaminated medical supplies and equipment, contaminated environmental surfaces, or contaminated air. HCP include, but are not limited to, physicians, nurses, nursing assistants, therapists, technicians, emergency medical service personnel, dental personnel, pharmacists, laboratory personnel, autopsy personnel, students and trainees, contractual personnel, home healthcare personnel, and persons not directly involved in patient care (e.g., clerical, dietary, house-keeping, laundry, security, maintenance, billing, chaplains, and volunteers) but potentially exposed to infectious agents that can be transmitted to and from HCP and patients. This guidance is not intended to apply to persons outside of healthcare settings for reasons discussed in the previous section.

Introduction

Influenza is an acute viral respiratory illness that is primarily transmitted in households and community settings. Each year, an estimated 5% to 20% of U.S. residents have influenza virus infection, an [estimated 3-11% are symptomatic](#) ², and many will seek medical care in ambulatory healthcare settings (e.g., pediatricians' offices, urgent-care clinics). In addition, an [estimated 100,000 to 710,000 persons are hospitalized each year for influenza-related complications](#) in the United States. Healthcare-associated influenza infections can occur in any healthcare setting and are most common when influenza is also circulating in the community. Therefore, the influenza prevention measures outlined in this guidance should be implemented in all healthcare settings. Supplemental measures may need to be implemented during influenza season if outbreaks of healthcare-associated influenza occur within certain facilities, such as long-term care facilities and hospitals.

KEEP READING[Interim Guidance for Influenza Outbreak Management in Long-Term Care and Post-Acute Care Facilities](#)

Influenza Modes of Transmission

Traditionally, influenza viruses have been thought to spread from person to person primarily through large-particle respiratory droplets and small particles in the air (e.g., when an infected person coughs or sneezes near a susceptible person). Transmission via large-particle droplets occurs during close contact between source and recipient persons, because droplets generally travel only short distances (approximately 6 feet or less) through the air. Indirect contact transmission via hand transfer of influenza virus from virus-contaminated surfaces or objects to mucosal surfaces of the face (e.g., nose, mouth) may also occur. Transmission over longer distances through the air, such as from one patient room to another has not been documented and is thought not to occur. All respiratory secretions and bodily fluids, including diarrheal stools, of patients with influenza are considered to be potentially infectious; however, the risk may vary by strain. Detection of influenza virus in blood or stool in influenza infected patients is very uncommon.

FOR EVERYONE[How Flu Spreads](#)

Fundamental Elements to Prevent Influenza Virus Transmission

Preventing transmission of influenza virus and other infectious agents within healthcare settings requires a multi-faceted approach. Spread of influenza virus can occur among patients, HCP, and visitors; in addition, HCP may acquire influenza from persons in their household or community. The core prevention strategies include:

- administration of influenza vaccine
- implementation of respiratory hygiene and cough etiquette
- appropriate management of ill HCP
- adherence to infection control precautions for all patient-care activities and aerosol-generating procedures
- implementing environmental and engineering infection control measures.

Successful implementation of many, if not all, of these strategies is dependent on the presence of clear administrative policies and organizational leadership that promote and facilitate adherence to these recommendations among the various people within the healthcare setting, including patients, visitors, and HCP. These administrative measures are included within each recommendation where appropriate. Furthermore, this guidance should be implemented in the context of a comprehensive infection prevention program to prevent transmission of all infectious agents among patients and HCP.

Recommendations

1. Promote and administer seasonal influenza vaccine

Annual influenza vaccination is the most important measure to prevent seasonal influenza infection. Achieving high influenza vaccination rates of HCP and patients is a critical step in preventing healthcare transmission of influenza from HCP to patients and from patients to HCP. According to current national guidelines, unless contraindicated, vaccinate all people aged 6 months and older, including HCP, patients and residents of long-term care facilities.

[Seasonal Influenza Vaccination Resources for Health Professionals](#)

Systematic strategies employed by some institutions to improve HCP vaccination rates have included providing incentives, providing influenza vaccine at no cost to HCP, improving access (e.g., offering influenza vaccination at work and during work hours), requiring personnel to sign declination forms to acknowledge that they have been educated about the benefits and risks of vaccination, and mandating influenza vaccination for all HCP without contraindication. Many of these approaches have been shown to increase influenza vaccination rates; [tracking influenza](#)

[vaccination](#) coverage among HCP can be an important component of a systematic approach to protecting patients and HCP. Regardless of the strategy used, strong organizational leadership and an infrastructure for clear and timely communication and education, and for program implementation, have been common elements in successful programs. More information on different HCP influenza vaccination strategies can be found in the Appendix: Influenza Vaccination Strategies.

2. Take Steps to Minimize Potential Exposures

A range of administrative policies and practices can be used to minimize influenza exposures before arrival, upon arrival, and throughout the duration of the visit to the healthcare setting. Measures include screening and triage of symptomatic patients and implementation of respiratory hygiene and cough etiquette. Respiratory hygiene and cough etiquette are measures designed to minimize potential exposures of all respiratory pathogens, including influenza virus, in healthcare settings and should be adhered to by everyone – patients, visitors, and HCP – upon entry and continued for the entire duration of stay in healthcare settings.

KEEP READING

[Preventing Transmission of Viral Respiratory Pathogens in Healthcare Settings](#)

Before Arrival to a Healthcare Setting

- When scheduling appointments, instruct patients and persons who accompany them to inform HCP upon arrival if they have symptoms of any respiratory infection (e.g., cough, runny nose, fever) and to take appropriate preventive actions (e.g., wear a facemask upon entry, follow triage procedure).
- During periods of increased influenza activity:
 - Take steps to minimize elective visits by patients with suspected or confirmed influenza. For example, consider establishing procedures to minimize visits by patients seeking care for mild influenza-like illness who are not at increased risk for complications from influenza (e.g., provide telephone consultation to patients with mild respiratory illness to determine if there is a medical need to visit the facility).

Upon Entry and During Visit to a Healthcare Setting

- Take steps to ensure all persons with symptoms of a respiratory infection adhere to respiratory hygiene, cough etiquette, hand hygiene, and triage procedures throughout the duration of the visit. These might include:
 - Posting visual alerts (e.g., signs, posters) at the entrance and in strategic places (e.g., waiting areas, elevators, cafeterias) to provide patients and HCP with instructions (in appropriate languages) about respiratory hygiene and cough etiquette, especially during periods when influenza virus is circulating in the community. Instructions should include:
 - How to use facemasks or tissues to cover nose and mouth when coughing or sneezing and to dispose of contaminated items in waste receptacles.
 - How and when to perform hand hygiene.
 - Implementing procedures during patient registration that facilitate adherence to appropriate precautions (e.g., at the time of patient check-in, inquire about presence of symptoms of a respiratory infection, and if present, provide instructions).
- Provide facemasks (definition of facemask in Appendix) to patients with signs and symptoms of respiratory infection.
- Provide supplies to perform hand hygiene to all patients upon arrival to facility (e.g., at entrances of facility, waiting rooms, at patient check-in) and throughout the entire duration of the visit to the healthcare setting.
- Provide space and encourage persons with symptoms of respiratory infections to sit as far away from others as possible. If available, facilities may wish to place these patients in a separate area while waiting for care.
- During periods of increased community influenza activity, facilities should consider setting up triage stations that facilitate rapid screening of patients for symptoms of influenza and separation from other patients.

3. Monitor and Manage Ill Healthcare Personnel

HCP who develop fever and respiratory symptoms should be:

- Instructed not to report to work, or if at work, to stop patient-care activities, don a facemask, and promptly notify their supervisor and infection control personnel/occupational health before leaving work.
- Reminded that adherence to respiratory hygiene and cough etiquette after returning to work is always important. If symptoms such as cough and sneezing are still present, HCP should wear a facemask during patient-care activities. The importance of performing frequent hand hygiene (especially before and after each patient contact and contact with respiratory secretions) should be reinforced.
- Excluded from work until at least 24 hours after they no longer have a fever (without the use of fever-reducing medicines such as acetaminophen). Those with ongoing respiratory symptoms should be considered for evaluation by occupational health to determine appropriateness of contact with patients.

- Considered for temporary reassignment or exclusion from work for 7 days from symptom onset or until the resolution of symptoms, whichever is longer, if returning to care for patients in a [Protective Environment \(PE\)](#) [PDF](#) such as hematopoietic stem cell transplant patients (HSCT).
 - Patients in these environments are severely immunocompromised, and infection with influenza virus can lead to severe disease. Furthermore, once infected, these patients can have prolonged viral shedding despite antiviral treatment and expose other patients to influenza virus infection. Prolonged shedding also increases the chance of developing and spreading antiviral-resistant influenza strains; clusters of influenza antiviral resistance cases have been found among severely immunocompromised persons exposed to a common source or healthcare setting.
- HCP with influenza or many other infections may not have fever or may have fever alone as an initial symptom or sign. Thus, it can be very difficult to distinguish influenza from many other causes, especially early in a person's illness. HCP with fever alone should follow workplace policy for HCP with fever until a more specific cause of fever is identified or until fever resolves.

HCP who develop acute respiratory symptoms without fever may still have influenza infection and should be:

- Considered for evaluation by occupational health to determine appropriateness of contact with patients. HCP suspected of having influenza may benefit from influenza antiviral treatment.
- Reminded that adherence to respiratory hygiene and cough etiquette after returning to work is always important. If symptoms such as cough and sneezing are still present, HCP should wear a facemask during patient care activities. The importance of performing frequent hand hygiene (especially before and after each patient contact) should be reinforced.
- Allowed to continue or return to work unless assigned to care for patients requiring a [PE](#) [PDF](#) such as HSCT; these HCP should be considered for temporary reassignment or considered for exclusion from work for 7 days from symptom onset or until the resolution of all non-cough symptoms, whichever is longer.

Facilities and organizations providing healthcare services should:

- Develop sick leave policies for HCP that are non-punitive, flexible and consistent with public health guidance to allow and encourage HCP with suspected or confirmed influenza to stay home.
 - Policies and procedures should enhance exclusion of HCPs who develop a fever and respiratory symptoms from work for at least 24 hours after they no longer have a fever, without the use of fever-reducing medicines.
- Ensure that all HCP, including staff who are not directly employed by the healthcare facility but provide essential daily services, are aware of the sick leave policies.
- Employee health services should establish procedures for tracking absences; reviewing job tasks and ensuring that personnel known to be at higher risk for exposure to those with suspected or confirmed influenza are given priority for vaccination; ensuring that employees have prompt access, including via telephone to medical consultation and, if necessary, early treatment; and promptly identifying individuals with possible influenza. HCP should self-assess for symptoms of febrile respiratory illness. In most cases, decisions about work restrictions and assignments for personnel with respiratory illness should be guided by clinical signs and symptoms rather than by laboratory testing for influenza because laboratory testing may result in delays in diagnosis, false negative test results, or both.

4. Adhere to Standard Precautions

During the care of any patient, all HCP in every healthcare setting should adhere to [Standard Precautions](#), which are the foundation for preventing transmission of infectious agents in all healthcare settings. Standard Precautions assume that every person is potentially infected or colonized with a pathogen that could be transmitted in the healthcare setting. Further information about Standard Precautions, including hand hygiene, respiratory hygiene and cough etiquette and proper selection and use of personal protective equipment can be found in [CDC's Core Infection Prevention and Control Practices for Safe Healthcare Delivery in All Settings](#).

5. Adhere to Droplet Precautions

- Droplet Precautions should be implemented for patients with suspected or confirmed influenza for 7 days after illness onset or until 24 hours after the resolution of fever and respiratory symptoms, whichever is longer, while a patient is in a healthcare facility. In some cases, facilities may choose to apply [Droplet Precautions](#) for longer periods based on clinical judgment, such as in the case of young children or severely immunocompromised patients, who may shed influenza virus for longer periods of time.
- Place patients with suspected or confirmed influenza in a private room or area. When a single patient room is not available, consultation with infection control personnel is recommended to assess the risks associated with other patient placement options (e.g., cohorting [i.e., grouping patients infected with the same infectious agents together to confine their care to one area and prevent contact with susceptible patients], keeping the patient with an existing roommate). For more information about making decisions on patient placement for Droplet Precautions, refer to the CDC Guideline for Isolation Precautions [\[section V.C.2\]](#).
- HCP should don a facemask when entering the room of a patient with suspected or confirmed influenza. Remove the facemask when leaving the patient's room, dispose of the facemask in a waste container, and perform hand hygiene.
 - If some facilities and organizations opt to provide employees with alternative personal protective equipment, this equipment should provide the same protection of the nose and mouth from splashes and sprays provided by facemasks (e.g., face shields and N95 respirators or powered air purifying respirators).

- If a patient under Droplet Precautions requires movement or transport outside of the room:
 - Have the patient wear a facemask, if possible, and follow respiratory hygiene and cough etiquette and hand hygiene.
 - Communicate information about patients with suspected, probable, or confirmed influenza to appropriate personnel before transferring them to other departments in the facility (e.g., radiology, laboratory) or to other facilities.
- Patients under Droplet Precautions should be discharged from medical care when clinically appropriate, not based on the period of potential virus shedding or recommended duration of Droplet Precautions. Before discharge, communicate the patient's diagnosis and current precautions with post-hospital care providers (e.g., home-healthcare agencies, long-term care facilities) as well as transporting personnel.

6. Use Caution when Performing Aerosol-Generating Procedures

Some procedures performed on patients with suspected or confirmed influenza infection may be more likely to generate higher concentrations of infectious respiratory aerosols than coughing, sneezing, talking, or breathing. These procedures potentially put HCP at an increased risk for influenza exposure. Although there are limited data available on influenza transmission related to such aerosols, many authorities such as the World Health Organization recommend that [additional precautions be used when such procedures are performed](#) ². These include some procedures that are usually planned ahead of time, such as bronchoscopy, sputum induction, elective intubation and extubation, and autopsies; and some procedures that often occur in unplanned, emergent settings and can be life-saving, such as cardiopulmonary resuscitation, emergent intubation and open suctioning of airways. Ideally, a combination of measures should be used to reduce exposures from these aerosol-generating procedures when performed on patients with suspected or confirmed influenza. However, it is appropriate to take feasibility into account, especially in challenging emergent situations, where timeliness in performing a procedure can be critical to achieving a good patient outcome. Precautions for aerosol-generating procedures include:

- Only performing these procedures on patients with suspected or confirmed influenza if they are medically necessary and cannot be postponed.
- Limiting the number of HCP present during the procedure to only those essential for patient care and support. As is the case for all HCP, ensure that HCP whose duties require them to perform or be present during these procedures are offered influenza vaccination.
- Conducting the procedures in an airborne infection isolation room (AIIR) when feasible. This will not be feasible for unplanned, emergent procedures, unless the patient is already in an AIIR. Such rooms are designed to reduce the concentration of infectious aerosols and prevent their escape into adjacent areas using controlled air exchanges and directional airflow. They are single patient rooms at negative pressure relative to the surrounding areas, and with a minimum of 6 air changes per hour (12 air changes per hour are recommended for new construction or renovation). Air from these rooms should be exhausted directly to the outside or be filtered through a high-efficiency particulate air (HEPA) filter before recirculation. Room doors should be kept closed except when entering or leaving the room, and entry and exit should be minimized during and shortly after the procedure. Facilities should [monitor and document](#) the proper negative-pressure function of these rooms.
- Considering use of portable HEPA filtration units to further reduce the concentration of contaminants in the air. Some of these units can connect to local exhaust ventilation systems (e.g., hoods, booths, tents) or have inlet designs that allow close placement to the patient to assist with source control; however, these units do not eliminate the need for respiratory protection for individuals entering the room because they may not entrain all of the room air. Information on air flow/air entrainment performance should be evaluated for such devices.
- HCP should wear gloves, a gown, and either a face shield that fully covers the front and sides of the face or goggles.
- HCP should wear respiratory protection equivalent to a fitted N95 filtering facepiece respirator or equivalent N95 respirator (e.g., powered air purifying respirator, elastomeric) during aerosol-generating procedures (definition of respirator in Appendix).
 - When respiratory protection is required in an occupational setting, respirators must be used in the context of a comprehensive respiratory protection program that includes fit-testing and training as required under OSHA's Respiratory Protection standard ([29 CFR 1910.134](#) ³).
- Unprotected HCP should not be allowed in a room where an aerosol-generating procedure has been conducted until sufficient time has elapsed to remove potentially infectious particles. More information on [clearance rates](#) under differing ventilation conditions is available.
- Conduct environmental surface cleaning following procedures (section on environmental infection control).

7. Manage Visitor Access and Movement Within the Facility

- For the safety of the visitor, in general, patients with influenza should be encouraged to limit in-person visitation while they are infectious. However, facilities should adhere to local, territorial, tribal, state, and federal regulations related to visitation.
 - Counsel patients and their visitor(s) about the risks of an in-person visit.
 - Encourage use of alternative mechanisms for patient and visitor interactions such as video-call applications on cell phones or tablets, when appropriate.
- Facilities should provide instruction, before visitors enter the patient's room, on hand hygiene, limiting surfaces touched, and use of PPE according to current facility policy.
- Visitors should be instructed to only visit the patient room. They should minimize their time spent in other locations in the facility.

- Visitors who have been in contact with the patient before and during hospitalization are a possible source of influenza for other patients, visitors, and staff.

8. Monitor Influenza Activity

Healthcare settings should establish mechanisms and policies by which HCP are promptly alerted about increased influenza activity in the community or if an outbreak occurs within the facility and when collection of clinical specimens for viral culture may help to inform public health efforts. Close communication and collaboration with local and state health authorities is recommended. Policies should include designations of specific persons within the healthcare facility who are responsible for communication with public health officials and dissemination of information to HCP.

9. Implement Environmental Infection Control

Detailed information on environmental cleaning in healthcare settings can be found in CDC's [Guidelines for Environmental Infection Control in Health-Care Facilities](#).

Standard cleaning and disinfection procedures (e.g., using cleaners and water to preclean surfaces prior to applying disinfectants to frequently touched surfaces or objects for indicated contact times) are adequate for influenza virus environmental control in all settings within the healthcare facility, including those patient-care areas in which aerosol-generating procedures are performed. Management of laundry, food service utensils, and medical waste should also be performed in accordance with standard procedures. There are no data suggesting these items are associated with influenza virus transmission when these items are properly managed. Laundry and food service utensils should first be cleaned, then sanitized as appropriate. Some medical waste may be designated as regulated or biohazardous waste and require special handling and disposal methods approved by the State authorities.

10. Implement Engineering Controls

Consider designing and installing engineering controls to reduce or eliminate exposures by shielding HCP and other patients from infected individuals. Examples of engineering controls include installing physical barriers such as partitions in triage areas or curtains that are drawn between patients in shared areas. Engineering controls may also be important to reduce exposures related to specific procedures such as using closed suctioning systems for airways suction in intubated patients. Another important engineering control is ensuring that [appropriate air-handling systems](#) are installed and maintained in healthcare facilities.

11. Train and Educate Healthcare Personnel

Healthcare administrators should ensure that all HCP receive job- or task-specific education and training on preventing transmission of infectious agents, including influenza, associated with healthcare during orientation to the healthcare setting. This information should be updated periodically during ongoing education and training programs. Competency should be documented initially and repeatedly, as appropriate, for the specific staff positions. A system should be in place to ensure that HCP employed by outside employers meet these [education and training requirements](#) through programs offered by the outside employer or by participation in the healthcare facility's program.

- Key aspects of influenza and its prevention that should be emphasized to all HCP include:
 - Influenza signs, symptoms, complications, and risk factors for complications. HCP should be made aware that, if they have conditions that place them at higher risk of complications, they should inform their healthcare provider immediately if they become ill with an influenza-like illness so they can receive early treatment if indicated.
 - Central role of administrative controls such as vaccination, respiratory hygiene and cough etiquette, sick policies, and precautions during aerosol-generating procedures.
 - Appropriate use of personal protective equipment including respirator fit testing and fit checks.
 - Use of engineering controls and work practices including infection control procedures to reduce exposure.

12. Administer Antiviral Treatment and Chemoprophylaxis of Patients and Healthcare Personnel when Appropriate

Refer to the CDC web site for the most current recommendations on the use of antiviral agents for influenza treatment and chemoprophylaxis. Both HCP and patients should be reminded that persons treated with influenza antiviral medications continue to shed influenza virus while on treatment. Thus, hand hygiene, respiratory hygiene and cough etiquette practices should continue while on treatment.

KEEP READING

[About Influenza Antiviral Medications](#)

13. Considerations for Healthcare Personnel at Higher Risk for Complications of Influenza

[HCP at increased risk](#) for complications from influenza include pregnant women and women up to 2 weeks postpartum, persons 65 years old and older, and persons with chronic diseases such as asthma, heart disease, diabetes, diseases that suppress the immune system, certain other

chronic medical conditions, and morbid obesity. Influenza vaccination for prevention and early treatment with antiviral medications if symptomatic are very important for HCP at higher risk for influenza complications because they can decrease the risk of hospitalizations and deaths. HCP at increased risk for influenza complications should check with their healthcare provider as soon as possible if they become ill with influenza so that they can receive early initiation of antiviral treatment if indicated.

Some HCP may identify themselves as being at increased risk of complications, and express concerns about their risks. These concerns should be discussed and the importance of careful adherence to these guidelines should be emphasized.

Appendix: Additional Information about Influenza

Information about Facemasks:

- FDA – [Masks and N95 Respirators, FDA](#)
- A facemask is a loose-fitting, disposable device that creates a physical barrier between the mouth and nose of the wearer and potential contaminants in the immediate environment. Facemasks may be labeled as surgical, laser, isolation, dental or medical procedure masks. They may come with or without a face shield. If worn properly, a facemask is meant to help block large-particle droplets, splashes, sprays or splatter that may contain germs (viruses and bacteria) from reaching your mouth and nose. Facemasks may also help reduce exposure of the wearer's saliva and respiratory secretions to others. While a facemask may be effective in blocking splashes and large-particle droplets, a facemask, by design, does not filter or block very small particles in the air that may be transmitted by coughs, sneezes or certain medical procedures.
- Facemasks are cleared by the U.S. Food and Drug Administration (FDA) for use as medical devices. Facemasks should be used once and then thrown away in the trash.

Information about Respirators:

- FDA – [Masks and N95 Respirators, FDA](#)
- A respirator is a personal protective device that is worn on the face, covers at least the nose and mouth, and is used to reduce the wearer's risk of inhaling hazardous airborne particles (including dust particles and infectious agents), gases, or vapors. Respirators are certified by the National Institute for Occupational Safety and Health (NIOSH), CDC, and N95 respirators may also be cleared or approved by FDA as medical devices, depending on the intended use. A commonly used respirator is a filtering facepiece respirator (often referred to as an N95).
- To work properly, respirators must be specially fitted for each person who wears one (this is called "fit-testing" and is usually done in a workplace where respirators are used).
- [OSHA Respiratory Protection eTool](#)

Resources

- [About Influenza](#)
- [Clinical Information \(signs and symptoms of influenza\)](#)
- [World Health Organization \(WHO\). Epidemic- and pandemic-prone acute respiratory diseases – Infection prevention and control in health care](#)
- [Interim Guidance for Influenza Outbreak Management in Long-Term Care and Post-Acute Care Facilities](#)
- [Seasonal Influenza Vaccination Resources for Health Professionals](#)
- [Guidance for Prevention and Control of Influenza in the Peri- and Postpartum Settings](#)
- [About Influenza Antiviral Medications](#)

SOURCES AND PAGE INFO

CONTENT SOURCE:
National Center for Immunization and Respiratory Diseases (NCIRD)

ABOUT THIS PAGE

Published: October 1, 2024

Updated: April 28, 2025
This page was last updated on this date. Updates may include minor edits, image changes, or other modifications to page content.

Reviewed: April 28, 2025

The information on this page was last reviewed by subject matter experts to ensure accuracy.