


## HW Assignments


- **Exam 3** will be on 30<sup>th</sup> November (Friday)
  - ✓ Covers the Chapters 8, 9, 10 and 13
  - ✓ 25 MCQs
  - ✓ I will post a **practice exam** for Exam 3 during the break
- **Chapter 10** HW Due 26<sup>th</sup> November at 11:55 pm
- **Chapter 13** HW will assign today
- **Quiz 5** (Chapters 9, 10) will post on 26<sup>th</sup> (Monday) and due Wednesday before midnight



## ***Chapter 13*** **Electrochemistry**

Edited by Dr. Katugampola

## Corrosion



- **Corrosion** is the degradation of metals by chemical reactions
  - ✓ **Uniform corrosion** occurs evenly over a large portion of the surface area of a metal
  - ✓ **Galvanic corrosion** occurs when two different metals contact each other in the presence of an electrolyte
  - ✓ **Crevice corrosion** occurs in machines

Copyright ©2019 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

3


## Oxidation–Reduction Reactions and Galvanic Cells



- Reactions involving the transfer of electrons are known as **oxidation–reduction** or **redox reactions**
  - ✓ **Oxidation** is the **loss of electrons** from some chemical species




- ✓ **Reduction** is the **gain of electrons** by some chemical species



Copyright ©2019 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

4

## Oxidation–Reduction and Half-Reactions






- When a clean copper wire is placed into a colorless solution of silver nitrate, it is quickly apparent that a chemical reaction occurs: Crystals of silver metal form on the copper wire, and the solution turns blue

5




- The solution's blue color is indicative of  $\text{Cu}^{2+}$  ions in solution
  - ✓  $\text{Cu}^{2+}$  is formed when a copper atom loses two electrons
  - ✓ The copper metal is oxidized
- The crystals formed on the surface of the copper wire are silver metal
  - ✓ Metallic silver is formed when a silver cation gains an electron
  - ✓ The silver in the solution is reduced



Two Half-Reactions.

- The electron transfer must be balanced, so the reduction half-reaction is multiplied by 2





## Oxidation–Reduction and Half-Reactions

(continued)





- Add the two half-reactions together and the electrons will cancel out, leaving the **net ionic equation** for the redox reaction




## Oxidation–Reduction and Half-Reactions

(continued)



- **Reducing agent:** The species undergoing oxidation  
✓ The Cu was oxidized and is the reducing agent
- **Oxidizing agent:** The species undergoing reduction  
✓ The  $\text{Ag}^+$  was reduced and is the oxidizing agent
- Writing the **molecular equation** by adding the spectator ions ( $\text{NO}_3^-$  in this case)





Copyright ©2019 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

9



## Problem



13.5 For the following oxidation-reduction reactions, identify the half reactions and label them as oxidation or reduction.




Answer:



Copyright ©2019 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

10

## Galvanic Cell




- The **salt bridge** contains a strong electrolyte that allows either cations or anions to migrate into the solution where they are needed to maintain charge neutrality. **Also, it closes the circuit**

11

## Terminology for Galvanic Cells

- Electrodes**
  - ✓ The electrode where oxidation occurs is the **anode**
  - ✓ The electrode where reduction occurs is the **cathode**
- Cell notation:** A shorthand notation for representing the specific chemistry of an electrochemical cell
  - Cell notation lists the metals and ions involved in the reaction
  - A single vertical line,  $|$ , denotes **a phase boundary**
  - A double vertical line,  $||$ , denotes **a salt bridge**
  - The **anode** is written on the left, the **cathode** on the right

## Terminology for Galvanic Cells (continued)



- General form of cell notation  
 $\text{anode} \mid \text{anode electrolyte} \parallel \text{cathode electrolyte} \mid \text{cathode}$
- For the previous example of copper and silver  
 $\text{Cu(s)} \mid \text{Cu}^{2+}(\text{aq}) (1 \text{ M}) \parallel \text{Ag}^{+}(\text{aq}) (1 \text{ M}) \mid \text{Ag}$ 
  - The electrolyte concentration is also given
- An electrochemical cell is at its **standard state** when the electrolyte concentrations are **1 M**
- For half-cells that generate or consume a gas, a partial pressure of **1 atm** is required for the standard state

Copyright ©2019 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

13

- Maximum work that can be obtained from an electrochemical cell


$$w_{\max} = qE$$

- ✓  $q$  is the charge
- ✓  $E$  is the cell potential or electromotive force (EMF)

Copyright ©2019 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

14

## Problem



In a classroom demonstration of the reaction between copper metal and silver ions, silver plates out on the copper wire. Which is the oxidizing agent in this reaction?

- Silver metal
- Silver ions
- Copper metal
- Copper ions

Answer: Silver ions

Copyright ©2019 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

15

## Problem



In a galvanic cell, where does oxidation occur?

- In the electrolyte
- At the anode
- At the cathode
- In the salt bridge



Answer: At the anode

Copyright ©2019 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

16



## Measuring cell potential




- Measurement of standard cell voltages for various combinations of the same half-reactions suggests that a characteristic potential can be associated with a particular half-reaction

17

## Standard component in cell potential measurements


- Standard hydrogen electrode** or **SHE**
- The cell is constructed using a platinum wire or foil as the electrode
  - The electrode is immersed in a 1 M HCl solution, and  $\text{H}_2$  gas is bubbled over the electrode at a pressure of 1 atm
- chosen as the **reference point for the scale of standard reduction potentials** and assigned a potential of exactly **zero volts**



Half-reaction,  
 $2 \text{H}^+ (\text{aq}) + 2 \text{e}^- \rightarrow \text{H}_2 (\text{g})$

Half-cell notation,  
 $\text{Pt}(\text{s}) \mid \text{H}_2 (\text{g}, 1 \text{ atm}) \mid \text{H}^+ (1 \text{ M})$

## Measuring Cell Potential



- For some galvanic cells, the SHE acts as the anode, and for other galvanic cells, the SHE acts as the cathode (anode is the site of oxidation and cathode is the site of reduction)

Copyright ©2019 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

19

## Standard Reduction Potentials




Here we list standard reduction potentials for several of the half-reactions involved in the cells discussed in the text. A more extensive table of potentials appears in Appendix I.

Half-Reaction	Standard Reduction Potential (V)
$\text{Zn}^{2+} + 2 \text{e}^{-} \rightarrow \text{Zn}$	-0.763
$\text{Fe}^{2+} + 2 \text{e}^{-} \rightarrow \text{Fe}$	-0.44
$2 \text{H}^{+} + 2 \text{e}^{-} \rightarrow \text{H}_2$	0.000
$\text{Cu}^{2+} + 2 \text{e}^{-} \rightarrow \text{Cu}$	+0.337
$\text{Fe}^{3+} + \text{e}^{-} \rightarrow \text{Fe}^{2+}$	+0.771
$\text{Ag}^{+} + \text{e}^{-} \rightarrow \text{Ag}$	+0.7994

20

## Standard Reduction Potentials (continued 1)



- ✓ A large, positive value for the standard reduction potential implies the substance is reduced readily and is a good oxidizing agent
- ✓ A large, negative value for the standard reduction potential implies the substance is oxidized readily and is a good reducing agent

Copyright ©2019 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

21

## Standard Reduction Potentials (continued 2)



- For a galvanic cell,
  - ✓ the half-reaction with the more positive reduction potential will be the cathode
  - ✓ The half-reaction with the more negative reduction potential will be the anode
- The standard cell potential for any pair of half-reactions ( $E^\circ_{\text{cell}}$ )



$$E^\circ_{\text{cell}} = E^\circ_{\text{red}} - E^\circ_{\text{ox}}$$

- ✓  $E^\circ_{\text{red}}$  is the standard reduction potential of the cathode
- ✓  $E^\circ_{\text{ox}}$  is the standard reduction potential of the anode

Copyright ©2019 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.

22

## Standard Reduction Potentials (continued 3)





- Arranging standard reduction potentials horizontally enables easy identification of the anode and cathode in a galvanic cell
- The reduction potential farthest to the left is the anode
- The reduction potential farthest to the right is the cathode

23


## Example Problem



- 1) Using standard reduction potentials, identify the anode and the cathode and determine the cell potential for a galvanic cell composed of copper and iron. Assume standard conditions



Iron must be oxidized,



By using,  $E^\circ_{\text{cell}} = E^\circ_{\text{red}} - E^\circ_{\text{ox}}$

$$E^\circ_{\text{cell}} = 0.337 \text{ V} - (-0.44 \text{ V}) = 0.78 \text{ V}$$

Copper is the cathode, because it is reduced  
 Iron is the anode, because it is oxidized

24

Copyright ©2019 Cengage Learning. All Rights Reserved. May not be scanned, copied or duplicated, or posted to a publicly accessible website, in whole or in part.