

QE2012 AC-3 ECE580 - Rhea

Print

ECE Ph.D. Qualifying Exam Automatic Control (AC) Question 3: Optimization

August 2012

Student answers and discussions for Part 1,2,3,4,5

1.(20 pts)

(i)(10 pts) Find the factor by which the uncertainty range is reduced when using the Fibonacci method. Assume that the last step has the form $1ho_{N-1}=rac{F_2}{F_3}=rac{2}{3}$,

where N-1 is the number of steps performed in the uncertainty range reduction process.

(ii)(10 pts) It is known that the minimizer of a certain function f(x) is located in the interval[-5, 15]. What is the minimal number of iterations of the Fibonacci method required to box in the minimizer within the range 1.0? Assume that the last useful value of the factor reducing the uncertainty range is 2/3, that is

$$1-
ho_{N-1}=rac{F_2}{F_3}=rac{2}{3}\,,$$

Click here to view student answers and discussions

Problem 2. (20pts) Employ the DFP method to construct a set of Q-conjugate directions using the function

$$\begin{split} f &= \frac{1}{2} \, \boldsymbol{x}^T \boldsymbol{Q} \boldsymbol{x} - \boldsymbol{x}^T \boldsymbol{b} + \boldsymbol{c} \\ &= \frac{1}{2} \, \boldsymbol{x}^T \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} \boldsymbol{x} - \boldsymbol{x}^T \begin{bmatrix} 2 \\ 1 \end{bmatrix} + 3 \end{split}$$

Where $x^{(0)}$ is arbitrary.

Click here to view student answers and discussions

Problem 3. (20pts) For the system of linear equations, Ax = b where

$$A = \begin{bmatrix} 1 & 0 & -1 \\ 0 & 1 & 0 \\ 0 & -1 & 0 \end{bmatrix}, b = \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix}$$

Find the minimum length vector $x^{(*)}$ that minimizes $\|Ax-b\|_2^2$

Click here to view student answers and discussions

Problem 4. (20pts) Use any simplex method to solve the following linear program.

$$\begin{array}{ll} \textit{Maximize} & x_1 + 2x_2 \\ \textit{S'ubject to} & -2x_1 + x_2 \leq 2 \\ & x_1 - x_2 \geq -3 \\ & x_1 \leq -3 \\ & x_1 \geq 0, x_2 \geq 0. \end{array}$$

wers and discussions

Problem 5.(20pts) Solve the following problem:

$$\begin{array}{ll} \textit{Minimize} & -x_1^2 + 2x_2 \\ \textit{Subject to} & x_1^2 + x_2^2 \leq 1 \\ & x_1 \geq 0 \\ & x_2 \geq 0 \end{array}$$

(i)(10pts) Find the points that satisfy the KKT condition.

(ii)(10pts)Apply the SONC and SOSC to determine the nature of the critical points from the previous part.

Click here to view student answers and discussions

Back to ECE QE page

Retrieved from "https://www.projectrhea.org/rhea/index.php?title=QE2012_AC-3_ECE580&oldid=58217" Categories: ECE | QE | CNSIP | Problem solving | Automatic control | Optimization

Shortcuts

Help Main Wiki Page Random Page Special

Pages Log in

Search

Search

Alumni Liaison

To all math majors: "Mathematics is a wonderfully rich subject."

Read more »



This page was last modified on 13 September 2013, at 09:17. This page has been accessed 22,370 times.