



ulm university universität
uulm

**Fakultät für
Naturwissenschaften**

Institute of Theoretical
Physics

Title of the work

Bachelor Thesis

Submitted by:

Jan Bulling
jan.bulling@uni-ulm.de
1109395

Supervised by:

Marit O. E. Steiner, Julen S. Pedernales, Martin B. Plenio

Abstract

Contents

1	Introduction	4
1.1	Feynman's Gedankenexperiment	4
2	A first look at the problem	5
2.1	Entanglement measures	5
3	Casimir effect	7
3.1	Proximity force approximation	7
3.2	Imperfect plate and spheres	7
3.3	Casimir forces between a conducting plate and a dielectric sphere	7
3.3.1	Polarizability of a dielectric sphere	7
4	Plate vibrations	11
	Bibliography	12
A	Proofs and other stuff	14
A.1	Negativity	14
A.2	Fidelity	15

1 Introduction

1.1 Feynman's Gedankenexperiment

2 A first look at the problem

Problem, general calculations from Julens paper without casimir

When is the system entangled (distance-measures, fidelity ??)




Figure 2.1: Simple Problem from Julens paper

Ignoring (for now) all other forms of interaction between the masses except gravity

The density $\rho(t) = |\psi(t)\rangle\langle\psi(t)|$ can be represented in the basis $\{|\psi_A^\uparrow\rangle, |\psi_A^\downarrow\rangle\} \otimes \{|\psi_B^\uparrow\rangle, |\psi_B^\downarrow\rangle\}$ as

$$\rho(t) = \frac{1}{4} \begin{pmatrix} 1 & & & 1 \\ & 1 & & \\ & & 1 & \\ 1 & & & 1 \end{pmatrix} \quad (2.1)$$

2.1 Entanglement measures

Why are they needed, what can one do?

Logarithmic negativity, properties, calculation




Figure 2.2: My problem




Figure 2.3: Entanglement dynamics quantified by the logarithmic negativity for two different orientations of the spatial superpositions relative to each other. The time of maximum entanglement t_{\max} for the parallel configuration is given by $t_{\max} = 8\pi\hbar L^3/(GM_A M_B d^2) \simeq 258$ ms.

3 Casimir effect

General introduction and comparison with retarded van der Waals forces

$$F_{\text{Casimir}} = -\frac{\hbar c \pi^2}{240 L^4} A \quad (3.1)$$

$$V_{\text{Casimir}} = \frac{\hbar c \pi^2}{720 L^3} A \quad (3.2)$$

Lifshitz:

$$F_{\text{DD}} = \frac{\hbar c \pi^2}{240 L^4} \left(\frac{\varepsilon_r - 1}{\varepsilon_r + 1} \right)^2 \varphi(\varepsilon_r) \quad (3.3)$$

$$F_{\text{DM}} = \frac{\hbar c \pi^2}{240 L^4} \frac{\varepsilon_r - 1}{\varepsilon_r + 1} \varphi(\varepsilon_r) \quad (3.4)$$

The numeric function φ is shown in fig. 3.1.

3.1 Proximity force approximation

3.2 Imperfect plate and spheres

Python numerical approach, gaussian modes (vibration modes of a spherical plane), perlin noise

3.3 Casimir forces between a conducting plate and a dielectric sphere

3.3.1 Polarizability of a dielectric sphere

The polarizability α is defined via

$$\mathbf{E}_{\infty} \alpha = \mathbf{p}, \quad (3.5)$$




Figure 3.1: Numeric casimir interaction $\varphi(\epsilon_r)$ between **(blue)** two dielectric plates and **(orange)** a dielectric and a conductor.




Figure 3.2: In the proximity force approximation the sphere is divided into infinitesimal plane areas dA which all exert a force dF according to eq. (3.1). All the contributions are added up together.

where \mathbf{p} is the induced dipole moment and \mathbf{E}_∞ is the external electric field that induces the dipole moment. For a linear and uniform dielectric, it is given as $\mathbf{p} = \mathcal{V}\varepsilon_0(\varepsilon_r - 1)\mathbf{E}_\text{in}$ [1, p. 220-226]. Here, \mathcal{V} is the volume of the object and \mathbf{E}_in is the electric field inside the dielectric. The electrostatic boundary conditions for the problem are given by

$$V_\text{in}|_{r=R} = V_\text{out}|_{r=R} \quad \text{and} \quad \varepsilon_r \varepsilon_0 \frac{\partial V_\text{in}}{\partial r} \Big|_{r=R} = \varepsilon_0 \frac{\partial V_\text{out}}{\partial r} \Big|_{r=R} \quad (3.6)$$

and the electric potential outside of the sphere at $r \rightarrow \infty$ should be equal to the external dipole-inducing field $V_\text{out}|_{r \rightarrow \infty} = -\mathbf{E}_\infty \cdot \mathbf{r} = -E_\infty r \cos \theta$. The electric potential inside and outside the sphere can be calculated using the spherical decomposition of the general electric potential $V \propto 1/|\mathbf{r} - \mathbf{r}'|$ into Legendre Polynomials P_l [1, p. 188-190]:

$$V_\text{in}(r, \theta) = -E_\infty r \cos \theta + \sum_{l=0}^{\infty} A_l r^l P_l(\cos \theta), \quad (3.7)$$

$$V_\text{out}(r, \theta) = -E_\infty r \cos \theta + \sum_{l=0}^{\infty} \frac{B_l}{r^{l+1}} P_l(\cos \theta). \quad (3.8)$$

Applying both boundary conditions, it follows that [1, p. 249-251]

$$\begin{cases} A_l = B_l = 0 & \text{for } l \neq 1, \\ A_1 = -\frac{3}{\varepsilon_r + 2} E_\infty, \quad B_1 = \frac{\varepsilon_r - 1}{\varepsilon_r + 2} R^3 E_\infty \end{cases} \quad (3.9)$$

and the resulting homogenous electric field $\mathbf{E}_\text{in} = -\nabla V_\text{in}$ inside the sphere is given as

$$\mathbf{E}_\text{in} = \frac{3}{\varepsilon_r + 2} \mathbf{E}_\infty. \quad (3.10)$$

3 Casimir effect

The field is shown on the right in fig. 3.3. The polarizability α of the sphere can be now be determined to

$$\alpha_{\text{sphere}} = 4\pi\epsilon_0 R^3 \left(\frac{\epsilon_r - 1}{\epsilon_r + 2} \right). \quad (3.11)$$




Figure 3.3: **left:** Electric potential V of a dielectric sphere in a external electric field $\mathbf{E}_\infty \parallel \mathbf{e}_z$. **right:** The corresponding electric field lines inside and outside the dielectric sphere.

4 Plate vibrations




Figure 4.1: Vibrational modes of a spherical plate fixed at the edge with $R/d = 1000$.

Bibliography

- [1] D. J. Griffiths, *Elektrodynamik, Eine Einführung*, edited by U. Schollwöck, 4th edition (Pearson, Hallbergmoos, 2018), 1711 pages.
- [2] J. S. Pedernales and M. B. Plenio, “On the origin of force sensitivity in tests of quantum gravity with delocalised mechanical systems”, *Contemporary Physics* **64**, 147–163 (2023) 10.1080/00107514.2023.2286074, arXiv:2311.04745.
- [3] M. Plenio, “Logarithmic negativity: a full entanglement monotone that is not convex.”, *Physical Review Letters* **95**, 090503 (2005) 10.1103/PhysRevLett.95.090503, arXiv:quant-ph/0505071.
- [4] M. B. Plenio and S. Virmani, “An introduction to entanglement measures”, *Quantum Information & Computation* **7**, 1–51 (2005), arXiv:quant-ph/0504163.
- [5] M. Hartmann, “Casimir effect in the plane-spheregeometry: Beyond the proximityforce approximation”, PhD thesis (Universität Augsburg, July 2018).
- [6] A. Canaguier-Durand, R. Guérout, P. A. M. Neto, A. Lambrecht, and S. Reynaud, “The Casimir effect in the sphere-plane geometry”, *International Journal of Modern Physics Conference Series* **14**, 250–259 (2012) 10.1142/s2010194512007374, arXiv:1202.3272.
- [7] G. Vidal and R. F. Werner, “A computable measure of entanglement”, *Phys. Rev. A* **65**, 032314 (2001) 10.1103/physreva.65.032314, arXiv:quant-ph/0102117.
- [8] M. A. Nielsen and I. L. Chuang, *Quantum computation and quantum information*, 10th anniversary ed. (Cambridge University Press, Cambridge, 2010), 1676 pages.
- [9] J. S. Pedernales, G. W. Morley, and M. B. Plenio, “Motional Dynamical Decoupling for Matter-Wave Interferometry”, *Phy. Rev. Lett.* **125**, 023602 (2019) 10.1103/physrevlett.125.023602, arXiv:1906.00835.
- [10] G. A. E. Vandenbosch, “The basic concepts determining electromagnetic shielding”, *American Journal of Physics* **90**, 672–681 (2022) 10.1119/5.0087295.
- [11] D. Carney, P. C. E. Stamp, and J. M. Taylor, “Tabletop experiments for quantum gravity: a user’s manual”, *Classical and Quantum Gravity* **36**, 034001 (2018) 10.1088/1361-6382/aaf9ca, arXiv:1807.11494.
- [12] L. H. Ford, “Casimir Force between a Dielectric Sphere and a Wall: A Model for Amplification of Vacuum Fluctuations”, *Phys. Rev. A* **58**, 4279–4286 (1998) 10.1103/physreva.58.4279, arXiv:quant-ph/9804055.

Bibliography

- [13] T. W. van de Kamp, R. J. Marshman, S. Bose, and A. Mazumdar, “Quantum Gravity Witness via Entanglement of Masses: Casimir Screening”, *Phys. Rev. A* **102**, 062807 (2020) 10.1103/physreva.102.062807, arXiv:2006.06931.
- [14] E. M. Lifshitz, “The theory of molecular attractive forces between solids”, *Sov. Phys. JETP* **2**, 73–83 (1956) 10.1016/b978-0-08-036364-6.50031-4.

A Proofs and other stuff

A.1 Negativity

Lemma A.1. *The trace norm $\|A\|_1 \equiv \text{tr} \sqrt{A^\dagger A}$ of a hermitian matrix A is equal to the sum of the absolute eigenvalues of A .*

Proof. This can be immediately seen by the spectral theorem:

$$\text{tr} \sqrt{A^\dagger A} = \text{tr} \sqrt{A^2} = \text{tr} \left\{ U \sqrt{\text{diag}(\lambda_1, \dots)^2} U^\dagger \right\} = \sum_i \sqrt{\lambda_i^2} = \sum_i |\lambda_i|.$$

□

Proposition A.1. *The **negativity** $\mathcal{N}(\rho)$ of a state ρ is given as the absolute sum of all negative eigenvalues of ρ :*

$$\mathcal{N}(\rho) \equiv \frac{\|\rho^{\Gamma_A}\|_1 - 1}{2} = \left| \sum_{\lambda_i < 0} \lambda_i \right|. \quad (\text{A.1})$$

Proof. The proof is in parts given by Vidal [7]. It is known that the density matrix is hermitian: $\rho = \rho^\dagger$. Using lemma A.1, the trace norm of the density matrix is given as $\|\rho\|_1 = \sum \lambda_i = \text{tr} \rho = 1$. The partial transpose ρ^{Γ_A} obviously also satisfies $\text{tr} \rho^{\Gamma_A} = 1$ but might have negative eigenvalues. Since ρ^{Γ_A} is still hermitian, the trace norm is given by

$$\|\rho^{\Gamma_A}\|_1 = \sum_i |\lambda_i| = \sum_{\lambda_i \geq 0} \lambda_i + \sum_{\lambda_i < 0} |\lambda_i| = \sum_i \lambda_i + 2 \sum_{\lambda_i < 0} |\lambda_i| = 1 + 2 \sum_{\lambda_i < 0} |\lambda_i|,$$

where in the last step $\sum \lambda_i = \text{tr} \rho^{\Gamma_A} = 1$ was used. The negativity can be defined as $\mathcal{N}(\rho) = \left| \sum_{\lambda_i < 0} \lambda_i \right|$ and the statement is shown. □

Remark. The **logarithmic negativity** [3] relates to the negativity as follows

$$E_N(\rho) = \log_2 \|\rho^{\Gamma_A}\|_1 = \log_2 (2\mathcal{N}(\rho) + 1) \quad (\text{A.2})$$

and can therefore be easily calculated by using the above proposition A.1. In comparison to the negativity, logarithmic negativity has additive properties [4]:

$$E_N(\rho \otimes \sigma) = E_N(\rho) + E_N(\sigma)$$

A.2 Fidelity

The *fidelity* of two quantum states ρ and σ is defined as [8, p. 409-412]

$$F(\rho, \sigma) = \text{tr} \sqrt{\sqrt{\rho} \sigma \sqrt{\rho}} \quad (\text{A.3})$$

and can be used as a distance measurement between quantum states. It is monotonic, concave and bounded between 0 and 1. If both states are equal $\rho = \sigma$, it is clear that $F(\rho, \sigma) = 1$, by using $\sqrt{\rho} \rho \sqrt{\rho} = \rho^2$. If both states commute, i.e. they are diagonalizable in the same orthogonal basis $\{|i\rangle\}$,

$$\rho = \sum_i r_i |i\rangle\langle i|; \quad \sigma = \sum_i s_i |i\rangle\langle i|,$$

the fidelity is given as [8]

$$F(\rho, \sigma) = \text{tr} \sqrt{\sum_i r_i s_i |i\rangle\langle i|} = \sum_i \sqrt{r_i s_i}.$$

This can be seen immediately by the use of the spectral theorem $\text{tr} \sqrt{\rho} = \text{tr} \{U \sqrt{\text{diag}(r_i)} U^\dagger\} = \text{tr} \text{diag}(\sqrt{r_i})$. Another special case is given for the fidelity of a pure state $\rho = |\psi\rangle\langle\psi|$ and an arbitrary state σ [8]:

$$F(|\psi\rangle, \sigma) = \text{tr} \sqrt{|\psi\rangle\langle\psi| \sigma |\psi\rangle\langle\psi|} = \sqrt{\langle\psi| \sigma |\psi\rangle}.$$

If the state $\sigma = |\phi\rangle\langle\phi|$ is also pure, the fidelity reduces to

$$F(|\psi\rangle, |\phi\rangle) = |\langle\psi|\phi\rangle| \leq 1, \quad (\text{A.4})$$

with equality being attained if the states are the same and only differ by a phase.