



ulm university universität  
**uulm**

**Fakultät für  
Naturwissenschaften**

Institute of Theoretical  
Physics

**Title of the work**

**Bachelor Thesis**

**Submitted by:**

Jan Bulling  
jan.bulling@uni-ulm.de  
1109395

**Supervised by:**

Marit O. E. Steiner, Julen S. Pedernales, Martin B. Plenio

## Abstract

# Contents

<b>1</b>	<b>Introduction</b>	<b>4</b>
1.1	Feynman's Gedankenexperiment . . . . .	4
<b>2</b>	<b>A first look at the problem</b>	<b>5</b>
2.1	Entanglement measures . . . . .	5
<b>3</b>	<b>Casimir effect</b>	<b>7</b>
3.1	Proximity force approximation . . . . .	7
3.2	Imperfect plate and spheres . . . . .	8
3.3	Casimir forces between a conducting plate and a dielectric sphere . . . . .	8
3.3.1	Polarizability of a dielectric sphere . . . . .	8
<b>4</b>	<b>Plate vibrations</b>	<b>10</b>
	<b>Bibliography</b>	<b>11</b>
<b>A</b>	<b>Proofs and other stuff</b>	<b>13</b>
A.1	Negativity . . . . .	13
A.2	Fidelity . . . . .	14


# **1 Introduction**

## **1.1 Feynman's Gedankenexperiment**

## 2 A first look at the problem

Problem, general calculations from Julens paper without casimir

When is the system entangled (distance-measures, fidelity ??)



**Figure 2.1:** Simple Problem from Julens paper

Ignoring (for now) all other forms of interaction between the masses except gravity

The density  $\rho(t) = |\psi(t)\rangle\langle\psi(t)|$  can be represented in the basis  $\{|\psi_A^\uparrow\rangle, |\psi_A^\downarrow\rangle\} \otimes \{|\psi_B^\uparrow\rangle, |\psi_B^\downarrow\rangle\}$  as

$$\rho(t) = \frac{1}{4} \begin{pmatrix} 1 & & & 1 \\ & 1 & & \\ & & 1 & \\ 1 & & & 1 \end{pmatrix} \quad (2.1)$$

### 2.1 Entanglement measures

Why are they needed, what can one do?

Logarithmic negativity, properties, calculation





Figure 2.2: My problem



**Figure 2.3:** Entanglement dynamics quantified by the logarithmic negativity for two different orientations of the spatial superpositions relative to each other. The time of maximum entanglement  $t_{\max}$  for the parallel configuration is given by  $t_{\max} = 8\pi\hbar L^3/(GM_A M_B d^2) \simeq 258$  ms.


## 3 Casimir effect

General introduction and comparison with retarded van der Waals forces

$$F_{\text{Casimir}} = -\frac{\hbar c \pi^2}{240 L^4} A \quad (3.1)$$

$$V_{\text{Casimir}} = \frac{\hbar c \pi^2}{720 L^3} A \quad (3.2)$$

### 3.1 Proximity force approximation



**Figure 3.1:** In the proximity force approximation the sphere is divided into infinitesimal plane areas  $dA$  which all exert a force  $dF$  according to eq. (3.1). All the contributions are added up together.

## 3.2 Imperfect plate and spheres

Python numerical approach, gaussian modes (vibration modes of a spherical plane), perlin noise

## 3.3 Casimir forces between a conducting plate and a dielectric sphere

### 3.3.1 Polarizability of a dielectric sphere

The polarizability  $\alpha$  is defined via

$$\mathbf{E}_\infty \alpha = \mathbf{p}, \quad (3.3)$$

where  $\mathbf{p}$  is the induced dipole moment and  $\mathbf{E}_\infty$  is the external electric field that induces the dipole moment. For a linear and uniform dielectric, it is given as  $\mathbf{p} = \mathcal{V} \varepsilon_0 (\varepsilon_r - 1) \mathbf{E}_\text{in}$  [1, p. 220-226]. Here,  $\mathcal{V}$  is the volume of the object and  $\mathbf{E}_\text{in}$  is the electric field inside the dielectric. The electrostatic boundary conditions for the problem are given by

$$V_\text{in}|_{r=R} = V_\text{out}|_{r=R} \quad \text{and} \quad \varepsilon_r \varepsilon_0 \frac{\partial V_\text{in}}{\partial r} \Big|_{r=R} = \varepsilon_0 \frac{\partial V_\text{out}}{\partial r} \Big|_{r=R} \quad (3.4)$$

and the electric potential outside of the sphere at  $r \rightarrow \infty$  should be equal to the external dipole-inducing field  $V_\text{out}|_{r \rightarrow \infty} = -\mathbf{E}_\infty \cdot \mathbf{r} = -E_\infty r \cos \theta$ . The electric potential inside and outside the sphere can be calculated using the spherical decomposition of the general electric potential  $V \propto 1/|\mathbf{r} - \mathbf{r}'|$  into Legendre Polynomials  $P_l$  [1, p. 188-190]:

$$V_\text{in}(r, \theta) = -E_\infty r \cos \theta + \sum_{l=0}^{\infty} A_l r^l P_l(\cos \theta), \quad (3.5)$$

$$V_\text{out}(r, \theta) = -E_\infty r \cos \theta + \sum_{l=0}^{\infty} \frac{B_l}{r^{l+1}} P_l(\cos \theta). \quad (3.6)$$

Applying both boundary conditions, it follows that [1, p. 249-251]

$$\begin{cases} A_l = B_l = 0 & \text{for } l \neq 1, \\ A_1 = -\frac{3}{\varepsilon_r + 2} E_\infty, \quad B_1 = \frac{\varepsilon_r - 1}{\varepsilon_r + 2} R^3 E_\infty \end{cases} \quad (3.7)$$


and the resulting homogenous electric field  $\mathbf{E}_\text{in} = -\nabla V_\text{in}$  inside the sphere is given as

$$\mathbf{E}_\text{in} = \frac{3}{\varepsilon_r + 2} \mathbf{E}_\infty. \quad (3.8)$$

The field is shown on the right in fig. 3.2. The polarizability  $\alpha$  of the sphere can be now be determined to


$$\alpha_\text{sphere} = 4\pi \varepsilon_0 R^3 \left( \frac{\varepsilon_r - 1}{\varepsilon_r + 2} \right). \quad (3.9)$$





**Figure 3.2:** **left:** Electric potential  $V$  of a dielectric sphere in a external electric field  $\mathbf{E}_\infty \parallel \mathbf{e}_z$ . **right:** The corresponding electric field lines inside and outside the dielectric sphere.

## 4 Plate vibrations



**Figure 4.1:** Vibrational modes of a spherical plate fixed at the edge with  $R/d = 1000$ .

# Bibliography

- [1] D. J. Griffiths, *Elektrodynamik, Eine Einführung*, edited by U. Schollwöck, 4th edition (Pearson, Hallbergmoos, 2018), 1711 pages.
- [2] J. S. Pedernales and M. B. Plenio, “On the origin of force sensitivity in tests of quantum gravity with delocalised mechanical systems”, *Contemporary Physics* **64**, 147–163 (2023) 10.1080/00107514.2023.2286074, arXiv:2311.04745.
- [3] M. Plenio, “Logarithmic negativity: a full entanglement monotone that is not convex.”, *Physical Review Letters* **95**, 090503 (2005) 10.1103/PhysRevLett.95.090503, arXiv:quant-ph/0505071.
- [4] M. B. Plenio and S. Virmani, “An introduction to entanglement measures”, *Quantum Information & Computation* **7**, 1–51 (2005), arXiv:quant-ph/0504163.
- [5] M. Hartmann, “Casimir effect in the plane-spheregeometry: Beyond the proximityforce approximation”, PhD thesis (Universität Augsburg, July 2018).
- [6] A. Canaguier-Durand, R. Guérout, P. A. M. Neto, A. Lambrecht, and S. Reynaud, “The Casimir effect in the sphere-plane geometry”, *International Journal of Modern Physics Conference Series* **14**, 250–259 (2012) 10.1142/s2010194512007374, arXiv:1202.3272.
- [7] G. Vidal and R. F. Werner, “A computable measure of entanglement”, *Phys. Rev. A* **65**, 032314 (2001) 10.1103/physreva.65.032314, arXiv:quant-ph/0102117.
- [8] M. A. Nielsen and I. L. Chuang, *Quantum computation and quantum information*, 10th anniversary ed. (Cambridge University Press, Cambridge, 2010), 1676 pages.
- [9] J. S. Pedernales, G. W. Morley, and M. B. Plenio, “Motional Dynamical Decoupling for Matter-Wave Interferometry”, *Phy. Rev. Lett.* **125**, 023602 (2019) 10.1103/physrevlett.125.023602, arXiv:1906.00835.
- [10] G. A. E. Vandenbosch, “The basic concepts determining electromagnetic shielding”, *American Journal of Physics* **90**, 672–681 (2022) 10.1119/5.0087295.
- [11] D. Carney, P. C. E. Stamp, and J. M. Taylor, “Tabletop experiments for quantum gravity: a user’s manual”, *Classical and Quantum Gravity* **36**, 034001 (2018) 10.1088/1361-6382/aaf9ca, arXiv:1807.11494.
- [12] L. H. Ford, “Casimir Force between a Dielectric Sphere and a Wall: A Model for Amplification of Vacuum Fluctuations”, *Phys. Rev. A* **58**, 4279–4286 (1998) 10.1103/physreva.58.4279, arXiv:quant-ph/9804055.

## *Bibliography*

- [13] T. W. van de Kamp, R. J. Marshman, S. Bose, and A. Mazumdar, “Quantum Gravity Witness via Entanglement of Masses: Casimir Screening”, *Phys. Rev. A* **102**, 062807 (2020) 10.1103/physreva.102.062807, arXiv:2006.06931.
- [14] E. M. Lifshitz, “The theory of molecular attractive forces between solids”, *Sov. Phys. JETP* **2**, 73–83 (1956) 10.1016/b978-0-08-036364-6.50031-4.

# A Proofs and other stuff

## A.1 Negativity

**Lemma A.1.** *The trace norm  $\|A\|_1 \equiv \text{tr} \sqrt{A^\dagger A}$  of a hermitian matrix  $A$  is equal to the sum of the absolute eigenvalues of  $A$ .*

*Proof.* This can be immediately seen by the spectral theorem:

$$\text{tr} \sqrt{A^\dagger A} = \text{tr} \sqrt{A^2} = \text{tr} \left\{ U \sqrt{\text{diag}(\lambda_1, \dots)^2} U^\dagger \right\} = \sum_i \sqrt{\lambda_i^2} = \sum_i |\lambda_i|.$$

□

**Proposition A.1.** *The **negativity**  $\mathcal{N}(\rho)$  of a state  $\rho$  is given as the absolute sum of all negative eigenvalues of  $\rho$ :*

$$\mathcal{N}(\rho) \equiv \frac{\|\rho^{\Gamma_A}\|_1 - 1}{2} = \left| \sum_{\lambda_i < 0} \lambda_i \right|. \quad (\text{A.1})$$

*Proof.* The proof is in parts given by Vidal [7]. It is known that the density matrix is hermitian:  $\rho = \rho^\dagger$ . Using lemma A.1, the trace norm of the density matrix is given as  $\|\rho\|_1 = \sum \lambda_i = \text{tr} \rho = 1$ . The partial transpose  $\rho^{\Gamma_A}$  obviously also satisfies  $\text{tr} \rho^{\Gamma_A} = 1$  but might have negative eigenvalues. Since  $\rho^{\Gamma_A}$  is still hermitian, the trace norm is given by

$$\|\rho^{\Gamma_A}\|_1 = \sum_i |\lambda_i| = \sum_{\lambda_i \geq 0} \lambda_i + \sum_{\lambda_i < 0} |\lambda_i| = \sum_i \lambda_i + 2 \sum_{\lambda_i < 0} |\lambda_i| = 1 + 2 \sum_{\lambda_i < 0} |\lambda_i|,$$

where in the last step  $\sum \lambda_i = \text{tr} \rho^{\Gamma_A} = 1$  was used. The negativity can be defined as  $\mathcal{N}(\rho) = \left| \sum_{\lambda_i < 0} \lambda_i \right|$  and the statement is shown. □

*Remark.* The **logarithmic negativity** [3] relates to the negativity as follows

$$E_N(\rho) = \log_2 \|\rho^{\Gamma_A}\|_1 = \log_2 (2\mathcal{N}(\rho) + 1) \quad (\text{A.2})$$

and can therefore be easily calculated by using the above proposition A.1. In comparison to the negativity, logarithmic negativity has additive properties [4]:

$$E_N(\rho \otimes \sigma) = E_N(\rho) + E_N(\sigma)$$

## A.2 Fidelity

The *fidelity* of two quantum states  $\rho$  and  $\sigma$  is defined as [8, p. 409-412]

$$F(\rho, \sigma) = \text{tr} \sqrt{\sqrt{\rho} \sigma \sqrt{\rho}} \quad (\text{A.3})$$

and can be used as a distance measurement between quantum states. It is monotonic, concave and bounded between 0 and 1. If both states are equal  $\rho = \sigma$ , it is clear that  $F(\rho, \sigma) = 1$ , by using  $\sqrt{\rho} \rho \sqrt{\rho} = \rho^2$ . If both states commute, i.e. they are diagonalizable in the same orthogonal basis  $\{|i\rangle\}$ ,

$$\rho = \sum_i r_i |i\rangle\langle i|; \quad \sigma = \sum_i s_i |i\rangle\langle i|,$$

the fidelity is given as [8]

$$F(\rho, \sigma) = \text{tr} \sqrt{\sum_i r_i s_i |i\rangle\langle i|} = \sum_i \sqrt{r_i s_i}.$$

This can be seen immediately by the use of the spectral theorem  $\text{tr} \sqrt{\rho} = \text{tr} \{U \sqrt{\text{diag}(r_i)} U^\dagger\} = \text{tr} \text{diag}(\sqrt{r_i})$ . Another special case is given for the fidelity of a pure state  $\rho = |\psi\rangle\langle\psi|$  and an arbitrary state  $\sigma$  [8]:

$$F(|\psi\rangle, \sigma) = \text{tr} \sqrt{|\psi\rangle\langle\psi| \sigma |\psi\rangle\langle\psi|} = \sqrt{\langle\psi| \sigma |\psi\rangle}.$$

If the state  $\sigma = |\phi\rangle\langle\phi|$  is also pure, the fidelity reduces to

$$F(|\psi\rangle, |\phi\rangle) = |\langle\psi|\phi\rangle| \leq 1, \quad (\text{A.4})$$

with equality being attained if the states are the same and only differ by a phase.