

Worksheet-1 in R

Worksheet for R Programming

Instructions:

- Use RStudio or the RStudio Cloud accomplish this worksheet. + Save the R script as *RWorksheet_lastname#1.R*.
- Create your own *GitHub repository* and push the R script as well as this pdf worksheet to your own repo.

Accomplish this worksheet by answering the questions being asked and writing the code manually.

Using functions:

`seq()`, `assign()`, `min()`, `max()`, `c()`, `sort()`, `sum()`, `filter()`

1. Set up a vector named `age`, consisting of 34, 28, 22, 36, 27, 18, 52, 39, 42, 29, 35, 31, 27, 22, 37, 34, 19, 20, 57, 49, 50, 37, 46, 25, 17, 37, 42, 53, 41, 51, 35, 24, 33, 41.

a. How many data points?

- **There are 35 data points**

b. Write the R code and its output.

```
[1] 34 28 22 36 27 18 52 39 42 29 35 31 27 22 37 34 19 20 57 49 50 37 46 25 17  
37 42 53 41
```

2. Find the reciprocal of the values for `age`.

Write the R code and its output.

```
> exe_age <- 1/age
```

```
> exe_age
```

```
[1] 0.02941176 0.03571429 0.04545455 0.02777778 0.03703704 0.05555556 0.01923077  
0.02564103 0.02380952 0.03448276  
[11] 0.02857143 0.03225806 0.03703704 0.04545455 0.02702703 0.02941176 0.05263158
```

0.05000000 0.01754386 0.02040816

[21] 0.02000000 0.02702703 0.02173913 0.04000000 0.05882353 0.02702703 0.02380952

0.01886792 0.02439024 0.01960784

[31] 0.02857143 0.04166667 0.03030303 0.02439024

3. Assign also `new_age <- c(age, 0, age)`.

What happen to the `new_age`?

```
[1] 34 28 22 36 27 18 52 39 42 29 35 31 27 22 37 34 19 20 57 49 50 37 46 25 17 37 42 53 41 51 35 24 33
41 0 34 28 22 36
[40] 27 18 52 39 42 29 35 31 27 22 37 34 19 20 57 49 50 37 46 25 17 37 42 53 41 51 35 24 33 41
```

4. Sort the values for `age`.

Write the R code and its output.

```
sort(age)
[1] 17 18 19 20 22 22 24 25 27 27 28 29 31 33 34 34 35 35 36 37 37 37 39 41 41 42 42 46 49 50 51 52 53
57
```

5. Find the minimum and maximum value for `age`.

Write the R code and its output.

```
> min(age)
[1] 17
> max(age)
[1] 57
```

6. Set up a vector named `data`, consisting of 2.4, 2.8, 2.1, 2.5, 2.4, 2.2, 2.5, 2.3, 2.5, 2.3, 2.4, and 2.7.

a. How many data points?

- **12 Data points**

b. Write the R code and its output.

```
> data <- c(2.4, 2.8, 2.1, 2.5, 2.4, 2.2, 2.5, 2.3, 2.5, 2.3, 2.4, 2.7)
> data
[1] 2.4 2.8 2.1 2.5 2.4 2.2 2.5 2.3 2.5 2.3 2.4 2.7
```

7. Generates a new vector for data where you double every value of the data. | What happen to the data?

```
> data * 2  
[1] 4.8 5.6 4.2 5.0 4.8 4.4 5.0 4.6 5.0 4.6 4.8 5.4
```

8. Generate a sequence for the following scenario:

8.1 Integers from 1 to 100.

```
> seq(1:100)
```

```
[1] 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29
[30] 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56
57 58
[59] 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85
86 87
[88] 88 89 90 91 92 93 94 95 96 97 98 99 100
```

8.2 Numbers from 20 to 60

```
> seq(20,60)
```

```
[1] 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47
48 49 50 51 52 53 54 55 56 57 58
[40] 59 60
```

*8.3 Mean of numbers from 20 to 60

```
> mean(20:60)
```

```
[1] 40
```

*8.4 Sum of numbers from 51 to 91

```
> sum(51:91)
```

```
[1] 2911
```

*8.5 Integers from 1 to 1,000

```
> seq(1:1000)
```

```
[1] 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23
[24] 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46
[47] 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69
[70] 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92
[93] 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113
114 115
[116] 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135
136 137 138
[139] 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158
159 160 161
[162] 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181
182 183 184
```

[185] 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204
 205 206 207
 [208] 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227
 228 229 230
 [231] 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250
 251 252 253
 [254] 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273
 274 275 276
 [277] 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296
 297 298 299
 [300] 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319
 320 321 322
 [323] 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342
 343 344 345
 [346] 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365
 366 367 368
 [369] 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388
 389 390 391
 [392] 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411
 412 413 414
 [415] 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434
 435 436 437
 [438] 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457
 458 459 460
 [461] 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480
 481 482 483
 [484] 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503
 504 505 506
 [507] 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526
 527 528 529
 [530] 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549
 550 551 552
 [553] 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572
 573 574 575
 [576] 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595
 596 597 598
 [599] 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618
 619 620 621
 [622] 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641
 642 643 644
 [645] 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664
 665 666 667
 [668] 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687
 688 689 690

[691] 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710
 711 712 713
 [714] 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733
 734 735 736
 [737] 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756
 757 758 759
 [760] 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779
 780 781 782
 [783] 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802
 803 804 805
 [806] 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825
 826 827 828
 [829] 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848
 849 850 851
 [852] 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871
 872 873 874
 [875] 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894
 895 896 897
 [898] 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917
 918 919 920
 [921] 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940
 941 942 943
 [944] 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963
 964 965 966
 [967] 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986
 987 988 989
 [990] 990 991 992 993 994 995 996 997 998 999 1000

a. How many data points from 8.1 to 8.4? _____

8.1 = 100 data points

8.2 = 41 data points

8.3 = 1 data point

8.4 = 1 data point

TOTAL OF 143 DATA POINT

b. Write the R code and its output from 8.1 to 8.4.

> data.frame(1:100)

X1.100

1 1

2 2

3 3

4 4

5	5
6	6
7	7
8	8
9	9
10	10
11	11
12	12
13	13
14	14
15	15
16	16
17	17
18	18
19	19
20	20
21	21
22	22
23	23
24	24
25	25
26	26
27	27
28	28
29	29
30	30
31	31
32	32
33	33
34	34
35	35
36	36
37	37
38	38
39	39
40	40
41	41
42	42
43	43
44	44
45	45
46	46
47	47
48	48

49	49
50	50
51	51
52	52
53	53
54	54
55	55
56	56
57	57
58	58
59	59
60	60
61	61
62	62
63	63
64	64
65	65
66	66
67	67
68	68
69	69
70	70
71	71
72	72
73	73
74	74
75	75
76	76
77	77
78	78
79	79
80	80
81	81
82	82
83	83
84	84
85	85
86	86
87	87
88	88
89	89
90	90
91	91
92	92

93	93
94	94
95	95
96	96
97	97
98	98
99	99
100	100

c. For 8.5 find only maximum data points until 10.

```
> data.frame(20:60)
```

	X20.60
1	20
2	21
3	22
4	23
5	24
6	25
7	26
8	27
9	28
10	29
11	30
12	31
13	32
14	33
15	34
16	35
17	36
18	37
19	38
20	39
21	40
22	41
23	42
24	43
25	44
26	45
27	46

28	47
29	48
30	49
31	50
32	51
33	52
34	53
35	54
36	55
37	56
38	57
39	58
40	59
41	60

9. *Print a vector with the integers between 1 and 100 that are not divisible by 3, 5 and 7 using filter option.

```
filter(function(i) { all(i %% c(3,5,7) != 0) }, seq(100))
```

Write the R code and its output.

```
> filter(function(i){all(i%%c(3, 5, 7) != 0)}, seq(100))
```

```
Error in attr("data, "tsp") <- c(start, end, frequency) :
```

```
object is not a matrix
```

10. Generate a sequence backwards of the integers from 1 to 100.

Write the R code and its output.

```
> seq(100:1)
```

```
[1] 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29
[30] 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56
57 58
[59] 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85
86 87
[88] 88 89 90 91 92 93 94 95 96 97 98 99 100
```

11. List all the natural numbers below 25 that are multiples of 3 or 5.

Find the sum of these multiples.

```
> sum((1:25)[((1:25)%%3==0) | ((1:25)%%5==0)])
```

```
[1] 168
```

a. How many data points from 10 to 11?

101 data points

b. Write the R code and its output from 10 and 11.

```
> data.frame(1:100)
```

```
  X1.100
```

```
1      1
2      2
3      3
```

4	4
5	5
6	6
7	7
8	8
9	9
10	10
11	11
12	12
13	13
14	14
15	15
16	16
17	17
18	18
19	19
20	20
21	21
22	22
23	23
24	24
25	25
26	26
27	27
28	28
29	29
30	30
31	31
32	32
33	33
34	34
35	35
36	36
37	37
38	38
39	39
40	40
41	41
42	42
43	43
44	44
45	45
46	46
47	47

48	48
49	49
50	50
51	51
52	52
53	53
54	54
55	55
56	56
57	57
58	58
59	59
60	60
61	61
62	62
63	63
64	64
65	65
66	66
67	67
68	68
69	69
70	70
71	71
72	72
73	73
74	74
75	75
76	76
77	77
78	78
79	79
80	80
81	81
82	82
83	83
84	84
85	85
86	86
87	87
88	88
89	89
90	90
91	91

92 92
93 93
94 94
95 95
96 96
97 97
98 98
99 99
100 100

12. Statements can be grouped together using braces '{' and '}'. A group of statements is sometimes called a **block**. Single statements are evaluated when a new line is typed at the end of the syntactically complete statement. Blocks are not evaluated until a new line is entered after the closing brace.

Enter this statement:

```
{ x <- 0+ x + 5 + }
```

Describe the output.

```
> { x <- 0+ x + 5 + }
```

```
Error: unexpected '}' in "{ x <- 0+ x + 5 + }"
```

13. *Set up a vector named score, consisting of 72, 86, 92, 63, 88, 89, 91, 92, 75, 75 and 77. To access individual elements of an atomic vector, one generally uses the x[i] construction.

Find x[2] and x[3]. Write the R code and its output.

```
2 86
3 92
```

14. *Create a vector a = c(1,2,NA,4,NA,6,7).

a. Change the NA to 999 using the codes print(a,na.print="-999").

```
> a = c(1, 2, NA, 4, NA, 6, 7)
> print(a, na.print="-999")
[1] 1 2 -999 4 -999 6 7
```

b. Write the R code and its output. Describe the output.

```
> a = c(1, 2, NA, 4, NA, 6, 7)
> print(a, na.print="-999")
[1] 1 2 -999 4 -999 6 7
```

The NA was replaced by -999

15. A special type of function calls can appear on the left hand side of the assignment operator as in > class(x) <- "foo".

Follow the codes below:

```
name = readline(prompt="Input your name: ")
age = readline(prompt="Input your age: ")
print(paste("My name is",name, "and I am",age,"years old. "))
print(R.version.string)
```

What is the output of the above code?

```
[1] "My name is Jan James Lennox Tahumar and I am 20 years old"
> print(R.version.string)
[1] "R version 4.2.1 (2022-06-23 ucrt)"
```