

UNIVERSITÄT BONN

Notes for the lecture

Algebraic Topology I

held by

Stefan Schwede

T_EXed by

Jan Malmström

WiSe 2025/26

Corrections and improvements

If you have corrections or improvements, contact me via (s94jmalm@uni-bonn.de).

Contents

Lecture	1
1.1 Blakers-Massay theorem/Homotopy excision	3
1.2 Feudenthal suspension theorem	8
1.3 Hurewicz-theorem	10
1.3.1 Relative Hurewicz theorem	16
1.4 CW-Approximation	18
1.5 Representativity of cohomology	22
1.6 Cohomology operations and the Steenrod algebra	26
 Appendix	 A
List of definitions	B
List of statements	C

Lecture

Organizatorial

For this term we will be doing unstable homotopy theory. Next term we will be doing stable homotopy theory. Note that there were 2 previous courses. Note that all important information is shared on the website <https://www.math.uni-bonn.de/people/schwede/at1-ws2526>. You can sign up for the previous topology courses and see the lecture videos for these courses there.

There are no lecture notes for this lecture specifically, but some similar materials are linked on the webpage.

Exercise sheets will be uploaded fridays and handed in 11 days later via eCampus. Registration for eCampus opens at 4 today.

For exam admission you will have to score 50% of the points on the exercise sheets and have presented 2 exercises in tutorial.

The first exam will be written in the last week of semester.

I fear I will not be able to copy pictures here.

1.1 Blakiers-Massy theorem/Homotopy excision

We start with a reminder on relative homotopy groups.

Definition 1.1: Relative Homotopy Groups

Let (X, A) be a space pair i.e. A is a subspace of a topological space X . We write

$$I = [0, 1] \quad I^n = [0, 1]^n \text{ the } n\text{-cube}$$

$$\partial(I^n) = \text{boundary of } I^n$$

$$I^{n-1} \subseteq I^n$$

via Inclusion on the first $n - 1$ coordinates.

$$J^{n-1} = I^{n-1} \times \{1\} \cup (\partial I^{n-1}) \times [0, 1]$$

He draws a picture for $n = 2$.

For $n \geq 1$ the n -th relative homotopy groups $\pi_n(X, A, x)$ is the set of triple homotopy classes of triple maps $x \in A \subseteq X$

$$(I^n, \partial I^n, J^{n-1}) \rightarrow (X, A, \{x\})$$

where a triple map takes each subset on the left into the subset on the right. A triple-homotopy must also conserve these conditions.

For $n \geq 2$ or $n = 1$ and $A = \{x\}$ the set $\pi_n(X, A, \{x\})$ has a group structure by concatenation in the first coordinate. He again draws a picture.

The group structure is commutative if $n \geq 3$ or $n = 2$ and $A = \{x\}$.

Definition 1.2: n -Connectedness

Let $n \geq 0$. A space pair (X, A) is n -connected, if the following equivalent conditions hold:

1. For all $0 \leq q \leq n$ every pair map $(I^q, \partial I^q) \rightarrow (X, A)$ is homotpic relative $\partial(I^q)$ to a map with image in A
2. For all $a \in A$, $\text{incl}_*: \pi_q(A, a) \rightarrow \pi_q(X, a)$ is bijective for $q \leq n$ and surjective for $q = n$.
3. $\pi_0(A) \rightarrow \pi_0(X)$ is bijective and for all $1 \leq q \leq n$ the relative homotopy group

$$\pi_q(X, A, x) \cong 0$$

Proof. You proof the equivalence using the LES¹ of homotopy groups. □

Let Y be a space, Y_1, Y_2 open subsets of $Y = Y_1 \cup Y_2$, $Y_0 := Y_1 \cap Y_2$.

Excision in homology shows that for all abelian groups B , $i \geq 0$

$$H_i(Y_2; Y_0, B) \rightarrow H_i(Y, Y_1; B)$$

is an isomorphism.

Excision does not generally hold for homotopy groups, i.e. for $x \in Y_0$

$$\text{incl}_*: \pi_i(Y_2, Y_0; x) \rightarrow \pi_i(Y, Y_1; x)$$

¹Long exact sequence

is **not** generally an isomorphism.

"Blakers Massey theorem implies that excision holds for homotopy groups in a range."

Theorem 1.3: Blakers Massey

Let Y be a space, Y_1, Y_2 open subsets with $Y = Y_1 \cup Y_2, Y_0 := Y_1 \cap Y_2$. Let $p, q \geq 0$, such that for all $y \in Y_0$

$$\pi_i(Y_1, Y_0, y) = 0 \text{ for all } 1 \leq i \leq p$$

and

$$\pi_i(Y_2, Y_0, y) = 0 \text{ for all } 1 \leq i \leq q$$

Then for all $y \in Y_0$, the map

$$\text{incl}_*: \pi_i(Y_2, Y_0, y) \rightarrow \pi_i(Y, Y_1, y)$$

is an isomorphism for $1 \leq i < p + q$ and surjective for $i = p + q$. He notes how the referenced literature uses different indices. They have proofs in more detail and pictures, however Lück's script contains typos

Proof. Schwede explains he doesn't like the proof, it is too technical and not very enlightening.

We define what cubes are

Cubes in \mathbb{R}^n , $n \geq 1$. $a = (a_1, \dots, a_n) \in \mathbb{R}^n$ the "lower left corner of the cube"

$\partial \in \mathbb{R}_{\geq 0}$ "side length of the cube"

$L \subset \{1, \dots, n\}$ "relevant dimensions"

$W = W(a, \delta, L) = \{x = (x_1, \dots, x_n) \in \mathbb{R}^n : a_i \leq x_i \leq a_i + \delta \text{ for all } i \in L, x_i = a_i \text{ for all } i \in \{1, \dots, n\} \setminus L\}$

² A face W' of W is a subset of the form

$$W' = \{x \in W : x_i = a_i \text{ for all } i \in L_0, x_i = a_i + \delta \text{ for all } i \in L_1\}$$

for some subsets $L_0, L_1 \subseteq L$

Let $1 \leq p \leq n$ we define two subsets of a cube $W = (a, \delta, L)$.

$$K_p(W) = \{x \in W : x_i < a_i + \delta/2 \text{ for at least } p \text{ values of } i \text{ in } L\}$$

We call these " p small coordinates"

$$G_p(W) = \{x \in W : x_i > a_i + \delta/2 \text{ for at least } p \text{ coordinates } i \text{ in } L\}$$

these are " p big coordinates".

For $p > \dim(W)$, $K_p(W) = G_p(W) = \emptyset$. If $p + q \geq \dim(W)$, then $K_p(W) \cap G_q(W) = \emptyset$.

He draws pictures.

²W weil Würfel

Lemma 1.4: 1.14

It is Lemma 1.14 in Lück's Script.

Let (Y, A) be a space pair, $W \subseteq \mathbb{R}^n$ a cube, $f: W \rightarrow Y$ continuous. Suppose that for some $p \leq \dim(W)$, $f^{-1}(A) \cap W' \subseteq K_p(W')$ for all proper¹ faces W' of W .

Then there is a continuous map $g: W \rightarrow Y$ homotopic to f relative ∂W such that all $g^{-1}(A) \subseteq K_p(W)$

¹subcube of the boundary

Proof. Wlog: $W = I^n = W(0, 1, \{1, \dots, n\})$

Let I_2^n be the subcube $[0, 1/2]^n$. He draws a picture. $x_4 = (1/4, \dots, 1/4) \in I_2^n$. We define a continuous map $h: I^n \rightarrow I^n$ by radical projection away from x_4 . Picture. Let $r(y)$ be the ray from x to y . We map all of $r(y) \cap I^n \setminus I_2^n$ to the intersection point of $r(y)$ and ∂I^n and the rest linearly extends as far as required.³

Obviously⁴ h is homotopic relative boundary ∂I^n to the identity.

We set $g: f \circ h: I^n \rightarrow Y$, which is then homotopic relative $\partial(I^n)$ ⁵ to f .

It remains to show that $g^{-1}(A) \subseteq K_p(W)$. Consider $z \in I^n$ with $g(z) \in A$.

Case 1 for all $i = 1, \dots, n, z_i < 1/2$, i.e. $z \in I_2^n$, then $z \in K_n(I^n) \subseteq K_p(I^n)$.

Case 2 There is an $i \in \{1, \dots, n\}$, s.t. $z \geq 1/2$. Then $h(z) \in \partial(I^n)$. Let W' be some proper face of W , with $h(z) \in W'$. Since $f(h(z)) = g(z) \in A$, by hypothesis, $h(z) \in K_p(W')$, so $h(z) < 1/2$ for at least p coordinates. By expansion⁶ property of h , also p coordinates of z are small coordinates.

□

Proposition 1.5. ⁷ Let Y_1, Y_2 be open subsets of $Y, Y_0 := Y_1 \cap Y_2$. Suppose that (Y_1, Y_0) is p -connected, (Y_2, Y_0) is q -connected. Let $f: I^n \rightarrow Y$ be continuous. Let $\mathcal{W} = \{W\}$ be a subdivision of I^n into subcubes of the same side length s.t. for all $W \in \mathcal{W}$ $f(W) \subseteq Y_1$ or $f(W) \subseteq Y_2$. Then there is a homotopy $h: I^n \times I \rightarrow Y$ with $h_0 = f$ such that for all $W \in \mathcal{W}$:

1. If $f(W) \subseteq Y_j, j \in 0, 1, 2$, then $h_t(W) \subseteq Y_j$ for all $t \in [0, 1]$
2. If $f(W) \subseteq Y_0$, then $h_t|_W = f|_W$, i.e. h is constant on W .
3. If $f(W) \subseteq Y_1$, then $h_1^{-1}(Y_1 \setminus Y_0) \subseteq K_{p+1}(W)$.
4. If $f(W) \subseteq Y_2$, then $h_1^{-1}(Y_2 \setminus Y_0) \subseteq G_{q+1}(W)$.

Proof. We let $C^k \subseteq I^n$ be the union of all cubes in \mathcal{W} of dimension at most k . We construct homotopies $h[k]: C_k \times I \rightarrow Y$, such that for all $W \in \mathcal{W}, W \subseteq C_k$ conditions 1. to 4. hold, and $h[k]$ is constant on $C_{k-1} \times I$. Then the final $h[n]$ does the job.

Note. If $W \in \mathcal{W}$ and $f(W) \subseteq Y_0$ and 2. holds, then also 3. and 4. hold.

$$h^{-1}(Y_1 \setminus Y_0) = h_1^{-1}(Y_2 \setminus Y_0) = \emptyset$$

If $W \in \mathcal{W}$, is such that $f(W) \subseteq Y_1$ and $f(W) \subseteq Y_2$, then $f(W) \subseteq Y_1 \cap Y_2 = Y_0$. So each $W \in \mathcal{W}$ is in exactly one of the following cases

- $f(W) \subseteq Y_0$

³I hope this description is clear, hard without the picture.

⁴Meaning he's too lazy to come up with formulas for the map.

⁵I am very inconsistent in remembering these parantheses with the boundary operator. Just imagine it always being as here.

⁶no idea if this is the word he wrote

⁷11.5 in Lück's notes

- $f(W) \subseteq Y_1$ and $f(W) \not\subseteq Y_1$
- $f(W) \subseteq Y_2$ and $f(W) \not\subseteq Y_2$

Inductive construction $k = 0$, i.e. vertexes of the cubes $w \in \mathcal{W}$. If $w \in Y_2$, take $h[0]_t = \text{const}_{w_0}$.

Suppose $f(W_0) \in Y_1$, but $f(w_0) \notin Y_2$. Since Y_1, Y_0 is 0-connected, there is a path $\pi: I \rightarrow Y_1$ from w_0 to a point in Y_0 . We take $h[0]$ as the path on w_0 . Analogous if $f(W_0) \in Y_2 \setminus Y_1$.

Inductive Step Let $W \in \mathcal{W}$ be a cube of exact dimension k . Then $\partial W = W \cap C_{k-1}$. Since $(W, \partial W)$ has the HEP, we can extend the previous homotopy $h[k-1]|_{\partial W}$ to some homotopy on W relative to $f|_W$. Let this be $h'[k]: C_k \times I \rightarrow Y$: this satisfies conditions 1. and 2. but not yet 3. and 4.

We produce another homotopy $h[k]''$ and set $h[k] = h[k]' * h[k]''$.

Consider a cube $W \in \mathcal{W}$ of dimension k .

If $f(W) \subseteq Y_0$ set $h[k]''$ as the constant homotopy on W .

If $h[k]'_1(W) \subseteq Y_1$, but $h[k]'_1(W) \not\subseteq Y_2$ there is a homotopy relative ∂W from $h[k]'_1$ to a map $f_1(W) \subseteq Y_0$.

If $k = \dim(W) > p$ then we use the lemma 1.4 for $f = h[k]'_1|_W$ and the resulting homotopy is $h[k]''|_W$.

If $h[k]'_1(W) \subseteq Y_2$ but $h[k]'_1(W) \not\subseteq Y_1$, use the complement case of the lemma⁸ □

[13.10.2025, Lecture 1]
[15.10.2025, Lecture 2]

Now for the actual proof of Blakers Massey Let $F(Y_1, Y, Y_2) = Y_1 \times_Y Y^{[0,1]} \times_Y Y_2$.

Let $F(Y_1, Y_1, Y_0)$ be the subspace of those $w \in F(Y_1, Y, Y_2)$ such that $w([0, 1]) \subseteq Y_1$.

Proposition 1.6. Assume that (Y_1, Y_0) is p -connected, (Y_2, Y_0) is q -connected. Then the pair $(F(Y_1, Y, Y_2), F(Y_1, Y_1, Y_0))$ is $(p + q - 1)$ -connected.

Proof. We consider a map of pairs

$$\phi: (I^n \partial I^n) \rightarrow (F(Y_1, Y, Y_2), F(Y_1, Y_1, Y_0))$$

for $1 \leq n \leq p + q + 1$. We want to homotop ϕ relative ∂I^n to a map with image in $F(Y_1, Y_1, Y_0)$. We use the adjoint

$$\text{maps}(X \times [0, 1], Z) \cong \text{maps}(X, Z^{[0,1]})$$

We let $\Phi: I^n \times I \rightarrow Y$ be the adjoint of ϕ , this is *admissable*, in the sense that

1. $\Phi(x, 0) \in Y_1$ for all $x \in I^n$
2. $\Phi(x, 1) \in Y_2$ for all $x \in I^n$
3. $\Phi(x, t) \in Y_1$ for all $x \in \partial I^n, t \in [0, 1]$

We want a homotopy of Φ through admissable maps to a map $\Phi': I^n \times I \rightarrow Y$ such that $\text{Im}(\Phi') \subseteq Y_1$.

Apply Proposition 11.5 to Φ (with $n+1$ instead of n). This gives a homotopy through admissable maps to $\Psi = g$. Let $h: I^n \times I \times I \rightarrow Y$ be a homotopy witnessing this. $h_0 = \Phi, h_1 F\Psi$.

Consider the projection $\text{pr}: I^n \times I \rightarrow I^n$ away from the last coordinate.

Claim. The image under pr of $\Psi^{-1}(Y \setminus Y_1)$ and $\Psi^{-1}(Y \setminus Y_2)$ are disjoint.

Suppose there is $y \in I^n$ in the intersection of the preimages, so $z_1 \in \Psi^{-1}(Y \setminus Y_1), z_2 \in \Psi^{-1}(Y \setminus Y_2)$ s.t $\text{pr}(x_1) = y = \text{pr}(z_2)$. Let $W = I^{n+1}$ be a subcube of the subdivision of Proposition 11.5 such

⁸rest of the proof next lecture. I am very sure some words won't make sense, as they were unreadable on the board.

that $z_1 \in W$. Since $z_1 \in \Psi^{-1}(Y \setminus Y_1)$, $z_1 \in K_{p+1}(W)$, so $y = \text{pr}(z_1) \in K_p(\text{Im}(W))$. Analogous $y = \text{pr}(z_2) \in G_q(\text{Im}(W))$. Since $p + q > n$, this is a contradiction and the claim is proven.

Claim. The intersection of $\text{pr}(\Psi^{-1}(Y \setminus Y_1))$ with ∂I^n is empty since Ψ is admissible, and thus $\Psi(\partial I^n) \subseteq Y_1$. So $\text{pr}(\Psi^{-1}(Y \setminus Y_1))$ and $\text{pr}(\Psi^{-1}(Y \setminus Y_2)) \cup (\partial I^n)$ are two disjoint closed subsets of the compact space I^n .

So there is a continuous separating function $\tau: I^n \rightarrow [0, 1]$, s.t. $\tau \equiv 0$ is $\text{pr}(\Psi^{-1}(Y \setminus Y_1))$ and $\tau \equiv 1$ is $\partial I^n \cup \text{pr}(Y \setminus Y_2)$

We define another homotopy starting with Ψ by

$$h: (I^n \times I) \times I \rightarrow Y$$

by

$$((x, t), s) \mapsto \Psi(x, (1 - s)t + s \cdot t\tau(x))$$

This is

- Homotopy through admissible maps
- $h(x, t, 0) = \Psi(x, t)$

Claim. $h(_, _, 1)$ has image in Y_1 .

$$h(x, t, 1) = \Psi(x, t \cdot \tau(x))$$

- if $x \in \Psi^{-1}(Y \setminus Y_1)$, then $\tau(x) = 0$, $h(x, t, 1) = \Psi(x, 0) \in Y_1$.
- if $x \in \Psi^{-1}(Y_1)$, then by admissibility $\Psi(x, t \cdot \tau(x)) \in Y$.

□

For the actual proof (Y_1, Y_0) p -connected (Y_2, Y_0) q -connected
some diagram about something being Serre fibration

We compare two Serre fibrations

$$F(\{y_0\}, Y_1, Y_0) \quad F(\{y_0\}, Y, Y_2)$$

$$F(Y_1, Y_1, Y_0) \longrightarrow F(Y_1, Y, Y_2)$$

$$Y_1$$

$$Y_1$$

partial 5-lemma shows that also the pair

$$(F(\{y_0\}, Y, Y_2), F(\{y_0\}, Y_1, Y_0))$$

is $(p + q - 1)$ -connected.

The following square commutes for all $n \geq 1$

$$\begin{array}{ccc} \pi_{n-1}(F(\{y_0\}, Y_1, Y_0), \text{const}_{y_0}) & \longrightarrow & \pi_{n-1}(F(\{y_0\}, Y, Y_1), x) \\ \downarrow \cong & & \downarrow \cong \\ \pi_n(Y_1, Y_0, y_0) & \xrightarrow{\text{id}_*} & \pi_n(Y, Y_1, y_0) \end{array}$$

□

1.2 Freudenthal suspension theorem

Section 6.4 in tom Dieck's book

Definition 1.7

Let X be a based space. The *unreduced suspension* is $X^\diamond = X \times [-1, 1] / \sim$ where \sim identifies all points with second variable -1 to S and all points with second variable 1 to N . We use S as the base point of X^\diamond .¹

¹south and north pole

Note. If X is well based, i.e. $\{x_0\} \hookrightarrow X$ has the HEP, then the quotient map $X^\diamond \rightarrow X = \frac{X \times [0, 1]}{X \times \{0, 1\} \cup \{x_0\} \times [0, 1]}$ is a homotopy equivalence.

The suspension homomorphism $S: \pi_k(X, x_0) \rightarrow \pi_{k+1}(X^\diamond, S)$ for $k \geq 1$ is $S[f: S^k \rightarrow X] := [f^\diamond: S^{k+1} = (S^k)^\diamond \rightarrow X^\diamond]$

Theorem 1.8: Freudenthal suspension theorem

Let X be an $(n-1)$ -connected space, $n \geq 1$. Then the suspension homomorphism is

- bijective for $i \leq 2n-2$
- surjective for $i = 2n-1$

Proof. We cover X^\diamond by $Y_1 = X^\diamond \setminus \{N\}$, $Y_2 = X^\diamond \setminus \{S\}$. We use repeatedly that Y_1 and Y_2 are contractible. Then $X \rightarrow Y_1 \cap Y_2 = Y_0$ is a homotopy equivalence. We claim without proof that the following diagram commutes:

I couldn't keep up.

So we may show that $\pi_{i+1}(Y_2, Y_0, x_0) \rightarrow \pi_{i+1}(Y, Y_1, y_0)$ is bijective for $i \leq 2n-2$ and surjective for $i \leq 2n-1$.

Because X is $(n-1)$ -connected

$$\pi_{k+1}(Y_1, Y_0, x_0) \xrightarrow[\partial]{\cong} \pi_k(Y_0, x_0)$$

So $\pi_i(Y_1, Y_0, x_0) = 0$ for $i \leq n$, i.e. (Y_1, Y_0) is n -connected. Also (Y_2, Y_0) is n -connected.

BM gives $\pi_k(Y_2, Y_0, y) \rightarrow \pi_k(X^\diamond, Y_1, y)$ is bijective for $k < 2n-1$ and surjective for $k = 2n$.

Setting $i = k-1$ ends the proof. \square

Take $X = S^n$

Corollary 1.9. The suspension homomorphism

$$S: \pi_i(S^n, *) \rightarrow \pi_{i+1}(S^{n+1}, *)$$

is bijective for $i \leq 2n-2$ and surjective for $i = 2n-1$.

Corollary 1.10. For all $n \geq 1$, $\pi_n(S^n, *) \cong \mathbb{Z}$, generated by the class of id_{S^n} . Moreover $\text{deg}: \pi_n(S^n, *) \rightarrow \mathbb{Z}$ is an isomorphism.

Proof. Induction on n . For $n = 1$ we have this by covering theory.

$n \geq 1$ By Freudenthal, the suspension homomorphism

$$S: \pi_n(S^n, *) \rightarrow \pi_{n+1}(S^{n+1}, *)$$

is surjective. The composite

$$\pi_n(S^n, *) \xrightarrow{S} \pi_{n+1}(S^{n+1}, *) \xrightarrow{\deg} \mathbb{Z}$$

is bijective by induction. So S is also injective and \deg in one dimension higher is also an isomorphism. \square

Recall. $\pi_3(S^2, *) \cong \mathbb{Z}\{\eta\}$, where $\eta: S^3 \rightarrow S^2$ is the Hopf map.

Proposition 1.11. For $n \geq 3$, $\pi_{n+1}(S^n, *) = \mathbb{Z}/2\{S^{n-2}(\eta)\}$ cycles of order two.

Proof. Only partial. For $n \geq 2$, $S: \pi_{n+1}(S^n, q) \mapsto \pi_{n+2}(S^{n+1}, S^{n+1}, *)$ is surjective.

$$\mathbb{Z}\{\eta\} = \pi_3(S^2, *) \rightarrow \pi_4(S^3, *) \xrightarrow{S} \pi_5(S^4, *) \dots$$

Claim. $S(2\eta) = 0$ in $\pi_4(S^3, *)$. Which gives $\pi_{n+1}(S^n, *)$ is either trivial or order 2.

Consider the commutative square

I did not manage to copy. Something complex conjugation

$$\implies [\eta] = [\eta \circ \text{complex conjugation}] = [d \circ \eta] \text{ in } \pi_3(S^2, * \cong \mathbb{Z}).$$

If $f: S^n \rightarrow S^n$ has degree k , then precomposition of $\pi_n(X, x) \rightarrow \pi_n(X, x)$ is multiplication by k .

Let $c: S^1 \rightarrow S^1$ be any map of degree -1 . Then $c \wedge S^2, S^1 \wedge d$: both have degree -1 . So $c \wedge S^2 \sim S^1 \wedge d$

two diagrams I did not copy.

We get $S(\eta) = S(d \circ \eta) = S(\eta \circ (c \wedge S^1)) = S(-\eta) = -S(\eta)$ and hence $2 \cdot S(\eta) = 0$. \square

[13.10.2025, Lecture 2]
[20.10.2025, Lecture 3]

We make a bit of a preview⁹ for stable homotopy theory. Following Lück's notes today rather closely

Definition 1.12

Let X be a based space. The n -th stable homotopy group of X is the colimit $\pi_n^S(X, *)$

$$\pi_n(X, *) \xrightarrow{S} \pi_{n+1}(S^1 \wedge X, *) \xrightarrow{S} \pi_{n+2}(S^2 \wedge X, *) \dots$$

Since $S^k \wedge X$ is $(k-1)$ -connected, the suspension stabilises (i.e. S is isomorphism for $\pi_{n+k}(S^k \wedge X, *)$ onwards).

$\pi_n^S()$ is a functor from based spaces to abelian groups. and it is homotopy invariant. It comes with a natural transformation $\pi_n(X, *) \rightarrow \pi_n^S(X, *)$.

Preview: $\{\pi_n^S\}_{n \in \mathbb{Z}}$ form a generalized homology theory on based spaces.

If (X, A) is a pair of based space, $x \in A \subseteq X$, we define

$$\pi_n^S(X, A, *) := \pi_n^S(X \cup_A CA, *)$$

The map collapsing X is

$$X \cup_A CA \rightarrow \frac{X \cup_A CA}{X} \cong \frac{CA}{A} \cong A^\diamond$$

⁹He just spammed random stuff, I don't think I copied enough for it to make sense.

induces an connecting homomorphism

$$\partial: \pi_n^S(X, A, x) = \pi_n^S(X \cup_A CA, *) \xrightarrow{p_*} \pi_n^S(A^\circ) \rightarrow \pi_n^S(A \wedge S^1) \xrightarrow{\cong} \pi_{n-1}^S(A)$$

The following sequence will then be exact:

$$\dots \pi_n^S(A, x) \xrightarrow{\text{incl}_*} \pi_n^S(X, x,) \xrightarrow{\text{incl}_*} \pi_n^S(X, A, *) \xrightarrow{\partial} \pi_{n-1}^S(A, *) \rightarrow \dots$$

Stable stems are the special case

$$\pi_n^S(S^0) = \text{colim}(\pi_n(S^0, *) \rightarrow \pi_{n+1}(S^1, *) \rightarrow \dots)$$

n	0	1	2	3	4	5	6	7	8
$\pi_n^S = \pi_n^S(S^0)$	\mathbb{Z}	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/24$	0	0	$\mathbb{Z}/2$	$\mathbb{Z}/240$	$(\mathbb{Z}/2)^2$
Generator	id	η	η^2	$\nu, \eta^3 = 12\nu$	-	-	ν^2	σ	$\eta\sigma, \epsilon$

There is a graded-commutative ring structure on $\pi_*^S = \{\pi_n^S\}_{n \in \mathbb{Z}}$. $\pi_n^S \times \pi_m^S \rightarrow \pi_{n+m}^S$

$$[f: S^{n+k} \rightarrow S^k \times S^k] \times [g: S^{m+l} \rightarrow S^l] = [SW_{n+m+k+l} \rightarrow S^{n+k} \wedge S^{m+l} \xrightarrow{f \wedge g} S^{k+l}]$$

I missed a bit more

Nishidas theorem says: every poitive dimensional element of π_*^3 is nilpotent.

From Serre spectral sequence we will see: For $m > n \geq 1$ $\pi_m(S^n, *)$ is finite except $\pi_{4k-1}(S^{2k}, *) \cong \mathbb{Z} \oplus \text{some finite group}$.

This was exercise 5.2 of last summer term: *Hopf invariant* $h: \pi_{2k-1}(S^k, *) \rightarrow \mathbb{Z}$ for $[f: S^{2k-1} \rightarrow S^k]$

$$H^*(S^k \cup_f D^{2k}, \mathbb{Z}) = \begin{cases} \mathbb{Z} & i = 0, k, 2k \\ 0 & \text{else} \end{cases}$$

we have $H^k(Cf, \mathbb{Z}) = \mathbb{Z}\{a\}$ $H^{2k}(Cf, \mathbb{Z}) \cong \mathbb{Z}\{b\}$ and $a \cup a = h(f) \cdot b$.

In general we can look at

$t_k: S^{2k-1} \rightarrow S^k \vee S^k$ the attaching map of the $2k$ -cell in the product minimal CW-structure of $S^k \times S^k$. And we get $h(t_k) = 2$.

The Hopf invariant 1 theorem tells us when we can realize Hopf invariant 1 and here we see that we always find Hopf invariant 2.

1.3 Hurewicz-theorem

This is a "trivial"¹⁰ Corollary of the Blakers-Massey theorem.

Definition 1.13

Let X be a based space. Choose an orientation of S^n , $n \geq 1$, i.e. $[S^n] \in H_n(S^n, \mathbb{Z})$. The Hurewicz homomorphism

$$h: \pi_n(X, x) \rightarrow H_n(X; \mathbb{Z})$$

is defined by

$$h[f: S^n \rightarrow X] = H_n(f, \mathbb{Z})[S^n].$$

This is a group homomorphism.

¹⁰meaning 1.5 lectures of intermediate steps

Group Homomorphism: Let $\Delta: S^n \rightarrow S^n \vee S^n$ be a pinch map. Group structure on $\pi_n(X, x)$ is given by $[f] + [g] = [S^n \xrightarrow{\Delta} S^n \vee S^n \xrightarrow{f+g} X]$. Then

$$\begin{array}{ccc} \pi_n(X, x) & & \\ \cong \downarrow & \searrow h & \\ \pi_n(X, y) & \xrightarrow{h} & H_n(X, \mathbb{Z}) \end{array}$$

$$([f] + [g])_* = [f]_* + [g]_*: H_n(S^n) \rightarrow H_n(X)$$

So Hurewicz is a homomorphism.

Lemma 1.14

Let $w: [0, 1] \rightarrow X$ be a path from $x = w(0)$ to $y = w(1)$. Then the following commutes:

$$\begin{array}{ccc} \pi_n(X, x) & & \\ \cong \downarrow & \searrow h & \\ \pi_n(X, y) & \xrightarrow{h} & H_n(X, \mathbb{Z}) \end{array}$$

Proof. f and $f * w$ are freely homotopic $h_n(_, \mathbb{Z})$ is homotopy invariant □

For $n = 1$ we have Poincaré: X based path connected, $h: \pi_1(X, x) \rightarrow H_1(X, \mathbb{Z})$ is surjective with kernel the commutator subgroup or equivalently

$$\pi_1(X, x)_{ab} \xrightarrow{\cong} H_1(X, \mathbb{Z})$$

Theorem 1.15: Hurewicz

Let X be an $(n - 1)$ -connected space, $n \geq 2$. Then the Hurewicz homomorphism

$$h: \pi_n(X, x) \rightarrow H_n(X, \mathbb{Z})$$

is an isomorphism.

We will need to work a bit for the proof.

Proposition 1.16 (11.9 in Lück's notes). Let $m, n \geq 0$, let

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ \downarrow i & & \downarrow \bar{i} \\ X & \xrightarrow{\bar{f}} & Y \end{array}$$

be a pushout square of spaces. Suppose $i: A \rightarrow X$ is a *cofibration*, i.e. a closed embedding with the HEP.

1. If f is n -connected, then so is \bar{f}
2. If f is n -connected, and i is m -connected, then for all $a \in A$, $\pi_i(f, \bar{f}): \pi_i(X, A, a) \rightarrow \pi_i(Y, B, f(a))$ is bijective for $1 \leq i < m + n$ and surjective for $i = m + n$.

Proof. "Basically reduction to Blakers-Massey"

We can replace X, B and Y up to homotopy equivalence by appropriate mapping cylinders:

$$\text{cyl}(f) = A \times [0, 1] \cup_{A \times 1, f} B$$

We get a homotopy commutative diagram

$$\begin{array}{ccccc} A & \longrightarrow & \text{Cyl}(f) & & \\ \downarrow & & \downarrow & \searrow \sim & \\ \text{Cyl}(i) & \longrightarrow & \text{cyl}(i) \cup_A \text{cyl}(f) & & B \\ & \searrow \sim & \downarrow \sim & \searrow \sim & \downarrow \bar{i} \\ & & X & \xrightarrow{\bar{f}} & Y \end{array}$$

This diagram does not commute. The homotopy equivalences shown are not trivial.

We apply Blakers-Massey then to the following open subspace of $W = \text{cyl}(i) \cup_A \text{cyl}(f)$.

$$W_2 = [0, 1/2) \times A \cup_{A \times 0} \text{cyl}(f), \quad W_1 = \text{cyl}(i) \cup_{A \times 0} [0, 1/2). \quad W_0 = W_1 \cap W_2 \sim A$$

Now having f is n -connected gives (W_2, W_0) is n -connected. i being m -connected gives (W_1, W_0) is m -connected.

Now BM gives $\pi(W_1, W_0, a) \rightarrow \pi_i(W, W_2, a)$ is bijective for $1 \leq i \leq m + n - 1$ and surjective for $i = m + n$.

Using the homotopy equivalences, we translate this back. □

Proposition 1.17 (11.1 in Lück). Let $m, n \geq 0$, let $c: A \rightarrow X$ be a cofibration. Suppose that i is m -connected and A is n -connected. Then

$$\pi_k(\text{pr}): \pi_k(X, A, a) \rightarrow \pi_k(X/A, *)$$

for all $a \in A$ is bijective for $1 \leq k \leq m + n$ and surjective for $k = m + n + 1$

Proof. We consider the pushout

$$\begin{array}{ccccc} A & \xrightarrow{j} & C(A) = A \times [0, 1] / A \times 1 \\ \downarrow i & & \downarrow & & \\ X & \longrightarrow & X \cup_A CA & \xrightarrow{\sim} & X/A \end{array}$$

Remark 1.18. X k -connected $\Leftrightarrow \{*\} \rightarrow X$ k -connected $\Leftrightarrow X \rightarrow \{*\}$ is $(n + 1)$ -connected. □

Proposition 1.19 (11.12 for Lück). Also an exercise (1.2) Let X, Y be well-pointed spaces. Let $m, n \geq 1$. Let X be m -connected, Y n -connected. Then

1. The inclusion $X \vee Y \rightarrow X \times Y$ induces isomorphisms

$$\pi_k(X \vee Y, *) \rightarrow \pi_k(X \times Y, *)$$

for all $0 \leq k \leq m + n$.

2. $\pi_k(X \times Y, X \vee D, *)$ and $\pi_k(X \wedge @, *)$ are trivial for all $0 \leq k \leq m + n + 1$.

3. The map $(pr_*^X, pr_*^Y): \pi_k(X \vee Y, x) \rightarrow \pi_k(X, x_0) \times \pi_k(Y, y_0)$ is an isomorphism for all $1 \leq k \leq m + n$.

[20.10.2025, Lecture 3]
[22.10.2025, Lecture 4]

Proposition 1.20. Let $n \geq 2$. Let $\{X_i\}_{i \in I}$ be a family of well-pointed, $(n-1)$ -connected spaces. Then the canonical map

$$\bigoplus_{i \in I} \pi_n(X_i, *) \rightarrow \pi_n\left(\bigvee_{i \in I} X_i, *\right)$$

is an isomorphism.

Proof. Step 1 If I is finite, we do Induction on $|I|$. Nothing to show if $I = \emptyset$, $|I| = 1$. Let $|I| \geq 2$. Wlog $I = \{1, 2, \dots, k\}$, $k \geq 2$.

By Proposition 1.19, $\pi_n(\bigvee_{i=1, \dots, k-1} X_i, *) \oplus \pi_n(X_k, *) \xrightarrow{\cong} \pi_n(\bigvee_{i=1, \dots, k} X_i, *)$ and the first part is isomorphic to $\bigoplus_{i=1, \dots, k} \pi_n(X_i)$ by induction.

Case 2 I is infinite: We first show injectivity. The projection

$$p_j: \bigvee_{i \in I} X_i \rightarrow X_j$$

induces an homomorphism

$$(p_j)_*: \pi_n\left(\bigvee_I X_i, *\right) \rightarrow \pi_n(X_j, *)$$

For varying $j \in I$, these assemble into a homomorphism

$$\bigoplus_{i \in I} \pi_n(X_i, *) \xrightarrow{\text{can}} \pi_n\left(\bigvee_{i \in I} X_i, *\right) \xrightarrow{((p_j)_*)} \prod_{j \in I} \pi_n(X_j, *)$$

and the composition is inclusion of sum into product, hence injective. So also the first map is injective.

For surjectivity let $f: S^n \rightarrow \bigvee_{i \in I} X_i$ be a continuous map that represents a class in $\pi_n(\bigvee_{i \in I} X_i)$. Because S^n is compact, there is a finite subset $J \subseteq I$ s.t. $\text{Im}(f) \subseteq \bigvee_{i \in J} X_i$ ¹¹. This implies $[f] \subseteq \pi_n(\bigvee_{j \in J} X_j, *) \rightarrow \pi_n(\bigvee_{i \in I} X_i, *)$.

$$\begin{array}{ccc} \pi_n(\bigvee_{j \in J} X_j, *) & \longrightarrow & \pi_n(\bigvee_{i \in I} X_i, *) \\ \text{Case 1} \uparrow & & \uparrow \text{can} \\ \bigoplus_{j \in J} \pi_n(X_j, *) & \longrightarrow & \bigoplus_{i \in I} \pi_n(X_i, *) \end{array}$$

so $[f]$ is in the image of the canonical map.

□

Theorem 1.21: Hurewicz

Let $n \geq 2$. Let X be $(n-1)$ -connected base space. Then the Hurewicz homomorphism $h: \pi_n(X, x) \rightarrow H_n(X, \mathbb{Z})$, $f \mapsto H_n(f, \mathbb{Z})[S^n]$ is an isomorphism.

¹¹he explains, why this is not easy to see. But it is point-set topology, so we won't do it.

- Proof.* • By CW-approximation (see later in this class) there is a CW-complex Y with one 0-cell and no cells in dimensions $1 \leq i \leq n-1$ and a weak homotopy equivalence $Y \xrightarrow{\simeq} X$.
- Every weak homotopy equivalence induces isomorphisms of $H_n(_, A)$ for all $n \geq 0$, for all abelian groups A . This will either be an exercise or proven later on.

We get a commutative diagram

$$\begin{array}{ccc} \pi_n(Y, y) & \xrightarrow[\cong]{f_*} & \pi_n(X, f(*)) \\ \downarrow h & & \downarrow h \\ H_n(Y, \mathbb{Z}) & \xrightarrow[\cong]{f_*} & H_n(X, \mathbb{Z}) \end{array}$$

So wlog we can assume that X admits a CW-structure with a single 0-cell and no cells of dimensions $1, \dots, n-1$. The inclusion of the $(n+1)$ -skeleton $X_{n+1} \rightarrow X$ induces isomorphisms

$$\pi_n(X_{n+1}, *) \rightarrow \pi_n(X, *)$$

by cellular approximation. Also

$$H_n(X_{n+1}, \mathbb{Z}) \rightarrow H_n(X, \mathbb{Z})$$

is an isomorphism for example by cellular homology.

It suffices to show the Hurewicz theorem for the $n+1$ -skeleton, i.e. X is a CW-complex with a single 0-cell, I many n -cells, J many $(n+1)$ -cells and no cells in any other dimension.

$$X = (\{x\} \cup_{I \times \partial D^n} D^n) \cup_{J \times \partial D^{n+1}} (J \times D^{n+1}) \cong (\bigvee_{i \in I} S^n) \cup_{J \times D^{n+1}} J \times D^{n+1}$$

We can assume that all attaching maps $\alpha: \partial D^{n+1} \rightarrow \bigvee_{i \in I} S^n$ are based.

We see this since $\bigvee_{i \in I} S^n$ is path connected, α is homotopic by HEP to a based map α' . Since homotopic attaching maps yield homotopy equivalent glueings.

So X can be written as a pushout

$$\begin{array}{ccc} \bigvee_{j \in J} S^n & \xrightarrow{f} & \bigvee_{i \in I} S^n \\ \downarrow & & \downarrow \\ \bigvee_{j \in J} D^{n+1} & \longrightarrow & X \end{array} = \begin{array}{ccc} & & X_n \\ & & \\ & & X_{n+1} \end{array}$$

Since $\bigvee_I S^n$ and $\bigvee_J S^n$ are $(n-1)$ -connected, $f: \bigvee_J S^n \rightarrow \bigvee_I S^n$ is $(n-1)$ -connected. Also $\bigvee_J S^n \rightarrow \bigvee_I D^{n+1}$ is n -connected, as $\bigvee_I D^{n+1}$ is contractible.

By Proposition ?? $\pi_k(\bigvee_J D^{n+1}, \bigvee_J S^n) \rightarrow \pi_k(X, X_n) = \pi_k(X, \bigvee_I S^n)$ is isomorphic for $1 \leq k \leq 2n-2$ and surjective for $k = 2n-1$. In particular $\pi_{n+1}(\bigvee_J D^{n+1}, \bigvee_J S^n) \rightarrow \pi_{n+1}(X, X_n)$ is surjective. We compare the long exact homotopy sequences of the vertical pairs:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \pi_{n+1}(\bigvee_J D^{n+1}, \bigvee_J S^n) & \xrightarrow[\cong]{\partial} & \pi_n(\bigvee_J S^n) & \longrightarrow & 0 \\ & & \downarrow g_* & & \downarrow f_* & & \\ & \longrightarrow & \pi_{n+1}(X, X_n) & \xrightarrow{\partial} & \pi_n(X_n) & \xrightarrow{\text{surjective by cell. approx}} & \pi_n(X, *) \end{array}$$

So the upper row is exact:

$$\begin{array}{ccccccc}
 \pi_n(\bigvee_J S^n) & \xrightarrow{f_*} & \pi_n(\bigvee_I S^n) & \xrightarrow{\text{incl}_*} & \pi_n(X) & \longrightarrow & 0 \\
 \downarrow h & & \downarrow h & & \downarrow h & & \\
 H_n(\bigvee_J S^n, \mathbb{Z}) & \xrightarrow{f_*} & H_n(\bigvee_I S^n, \mathbb{Z}) & \longrightarrow & H_n(X, \mathbb{Z}) & \longrightarrow & 0
 \end{array}$$

We claim the first 2 h are isomorphisms. We have a long exact sequence by excision. For all sets I

$$\begin{array}{ccc}
 \bigoplus_{i \in I} \pi_n(S^n, *) & \xrightarrow{\cong} & \pi_n(\bigvee_{i \in I} S^n, *) \\
 \downarrow \bigoplus h & & \downarrow h \\
 \bigoplus_{i \in I} H_n(S^n, \mathbb{Z}) & \xrightarrow{\cong} & H_n(\bigvee_I S^n, \mathbb{Z})
 \end{array}$$

so $h: \pi_n(X, x) \rightarrow H_n(X, \mathbb{Z})$ is an isomorphism by 5-lemma. \square

Some applications

Recall. If X is simply connected, then $H_1(X, \mathbb{Z}) \cong \pi_1(X, x)_{\text{ab}} = 0$. But if X is path connected and $H_1(X, \mathbb{Z}) = 0$, X need not be simply connected, because $\pi_1(X, x)$ could be non-trivial and perfect (= abelianization is trivial) (E.g. A_5).

Proposition 1.22 (12.7(i) for Lück). Let X be simply connected, $n \geq 1$. then the following are equivalent

1. X is n -connected.
2. $H_i(X, \mathbb{Z}) = 0$ for all $2 \leq i \leq n$.

Proof. By induction on n . Nothing to show for $n = 1$. The induction step is the Hurewicz theorem $\pi_n(X, x) \cong H_n(X, \mathbb{Z})$. \square

Proposition 1.23 (12.7 (ii) for Lück). Let X be simply connected. Then the following are equivalent:

1. X is weakly contractible¹²
2. $H_i(X, \mathbb{Z}) = 0$ for all $i \geq 2$.

Warning. There exist acyclic spaces, i.e. non-contractible CW-complexes, path connected with $H_i(X, \mathbb{Z}) = 0$ for all $i \geq 1$.

Remark 1.24. There is a slightly better version of the Hurewicz theorem: If X is $n - 1$ -connected, $n \geq 2$, then $h: \pi_n(X, x) \rightarrow H_n(X, \mathbb{Z})$ is isomorphism and $h: \pi_{n+1}(X, x) \rightarrow H_{n+1}(X, \mathbb{Z})$ is surjective.

¹²All homotopy groups vanish

1.3.1 Relative Hurewicz theorem

Definition 1.25: Relative Hurewicz map

Let (X, A) be a space pair, $a \in A$. Choose a generator $[D^n, S^{n-1}] \in H_n(D^n, S^{n-1}, \mathbb{Z})$. The relative Hurewicz homomorphism is

$$h: \pi_n(X, A, a) \rightarrow H_n(X, A, \mathbb{Z})$$

is defined by

$$[f: (D^n, S^{n-1}, z) \rightarrow (X, A, \{a\})] \mapsto h[f] := H_n(f, \mathbb{Z})[D^n, S^{n-1}]$$

The following diagram commutes:

$$\begin{array}{ccccc} \pi_n(X, a) & \longrightarrow & \pi_n(X, A, a) & \xrightarrow{\partial} & \pi_{n-1}(A, a) \\ \downarrow h & & \downarrow h & & \downarrow h \\ H_n(X, \mathbb{Z}) & \longrightarrow & H_n(X, A, \mathbb{Z}) & \xrightarrow{\partial} & H_{n-1}(A, \mathbb{Z}) \end{array}$$

Theorem 1.26: relative Hurewicz, (simply connected case)

Let $n \geq 2$. Let (X, A) be a space pair, such that X and A are simply connected and (X, A) is $(n-1)$ -connected. Then

1. The Hurewicz homomorphism $h: \pi_n(X, A, a) \rightarrow H_n(X, A, \mathbb{Z})$ is an isomorphism, and
2. The group $H_i(X, A, \mathbb{Z}) = 0$ for all $0 \leq i \leq n-1$.

Proof. We will deduce this from the absolute version and some other things we already did.

By replacing X by the mapping cylinder $A \times [0, 1] \cup_{A \times 1} X$ and replacing A by $A \times 0$, we can assume wlog that the inclusion $i: A \rightarrow X$ is a cofibration. Since A is 1-connected, and (X, A) is $(n-1)$ -connected,

$$pr: \pi_k(X, A, a) \rightarrow \pi_k(X/A, *)$$

is bijective for $1 \leq k \leq n$ and surjective for $k = n+1$.

X/A is simply connected by the van Kampen theorem.

Since $\pi_k(X, A, *) = 0$ for $k \leq n-1$ by hypothesis, we get that $\pi_k(X/A, *) = 0$ for $k \leq n-1$. So X/A is $(n-1)$ -connected. By the absolute Hurewicz theorem for X/A , $H_k(X/A, \mathbb{Z}) = 0$ for $1 \leq k \leq n-1$ and

$$\begin{array}{ccc} \pi_n(X, A, a) & \xrightarrow[\cong]{pr_*} & \pi_n(X/A, *) \\ \downarrow & & \downarrow h \cong \\ H_n(X, A, \mathbb{Z}) & \xrightarrow[\cong]{\text{excision}} & H_n(X/A, \mathbb{Z}) \end{array}$$

□

[22.10.2025, Lecture 4]
[27.10.2025, Lecture 5]

Proposition 1.27. Let $f: X \rightarrow Y$ be a map of simply connected spaces. Let $n \geq 1$. Then TFAE:

1. f is n -connected
2. $f_*: H_i(X, \mathbb{Z}) \rightarrow H_i(Y, \mathbb{Z})$ is bijective for $2 \leq i \leq n-1$ and surjective for $i = n$.

Proof. By replacing Y by the mapping cylinder of $f: X \rightarrow Y$, we may assume f is the inclusion $i: A \hookrightarrow X$ of a closed subspace (X, A) is n -connected is equivalent to $\pi_k(X, A, a) = 0$ for all $k \leq n$ which is then by Hurewicz equivalent to $H_k(X, A, \mathbb{Z}) = 0$ for all $k \leq n$. But this is equivalent to 2. \square

Theorem 1.28: Whitehead theorem

Let $f: X \rightarrow Y$ be a continuous map between simply connected CW-complexes. Then the following are equivalent:

1. f is a homotopy equivalence.
2. f is a weak homotopy equivalence.
3. $f_*: H_i(X, \mathbb{Z}) \rightarrow H_i(Y, \mathbb{Z})$ is an isomorphism for all $i \geq 0$.

Note. $1 \Leftrightarrow 2$ without simply connectedness is what we previously called the Whitehead theorem.

Theorem 1.29

Let $f: X \rightarrow Y$ be a continuous map between path connected CW-complexes. Suppose that for some (hence any) $x \in X$ $f_*: \pi_1(X, x) \rightarrow \pi_1(Y, f(x))$ is an isomorphism. Let $\tilde{f}: \tilde{X} \rightarrow \tilde{Y}$ be a lift to universal covers.

$$\begin{array}{ccc} \tilde{X} & \xrightarrow{f} & \tilde{Y} \\ \downarrow q & & \downarrow q \\ X & \xrightarrow{f} & Y \end{array}$$

Then TFAE:

1. f is a homotopy equivalence
2. $\tilde{f}_*: H_i(\tilde{X}, \mathbb{Z}) \rightarrow H_i(\tilde{Y}, \mathbb{Z})$ is an isomorphism for all $i \geq 0$.

Proof. $p_*: \pi_i(\tilde{X}, \tilde{x}) \rightarrow \pi_i(X, x)$ is an isomorphism for $i \geq 2$. So f is a weak homotopy equivalence iff $\tilde{f}: \tilde{X} \rightarrow \tilde{Y}$ is a weak homotopy equivalence. and now this is equivalent too $H_*(\tilde{f}, \mathbb{Z})$ is an isomorphism for all $* \geq 0$. \square

Theorem 1.30: 12.16 for Lück

Let X be a path connected CW-complex, $n \geq 2$. Then the following are equivalent:

1. X is homotopy equivalent to S^n
2. X is simply connected and $H_i(X, \mathbb{Z}) \cong \begin{cases} \mathbb{Z} & i = 0, n \\ 0 & \text{else} \end{cases}$

Proof. 1. \implies 2.

2. \implies 1. By the Hurewicz Theorem $h: \pi_n(X, x) \rightarrow H_n(X, \mathbb{Z}) \cong \mathbb{Z}$ is an isomorphism. Let $f: S^n \rightarrow X$ represent a generator of $\pi_n(X, x)$. Then f induces an isomorphism of all integral homology groups. Since S^n and X are simply connected CW-complexes, f is a homotopy equivalence.

□

We will not proof a even more general relative Hurewicz theorem:

Let (X, A) be a space pair, $a \in A$. Recall that $\pi_1(A, a)$ acts on $\pi_n(X, A, a)$ for $n \geq 1$:

With $f: (I^n, \partial I^n, J^{n-1}) \rightarrow (X, A, a)$ and $w: ([0, 1], \{0, 1\}) \rightarrow (A, a)$ with $w * f$, which he explains by a picture. Note $w * f$ is pair homotopic to f (but **Not** triple homotopic).

If $[I^n, \partial I^n] \in H_n(I^n, \partial I^n, \mathbb{Z}) \cong \mathbb{Z}$ is a generator, then

$$f_*[I^n, \partial I^n] = (w * f)_*[I^n, \partial I^n]$$

so $h: \pi_n(X, A, a) \rightarrow H_n(X, A, \mathbb{Z})$ satisfies $h[f] = h[w * f]$.

Definition 1.31

Let (X, A, a) be a space triple, $n \geq 2$. Set

$$\pi_n(X, A, a)^\dagger := \text{quotient of } \pi_n(X, A, a) \text{ by the normal subgroup generated by } [w * f] \cdot [f]^{-1}$$

Note. For $n \geq 3$, the group $\pi_n(X, A, a)$ is abelian, hence so is $\pi_n(X, A, a)^\dagger$. For $n = 2$ $\pi_2(X, A, a)$ need not be abelian, but $\pi_2(X, A, a)^\dagger$ is.

Let $f, g: (I^2, \partial I^2, J^1) \rightarrow (X, A, a)$, set $w = g|_{[0,1]}: ([0, 1], \{0, 1\}) \rightarrow (A, a)$. The rest of the proof is pictures, so good luck understanding without them. I'm sorry. We have $[g]^{-1} \cdot [f] \cdot [g] = [w] * [f]$ See this in Tom Dieck Prop. 6.2.6.

So in particular $[g]^{-1} \cdot [f] \cdot [g] \equiv [f]$ in $\pi_2(X, A, a)^\dagger$, so it is abelian.

Theorem 1.32: Relative Hurewicz with π_1

Let (X, A) be a path connected space pair. Suppose for all $a \in A$, $\text{incl}_*: \pi_1(A, a) \rightarrow \pi_1(X, a)$ is an isomorphism. Let $n \geq 2$ be such that $\pi_i(X, A, a) = 0$ for all $1 \leq i \leq n - 1$. Then $H_i(X, A, \mathbb{Z}) = 0$ for $0 \leq i \leq n - 1$ and the modified Hurewicz map

$$h^\dagger: \pi_n(X, A, a)^\dagger \rightarrow H_n(X, A, \mathbb{Z})$$

is an isomorphism.

Warning. The hypothesis is on $\pi_i(X, A, a)$, but the conclusion on $\pi_n(X, A, a)^\dagger$!

1.4 CW-Approximation

"Every topological space can be approximated by a CW-complex"¹³. More detailed, you can find a CW-complex with a weak homotopy equivalence to your space.

we state a relative refined version of CW-approximation and proof that.

¹³See 6 in Lücks notes

Definition 1.33

Let (Y, A) be a space pair, $n \geq 0$. A n -CW-model for (Y, A) is a relative CW-complex (Z, A) and a certain map $f: Z \rightarrow Y$ such that

- $f|_A = \text{inclusion } A \hookrightarrow Y$
- (Z, A) is n -connected,
- The map $f_*: \pi_i(Z, z) \rightarrow \pi_i(Y, f(z))$ is injective for $i = n$ and bijective for $i > n$ for all $z \in Z$.

We have

$$A \xleftarrow[n\text{-connected}]{\text{relative CW}} Z \xrightarrow[n\text{-coconnected}]{} Y$$

Theorem 1.34

Let (Y, A) be a space pair, $A \neq \emptyset$, such that A is Hausdorff, $n \geq 0$. Then there is a n -CW-model (A, Z, f) , s.t. (Z, A) has no relative cells of dimension $\leq n$.

Addendum. If A comes with a CW-structure, then Z can be chosen as a CW-complex that contains A as a subcomplex.

Special Case. Y is n -connected, $n \geq 0$, $A = \{y_0\}$. Let $(Z, \{y_0\})$ be a n -CW-model with $f: Z \rightarrow Y$, without relative i -cells for $0 \leq i \leq n$. Then both Y and Z are n -connected, $f: \pi_i(Z, z) \rightarrow \pi_i(Y, z)$ is an isomorphism for $i > n$, hence f is a weak homotopy equivalence

Proof. We will inductively construct

$$A = Z_n \subseteq Z_{n+1} \subseteq \dots$$

and $f_i: Z_i \rightarrow Y$, $i \geq n$, such that

- for $i > n$, Z_i is obtained from Z_{i-1} by attaching i -cells.
- $f_i|_{Z_{i-1}} = f_{i-1}$, $f_n = \text{incl}: A \hookrightarrow Y$
- For all $z \in Z_i$, $\pi_j(f, z): \pi_j(Z_i, z) \rightarrow \pi_j(Y, f_i(z))$ is
 - injective for $j = n$
 - bijective for $n < j < i$
 - surjective for $j = i$

Given this, we take $Z = \bigcup_{i \geq n} Z_i$ then (Z, A) is a relative CW-complex with cells of dimensions $\geq n + 1$, and $f = \bigcup_{i \geq n} f_i$ has the desired property.

(Z, A) is n -connected by cellular approximation

$$\begin{array}{ccc} f_*: \pi_j(Z, z) & \longrightarrow & \pi_j(Y, f(z)) \\ \uparrow & \nearrow \cong & \\ \pi_j(Z_{j+1}, z) & & (f_{j+1})_* \end{array}$$

where the left up map is an isomorphism by cellular approximation.

For the inductive Step A: make π_i injective

Step B: make π_{i+1} surjective.

Suppose $i \geq n$ and $Z_i \xrightarrow{f_i} Y$ have been constructed with the desired properties. For each path component C of A choose a basepoint $x_c \in A$ in that component. For each element in the kernel of $(f_i)_*: \pi_i(Z_i, x_c) \rightarrow \pi_i(Y, f_i(x_c))$ choose a based continuous map $q_{c,u}: S^i \rightarrow Z_i$, s.t. $[g_{c,u}] = u$. Define Z'_{i+1} as the pushout

$$\begin{array}{ccc} \coprod_{C,u \in \text{Ker}(f_{i*})} S^i & \xrightarrow{\coprod q_{c,u}} & Z_i \xrightarrow{f_i} Y \\ \downarrow & & \downarrow \quad \nearrow f_{i+1} \\ \coprod_{C,u} D^{i+1} & \longrightarrow & Z'_{i+1} \end{array}$$

For $j \leq i+1$ this is bijective for $j < i$, surjective for $j = i$ by cellular approximation. I missed a bit here.

This was step A, now comes step B.

For each $C \in \pi_0(A)$ and each element v of $\pi_{i+1}(Y, f(x_c))$ choose a representation $q_{c,v}: S^{i+1} \rightarrow Y$ s.t. $[q_{c,v}] = v$. Define $Z_{i+1} = Z'_{i+1} \vee \coprod_{\substack{C \in \pi_0 A \\ v \in \pi_{i+1}(Y, f(x_c))}} S^{i+1} \xrightarrow{f_{i+1} = f'_{i+1} \vee q_{c,v}} Y$.

This now has all required properties, missed the diagram with explaining everything. \square

[27.10.2025, Lecture 5]

[29.10.2025, Lecture 6]

Was ill, might add in later.

[29.10.2025, Lecture 6]

[3.11.2025, Lecture 7]

We want to show, that for CW-complexes, Eilenberg-MacLane-spaces are unique up to homotopy.

Theorem 1.35

Let $n \geq 1$. Let Y be an $(n-1)$ -connected CW-complex. Let Z be a based space s.t. $\pi_m(Z, z) = 0$ for $m > n$. Then for every group-homomorphism

$$\Phi: \pi_n(Y, y) \rightarrow \pi_n(Z, z)$$

there is a based continuous map $f: Y \rightarrow Z$, s.t. $\pi_n(f) = \Phi$. and such an f is unique up to based homotopy. Equivalently: the map

$$\pi_n: [Y, Z]_* \rightarrow \text{Hom}_{\text{Grp}}(\pi_n(Y, y), \pi_n(Z, z))$$

is bijective.

Proof. If Y is based homotopy equivalent to Y' and the claim holds for Y' , then it holds for Y . Then we can assume wlog, that Y has one 0-cell and no cells in dimensions $1 \leq i \leq n-1$. So $Y_n = \bigvee_I D^n / S^{n-1}$.

Step 1 we construct a continuous map $f_n: Y_n \rightarrow Z$ such that the following commutes:

$$\begin{array}{ccc} \pi_n(Y_n, *) & \xrightarrow{(f_n)_*} & \pi_n(Z, z) \\ & \searrow \text{incl}_* \quad \nearrow \Phi & \\ & \pi_n(Y, y) & \end{array}$$

We choose a homeomorphism $S^n \cong D^n/\partial D^n$ and characteristic maps $\chi_i: D^n \rightarrow Y$ for all $i \in I$. then the $S^n \cong D^n/\partial D^n \xrightarrow{\chi_i} Y_n \hookrightarrow Y$ represent a class $[\chi_i] \in \pi_n(Y, y)$, so $\Phi[\chi_i] \in \pi_n(Z, z)$. Let $w_i: S^n \rightarrow Z$ be a based map such that $[w_i] = \Phi[\chi_i]$. We define

$$f_n: Y_n = \bigvee_{i \in I} D^n/\partial D^n \cong \bigvee_{i \in I} S^n \rightarrow Z$$

as w_i on the wedge summand indexed by i . Then the maps commute by construction on the classes $[\chi_i]$. Since the class $[\chi_i]$ for $i \in I$ generates the group $\pi_n(Y_n, *)$ (Hurewicz theorem). Since all the maps in the commutative diagram are group homomorphisms and commute on generators, the diagram commutes.

Step 2 We extend f_n continuously to $f_{n+1}: Y_{n+1} \rightarrow Z$.

Let J be an index set for the $(n+1)$ -cells of Y . For $j \in J$ let $\chi_j: D^{n+1} \rightarrow Y$ be a characteristic map for the j -th $n+1$ -cell. The associated attaching map is $\chi_j|_{S^n}: S^n \rightarrow Y_n \cong \bigvee_I D^n/\partial D^n \cong \bigvee_I S^n$. Since Y_n is path-connected $\chi_j|_{S^n}$ is homotopic to a based map. By the HEP for (D^{n+1}, S^n) , we can extend the homotopy between $\chi_j|_{S^n}$ and the based map $\alpha: S^n \rightarrow Y_n$ to the $(n+1)$ -cell. So

$$[S^n \xrightarrow{\alpha} Y_n \hookrightarrow Y] \in \pi_n(Y, y)$$

is the zero homotopy class, i.e. $[\alpha] \in \pi_n(Y_n, y)$ lies in the kernel of incl_* . By commutativity of the (way above) commutative diagram, we get $(f_n)_*[\alpha] = 0$ in $\pi_n(Z, z)$. So the composite

$$S^n \xrightarrow{\alpha} Y_n \xrightarrow{f_n} Z$$

is nullhomotopic. Hence also $S^n \xrightarrow{\chi_j|_{S^n}} Y_n \xrightarrow{f_n} Z$ is nullhomotopic. We choose a continuous extension $\beta_j: D^{n+1} \rightarrow Z$ of $\chi_j|_{S^n}$ and define $f_{n+1}: Y_{n+1} = (\bigvee_I S^n) \cup_{J \times S^n} J \times D^{n+1} \rightarrow Z$ by f_n on Y_n and by β_j on the j -th $(n+1)$ -cell.

Step 3 We apply the previous lemma to the relative CW-complex (Y, Y_{n+1}) of which all relative cells have dimension $n \geq 2$. Since $\pi_i(Z, z) = 0$ for all $i \geq n+1$, the map $f_{n+1}: Y_{n+1} \rightarrow Z$ extends to a map $f: Y \rightarrow Z$.

We have $\Phi = f_*$ since incl_* is surjective and we can cancel it on the left.

We still have to show uniqueness up to based homotopy. Let $f, f': Y \rightarrow Z$ be based continuous maps, such that $\pi_n(f) = \pi_n(f'): \pi_n(Y) \rightarrow \pi_n(Z)$. Since $Y_{n-1} = *$, they agree on Y_{n-1} . Let I be an index set as before for the n -cells, $\chi_i: D^n \rightarrow Y$ characteristic maps for the i -th n -cell. $[S^n \cong D^n/\partial D^n \xrightarrow{\chi_i} Y] \in \pi_n(Y, y)$ which implies $[f \circ \chi_i] = f_*[\chi_i] = f'_*[\chi_i] = [f' \circ \chi_i]$. So $f \circ \chi_i, f' \circ \chi_i: D^n/\partial D^n \cong S^n \rightarrow Z$ are based homotopic. Choose such a homotopy for each $i \in I$, and glue them into a homotopy H

$$f|_{Y_n} \sim f'|_{Y_n}: \bigvee_I S^n \rightarrow Z$$

We apply the lemma from last time to the relative CW-complex $(Y \times [0, 1], Y \times \{0\} \cup Y_n \times [0, 1] \cup Y \times \{1\})$ all of whose relative cells have dimension $\geq n+2$. So by the lemma, the map

$$f \cup H \cup f': Y \times \{0\} \cup Y_n \times [0, 1] \cup Y \times \{1\}$$

has a continuous extension $K: Y \times [0, 1] \rightarrow Z$. This map is a based homotopy from f to f' . \square

Corollary 1.36. Let $n \geq 1$, A a group, abelian if $n \geq 2$. Let $(X, \phi), (Y, \psi)$ be Eilenberg-MacLane spaces of type (A, n) . If X is a CW-complex, then there is a based continuous map $f: X \rightarrow Y$,

unique up to based homotopy, such that

$$\begin{array}{ccc} \pi_n(X, x) & \xrightarrow[\cong]{f_*} & \pi_n(Y, y) \\ & \searrow \phi \quad \swarrow \psi & \\ & A & \end{array}$$

commutes.

Moreover f is a weak homotopy equivalence, and a homotopy equivalence if Y is also CW.

1.5 Representativity of cohomology

Here A is abelian group and $n \geq 0$. The Aim of this subsection is a natural transformation of functors in Y

$$[Y, K(A, n)] \rightarrow H^n(Y; A)$$

that is an isomorphism for CW-complexes. We want to construct a group structure on $[Y, K(A, n)]$, the "Homotopy group structure on $K(A, n)$ ".

Our hypothesis are $n \geq 1$, A abelian group, $K(A, n)$ some EM-space of type (A, n) that is a CW-complex. Because A is abelian, the addition $A \times A \rightarrow A$, $(a, b) \mapsto a + b$ and inverse $A \rightarrow A$, $a \mapsto a^{-1}$ are group homomorphisms. We apply the previous theorem to $Y = K(A, n) \times K(A, n)$ (an EM-space of type $(A \times A, n)$) and $ZFK(A, n)$ and the momomorphism

$$\pi_n(K(A, n) \times K(A, n)) \cong \pi_n(K(A, n)) \times \pi_n(K(A, n)) \xrightarrow{\cong} A \times A \xrightarrow{+} A \xrightarrow{\cong} \pi_n(K(A, n))$$

So there is a continuous map $\mu: K(A, n) \times K(A, n) \rightarrow K(A, n)$ that induces the addition on π_n ¹⁴.

Similarly, the composite $\pi_n(K(A, n)) \cong A \xrightarrow{a \mapsto -a} A \cong \pi_n(K(A, n))$ is realized by a based continuous map $i: K(A, n) \rightarrow K(A, n)$ unique up to based homotopy.

μ is associative up to based homotopy:

$$\mu \circ (\mu \times \text{Id}), \mu \circ (\text{id} \times \mu): K(A, n)^3 \rightarrow K(A, n)$$

Bothe induce the same map on π_n , so by the theorem, $\mu \circ (\mu \times \text{Id}) \sim \mu \circ (\text{id} \times \mu)$, i.e. μ is homotopy associative.

Let $\tau: K(A, n) \times K(A, n) \rightarrow K(A, n) \times K(A, n)$ be the flip map $\tau(x, y) = (y, x)$. Then $\mu \circ \tau, \mu: K(A, n) \times K(A, n) \rightarrow K(A, n)$ are by applying the theorem homotopic, i.e. μ is homotopy commutative.

Since the constant map $*$: $K(A, n) \rightarrow K(A, n)$ to the basepoint is 0 on π_n , the theorem shows that $\mu \circ (\text{id}, i) = *$. So $i: K(A, n) \rightarrow K(A, n)$ is a homotopy inverse for μ .

Upshot. $K(A, n)$ is a "homotopy abelian group".

Remark. It is possible to even take $K(A, n)$ as a topological abelian group. E.g. $|\tilde{A}[\Delta^n / \partial \Delta^n]|$.

Construction 1.37. For every (compactly generated) space Y , the set of homotopy classes of maps $[Y, K(A, n)]$ becomes an abelian group via

$$+ : [Y, K(A, n)] \times [Y, K(A, n)] \rightarrow [Y, K(A, n)]$$

$$([f: Y \rightarrow K(A, n)], [g: Y \rightarrow K(A, n)]) \mapsto [Y \xrightarrow{(f, g)} K(A, n) \times K(A, n) \xrightarrow{\mu_*} [Y, K(A, n)]]$$

¹⁴some mumbling about uncountable A and compactly generated topology, retopologizing $K(A, n) \times K(A, n)$

The homotopy associativity, homotopy commutativity and homotopy inverse properties of (μ, i) imply that $+$ is associative, commutative, and has an inverse $-[f] = [i \circ f]$.

[3.11.2025, Lecture 7]
[05.11.2025, Lecture 8]

Example 1.38. We have in this class not talked about Vector- and Line-bundles. Schwede encourages us to do so. $\mathbb{R}P^\infty$ and $\mathbb{C}P^\infty$ are EM-spaces of type $(\mathbb{Z}, 1)$ and $(\mathbb{Z}, 2)$ respectively, so they have "homotopy group structures".

So we get abelian group structures on $[Y, \mathbb{R}P^\infty]$, and $[Y, \mathbb{C}P^\infty]$.

Also $\mathbb{R}P^\infty$ and $\mathbb{C}P^\infty$ classify real/complex line bundles for compact spaces Y ,

$$[Y, \mathbb{R}P^\infty] \xrightarrow{\cong} \text{Pic}_{\mathbb{R}}(Y) = \text{groups under } \otimes \text{ of isoclasses of line bundles over } Y$$

$$[f: Y \rightarrow \mathbb{R}P^\infty \mapsto f^*(\gamma)]$$

That is an iso of abelian groups $\mu^*(\gamma) \cong p_1^*(\gamma) \otimes p_2^*(\gamma)$ as line bundles over $\mathbb{R}P^\infty \times \mathbb{R}P^\infty$.

Similarly $[Y, \mathbb{C}P^\infty] \cong \text{Pic}_{\mathbb{C}}(Y)$.

Example 1.39. A strictly associative & commutative model for $K(\mathbb{Z}/2, 1)$ and $K(\mathbb{Z}/2, 2)$.

The ?? as polynomial algebra on $\mathbb{C}[x]$ has no zero-divisors. So multiplication restricts to a commutative and associative operation

- $(\mathbb{C}[x] \setminus \{0\}) \times (\mathbb{C}[x] \setminus \{0\}) \rightarrow \mathbb{C}[x] \setminus \{0\}$ For $z \in \mathbb{C} \setminus 0$ $z(fg) = (zf)g = f(zg)$ for $f, g \in \mathbb{C}[x]$
So this descends to a well defined map

$$(\mathbb{C}[x] \setminus \{0\})/\mathbb{C}^* \times \mathbb{C}[x] \setminus \{0\}/\mathbb{C}^* \rightarrow (\mathbb{C}[x] \setminus \{0\})/\mathbb{C}^*$$

which then is a multiplication on $\mathbb{C}P^\infty$. This structure is associative and commutative, but not invertible.

Lemma 1.40: Fundamental class (non-standard-notation)

Let (X, ϕ) be an EM-space of type (A, n) . Then there is a unique class $\iota = \iota_{A, n} \in H^n(X, A)$ such that the composite

$$\pi_n(X, *) \xrightarrow{\text{Hurewicz}} H_n(X, \mathbb{Z}) \xrightarrow{\Phi(\iota)} A$$

equals $\phi: \pi_n(X, x) \xrightarrow{\cong} A$. Where we have

$$\Phi: H^n(X, A) \rightarrow \text{Hom}(H_n(X, \mathbb{Z}), A)$$

is the map from UCT.

ι is called the fundamental class.

Proof. Since X is $(n-1)$ -connected and A abelian, the Hurewicz homomorphism $\pi_n(X, *) \rightarrow H_n(X, \mathbb{Z})$ is an isomorphism. So

$$\phi \circ h^{-1}: H_n(X, \mathbb{Z}) \xrightarrow{\cong} A \in \text{Hom}(H_n(X, \mathbb{Z}), A)$$

Since X is $(n-1)$ -connected, the $\Phi: H^n(X, A) \xrightarrow{\cong} \text{Hom}(H_n(X, \mathbb{Z}), A)$ is an isomorphism and $\iota \mapsto \phi \circ h^{-1}$. \square

Construction 1.41. An EM-space of type $(A, 0)$ is the group A with the discrete topology and $\iota_{A, 0} \in H^0(A, A)$ is the class represented by $\text{id}: A \rightarrow A$.

Theorem 1.42: Group homomorphism

Let $n \geq 0$, A any abelian group. Then for all (compactly generated) Y , the map $[Y, K(A, n)] \rightarrow H^n(Y, A)$, $[f: Y \rightarrow K(A, n)] \mapsto f^*(\iota_{A, n})$ is an homomorphism of groups.

Proof. In 2 steps

Step 1 The universal example: $Y = K(A, n) \times K(A, n)$ and $f = \mu: K(A, n)^2 \rightarrow K(A, n)$. Let $p_1, p_2: K(A, n)^2 \rightarrow K(A, n)$ denote the two projections. Then

$$[p_1] + [p_2] = [K(A, n)^2 \xrightarrow[\text{id}]{(p_1, p_2)} K(A, n)^2 \xrightarrow{\mu} K(A, n)]$$

So we need to show that $\mu^*(\iota) = p_1^*(\iota) + p_2^*(\iota)$ in $H^n(K(A, n) \times K(A, n), A)$. By UCT and Hurewicz

$$\begin{aligned} H^n(K(A, n) \times K(A, n), A) &\stackrel{\Phi}{\cong} \text{Hom}(H_n(K(A, n) \times K(A, n), \mathbb{Z}), A) \\ &\stackrel{\text{Hurewicz}}{\cong} \text{Hom}(\pi_n(K(A, n), *)^2, A) \cong \text{Hom}(A \times A, A) \end{aligned}$$

The composit isomorphism sends $\mu^*(\iota) \mapsto \text{Addition } A \times A \rightarrow A$ and $p_1^*(\iota) \mapsto p_1: A \times A \rightarrow A$ and $p_2^*(\iota) \mapsto p_2: A \times A \rightarrow A$. In $\text{Hom}(A \times A, A)$ the relation $\text{Addition} = p_1 + p_2$ holds.

Step 2 General case: Let Y be a compactly generated space, $f, g: Y \rightarrow K(A, n)$. Then

$$\begin{aligned} ([f] + [g])^*(\iota) &= [\mu \circ (f, g)]^*(\iota) = (f, g)^*(\mu^*(\iota)) \\ &\stackrel{\text{step 1}}{=} (f, g)^*(p_1^*(\iota) + p_2^*(\iota)) \\ &= (f, g)^*(p_1^*(\iota)) + (f, g)^*(p_2^*(\iota)) \\ &= (p_1 \circ (f, g))^*(\iota) + (p_2 \circ (f, g))^*(\iota) = f^*(\iota) + g^*(\iota) \end{aligned}$$

□

Lemma 1.43

Let $n \geq 1$, A an abelian group. Y an based CW-complex. Then the forgetful map

$$[Y, K(A, n)_*] \rightarrow [Y, K(A, n)]$$

is bijective.

Proof. Earlier we have done an exercise: Y is a non-degenerately¹⁵ based space, Z any based space. Then the forgetful map

$$[Y, Z]_* \rightarrow [Y, Z]$$

is surjective if Z is path-connected and bijective if Z is simply connected.

This implies the lemma for $n \geq 2$ and surjectivity for $n = 1$. Injectivity for $n = 1$: Let $f, g: Y \rightarrow K(A, 1)$ be freely homotopic. Then $f_* = g_*: H_1(Y, \mathbb{Z}) \rightarrow H_1(K(A, 1), \mathbb{Z})$. Since A is abelian, $\pi_1(K(A, 1), *) \xrightarrow{\cong} H_1(K(A, 1), \mathbb{Z})$. So $\pi_1(f) = \pi_1(g)$. So by the earlier theorem 1.35, f and g are based homotopic. □

¹⁵inclusion of base point has HEP

Theorem 1.44

Let $n \geq 0$, A an abelian group. Then for all CW-complexes Y , the homomorphism

$$[Y, K(A, n)] \mapsto H^n(Y, A), \quad [f] \mapsto f^*(\iota_{A, n})$$

is an isomorphism.

Proof. Both $[_, K(A, n)]$ and $H^n(_, A)$ takes disjoint unions in Y to products of abelian groups. Every CW-complex Y is the disjoint union of its path-components. So we can assume wlog, that Y is path-connected.

Induction on n . For $n = 0$ $[Y, K(A, 0)] = [Y, A^{\text{disc}}] \cong A$ and also $H^0(Y, A) = \text{Hom}(\pi_0(Y), A) \cong A$. We do not check, that the map is indeed the identity, which we get between these two.

Now for $n \geq 1$. By the lemma we can replace free homotopy classes by based homotopy classes. We first treat a special case:

Special Case Y is $n - 1$ -connected. Then $H_{n-1}(Y, \mathbb{Z})$ is trivial for $n \geq 2$ or free $n = 1$, so $\text{Ext}(H_{n-1}(Y, \mathbb{Z}), A) = 0$. So the UCT provides an isomorphism

$$\Phi: H^n(Y, A) \rightarrow \text{Hom}(H_n(Y, \mathbb{Z}), A)$$

also the Hurewicz map $\pi_n(Y, *) \rightarrow H_n(Y, \mathbb{Z})$ is an isomorphism for $n \geq 2$ or the universal homomorphism into an abelian group for $n = 1$. In any case

$$h^*: \text{Hom}(H_n(Y, \mathbb{Z}), A) \xrightarrow{\cong} \text{Hom}(\pi_n(Y, *), A)$$

is bijective. The composite

$$[Y, K(A, n)]_* \xrightarrow{[f] \mapsto f^*(\iota)} H^n(Y, A) \xrightarrow[\cong]{\Phi} \text{Hom}(H_n(Y, \mathbb{Z}), A) \xrightarrow[h^*]{\cong} \text{Hom}(\pi_n(Y, *), A)$$

sends f to $\phi \circ \pi_n(f): \pi_n(Y, *) \rightarrow A$. So again by theorem 1.35, the composite is bijective. Also Φ and h^* are bijective. So the first map is bijective

General case We consider $Y \cup_{Y_{n-1}} C(Y_{n-1}) = Y \cup_{Y_{n-1} < 0} Y_{n-1} \times [0, 1] \cup_{Y_{n-1} \times 1} \{*\}$. This is $n - 1$ -connected. We consider the following commutative diagram:

$$\begin{array}{ccccccc} [\Sigma Y_{n-1}, K(A, n)]_* & \xrightarrow{p^*} & [Y \cup_{Y_{n-1}} CY_{n-1}, K(A, n)]_* & \xrightarrow{\iota^*} & [Y, K(A, n)]_* & \longrightarrow & 0 \\ \downarrow & & \downarrow \text{(special case)} \cong & & \downarrow & & \\ H^n(\Sigma Y_{n-1}, A) & \xrightarrow{p^*} & H^n(Y \cup_{Y_{n-1}} CY_{n-1}, A) & \longrightarrow & H^n(Y, A) & \longrightarrow & 0 \end{array}$$

Here i and p are inclusion and projection $p: Y \cup_{Y_{n-1}} CY_{n-1} \rightarrow \Sigma Y_{n-1}$.

Claim. The upper row is exact:

Surjectivity of i^* : all relative cells of $(Y \cup_{Y_{n-1}} CY_{n-1}, Y)$ have relative cells of dimension at most n . But $\pi_i(K(A, n), *) = 0$ for all $i < n$. By a previous "extension lemma", every continuous map $Y \rightarrow K(A, n)$ admits a continuous extension to $Y \cup_{Y_{n-1}} CY_{n-1}$. Since $p \circ i$ is nullhomotopic, $i^* \circ p^* = 0$. Suppose $f: Y \cup_{Y_{n-1}} CY_{n-1} \rightarrow K(A, n)$ is based s.t. $i^*[f] = 0$. So $f|_Y: Y \rightarrow K(A, n)$ is based nullhomotopic. The HEP of the pair $(Y \cup_{Y_{n-1}} CY_{n-1}, Y)$ lets us replace f by a based homotopic map f' , s.t. $f'|_Y = \text{const}_*$. So $f' ??$

The lower sequence is also exact by the l.e.s. for a CW pair $(Y \cup_{Y_{n-1}} CY_{n-1}, Y)$.

The left map in the diagram is surjective. We let $\rho: K(A, n-1) \rightarrow \Omega K(A, n)$ be the unique up to based homotopy map that induces

$$\pi_{n-1}(K(A, n-1)) \xrightarrow{\rho_*} \pi_{n-1}(\Omega K(A, n)) \cong \pi_n(K(A, n)) \cong A$$

Such that the composition is ϕ_{n-1} .

Let $\kappa_n: \Sigma K(A, n-1) \rightarrow K(A, n)$ be the adjoint of ρ .

Then the following commutes: I did not manage to copy the diagram.

So the right vertical map is an isomorphism by the 5-lemma.

□

[05.11.2025, Lecture 8]
[10.11.2025, Lecture 9]

Schwede remarks that when he was a student someone cleaned the board for the professor.

Example 1.45. If you have already heard about vector bundles somewhere:

$K(\mathbb{Z}/2, 1) = \mathbb{R}P^\infty$. For X a CW-complex, Also $\text{Pic}(X) \xleftarrow{\cong} [X, \mathbb{R}P^\infty] \xrightarrow{\cong} H^1(X, \mathbb{Z}/2)$. $[\gamma: E \rightarrow X] \mapsto w_1(\gamma)$ is called the first Stiefel-Whitney class. So real line bundles are completely classified by their first Stiefel-Whitney class.

Example 1.46. $K(\mathbb{Z}, 2) = \mathbb{C}P^\infty$. Then

$$\text{Pic}^\mathbb{C}(X) \xleftarrow{\cong} [X, \mathbb{C}P^\infty] \xrightarrow{\cong} H^2(X, \mathbb{Z})$$

so complex line bundles $[\gamma: E \rightarrow X] \mapsto c_1(\gamma)$ are classified by their first Chern class.

1.6 Cohomology operations and the Steenrod algebra

Aim. Define "Steenrod squares", natural homomorphisms

$$\text{Sq}^i: H^n(X, \mathbb{F}_2) \rightarrow H^{n+i}(X, \mathbb{F}_2)$$

together with properties and applications.

Schwede will follow some notes of his own which are not yet finalized and he will eventually publish on his website.

Definition 1.47: Cohomology operation

Let A, B be abelian groups. A *cohomology operation* of type (A, n, B, m) is a natural transformation

$$\tau = \{\tau_X: H^n(X, A) \rightarrow H^m(X, B)\}$$

of set valued functors on topological spaces. Specifically these operations need not be group homomorphisms.

We write

$$\text{Oper}(A, n, B, m) = \text{abelian group under pointwise addition of all such operations.}$$

This is not standard notation.

Lemma 1.48

The map $\text{Oper}(A, n, B, m) \rightarrow H^m(K(A, n), B)$ given by $\tau \mapsto \tau_{K(A, n)}(\iota_{A, n})$ is an isomorphism of groups.

Proof. On the homotopy category of CW-complexes, the pair $(K(A, n), \iota_{A, n})$ represents the functor $H^n(_, A)$. For any other functor $F: Ho(\text{CW-complexes}) \rightarrow \mathbf{Sets}$ we have

$$\text{Nat}(H^n(_, A), F) \xrightarrow{\cong} F(K(A, n))$$

by the Yoneda lemma. Apply this to $F = H^n(_, B)$ yields a bijection

$$\text{Nat}_{Ho\text{CW} \rightarrow \mathbf{Sets}}(H^n(_, A), H^m(_, B)) \cong H^m(K(A, n), B)$$

CW-approximation $X^{CW} \xrightarrow{\sim} X$ of spaces X are natural and unique up to homotopy, and we are considering functors on **Top** which are homotopy invariant and take weak equivalences to isomorphisms.

$$\begin{array}{ccc} H^n(X^{CW}, A) & \xrightarrow{\tau_{X, CW}} & H^m(X^{CW}, B) \\ \downarrow \cong & & \downarrow \cong \\ H^n(X, A) & \xrightarrow{\tau_X} & H^m(X, B) \end{array}$$

So

$$\text{Nat}_{\mathbf{Top}}(H^n(_, A), H^m(_, B)) \xleftarrow{\cong} \text{Nat}_{Ho(\mathbf{Top})}(H^n(_, A), H^m(_, B)) \xrightarrow{\cong} \text{Nat}_{Ho(\mathbf{CW})}(H^n(_, A), H^m(_, B))$$

using weak homotopy equivalences and CW-approximation \square

Example 1.49. 1. $K(A, n)$ is simply connected, so $H^0(K(A, n), B) \cong B$ for $n \geq 1$ and

$$H^i(K(A, n), B) = 0 \text{ for } 1 \leq i \leq n-1.$$

so the only operations of type $(A, n, B, 0)$ are the constant functions with image in B .

For $1 \leq i \leq n-1$, $\text{Oper}(A, n, B, i) = \{b\}$

2. Let $f: A \rightarrow B$ be a group homomorphism. This induces a natural group homomorphism
- 3.

$$f_*: H^n(X, A) \rightarrow H^n(X, B)$$

This is a additive cohomology operation for all $n \geq 0$. So we have

$$\text{Hom}_{\text{grp}}(A, B) \rightarrow \text{Oper}(H^n(_, A), H^n(_, B)) \cong H^n(K(A, n), B) \xrightarrow[\text{UCT}]{\cong} \text{Hom}(H_n(K(A, n), \mathbb{Z}), B) \cong_{\text{Hurewicz}}$$

4. Bockstein Homomorphisms associated with a short exact sequence

$$0 \rightarrow B \rightarrow E \rightarrow A \rightarrow 0$$

of abelian groups are cohomology operations $\beta: H^n(X, A) \rightarrow H^{n+1}(X, B)$ for all $n \geq 0$.

β only depends on the equivalence class of the s.e.s. in $\text{Ext}(A, B)$. So we get

$$\text{Ext}(A, B) \rightarrow \text{Oper}(A, n, B, n+1) \xrightarrow{\cong} H^{n+1}(K(A, n), B)$$

where for $n \geq 2$ the homomorphism from UCT is a isomorphism.

These are all cohomology operations for $n \geq 2$.

5. The group $H_2(K(A, 1), \mathbb{Z})$ is not generally trivial. For a group G (not necessarily abelian)

$$H^2(K(G, 1), B) = \text{isomorphism classes of centric extensions, i.e.}$$

short exact sequences of groups

$$0 \rightarrow BE \rightarrow G \rightarrow 0$$

such that B is central in E . If G is abelian, then

$$\text{abelian extensions} = \text{Ext}(A, B) \rightarrow H^2(K(A, 1), B) = \text{Central extensions}$$

As an exercise we will see a sort of Bockstein that gives a cohomology operation from degree 1 to degree 2.

So far we have

$$\text{Oper}(A, n, B, i) \cong \begin{cases} B & i = 0 \\ 0 & 1 \leq i \leq n-1 \\ \text{Hom}(A, B) & i = n \\ \text{Ext}(A, B) & i = n+1 \geq 3 \end{cases}$$

6. Let R be a ring, $k \geq 0$. Then

$$H^n(X, R) \rightarrow H^{kn}(X, R) \quad x \mapsto x^k = x \cup \cdots \cup x$$

is a cohomology operation of type (R, n, R, kn) . This is typically not additive.

For example

$$H^n(K(\mathbb{F}_2, 1), \mathbb{F}_2) \cong \text{Oper}(\mathbb{F}_2, 1, \mathbb{F}_2, k) \in \{x \mapsto x^k\}$$

where this is equal to

$$H^n(\mathbb{R}P^\infty, \mathbb{F}_2)$$

and

$$H^*(\mathbb{R}P^\infty, \mathbb{F}_2) = \mathbb{F}_2[\iota_{\mathbb{F}_2, 1}]$$

So $\text{Oper}(\mathbb{F}_2, 1, \mathbb{F}_2, k) = \{0, x \mapsto x^k\}$.

We also calculated $H^*(\mathbb{C}P^\infty, \mathbb{Z}) = \mathbb{Z}[\iota_{\mathbb{Z}, 2}]$. So this gives

$$\text{Oper}(\mathbb{Z}, 2, \mathbb{Z}, k) = \begin{cases} 0 & k \text{ odd} \\ \mathbb{Z}\{x \mapsto x^m\} & k = 2m \end{cases}$$

7. $\text{Oper}(\mathbb{Z}, 1, B, n) = ?$ We need to compute the cohomology of $K(\mathbb{Z}, 1) = S^1$.

$$H^n(S^1, B) = \begin{cases} B & n = 0, 1 \\ 0 & \text{else} \end{cases}$$

This is now the operations.

This completes the operations we can really calculate.

Reminder Let R be a commutative ring. We discussed

$$\cup_1 : C^n(X, R) \otimes C^m(X, R) \rightarrow C^{n+m-1}(X, R)$$

with coboundary formula $\delta(f \cup_1 g) = (\delta f) \cup_1 g + (-1)^n f \cup_1 (\delta g) - (-1)^{n+m} f \cup g - (-1)^{(n+1)(m+1)} (g \cup f)$. This was how we proofed commutativity of the cup product. We can use this to produce new cohomology operations.

Suppose n is even and $f \in C^n(X, R)$ is a cocycle, $\delta f = 0$. Then $\delta(f \cup_1 f) = 0$, and the cohomology class $[f \cup_1 f] \in H^{2n-1}(X, R)$ only depends on the cohomology class of f . Same for n odd and $2 = 0$ in R .

So we get cohomology operations $Sq_1: H^n(X, R) \rightarrow H^{2n-1}(X, R)$ for n even and

$$Sq_1: H^n(X, R) \rightarrow H^{2n-1}(X, R/2)$$

for n odd. They are defined by $Sq_1[f] := [f \cup_1 f]$.

Preview. We will define Stable cohomology operations of type $(A, B, n) = \{\tau_i: H^i(_, A) \rightarrow H^{n+i}(_, B)\}$ and some compatibility with a suspensions isomorphism.

Definition 1.50: Reduced cohomology operation

A *reduced cohomology operation* of type (A, n, B, m) is a natural transformation $\tilde{H}^n(_, A) \rightarrow \tilde{H}^m(_, B)$ of set valued functors of based spaces.

Then $\text{redOper}(A, n, B, m) \cong \tilde{H}^m(K(A, n), B) = H^m(K(A, n), B) = \text{Oper}(A, n, B, m)$ for $m \geq 1$. And for $m = 0$ the constant operations induced by $B \setminus \{0\}$ vanish.

Construction 1.51. Let (X, ϕ) and (Y, ψ) be EM-spaces of type (A, n) and $(A, n+1)$, respectively for $n \geq 1$. By an earlier theorem, there is a unique-up-to-based homotopy map $\rho: X \rightarrow \Omega Y$, such that $\rho_*: \pi_n(X, *) \rightarrow \pi_n(\Omega Y, *)$ agrees with the composite

$$\pi_n(X, *) \xrightarrow{\phi} A \xrightarrow{\psi^{-1}} \pi_{n+1}(Y, *) \cong \pi_n(\Omega Y, *)$$

we have $S^n \wedge S^1 \cong S^{n+1}$ and by understanding $S^n = \mathbb{R}^n \cup \{\infty\}$, this isomorphism is by coordinates.

Further $\Sigma X = X \wedge S^1$ and $\Omega X = \text{maps}_*(S^1, X)$

We let $\varepsilon: \Sigma X \rightarrow Y$ be the adjoint of ρ .

Lemma 1.52

Let (X, ϕ) , (Y, ψ) be EM-spaces of type (A, n) and $(A, n+1)$ respectively.

1. The following commutes:

$$\begin{array}{ccccc} \pi_n(X, *) & \xrightarrow{\Sigma} & \pi_{n+1}(\Sigma X, *) & \xrightarrow{\varepsilon} & \pi_{n+1}(Y, *) \\ & \searrow \phi & & & \downarrow \psi \cong \\ & & & & A \end{array}$$

2. The fundamental class $\iota_{A,n} \in H^n(X, A)$, $\iota_{A,n+1} \in H^{n+1}(Y, A)$ fulfill

$$\varepsilon^*(\iota_{A,n+1}) = \Sigma(\iota_{A,n}) \in H^{n+1}(\Sigma X, A)$$

Proof. 1.

$$\begin{array}{ccc} \pi_n(X, *) & \xrightarrow{\Sigma} & \pi_{n+1}(\Sigma X, *) \\ \downarrow \rho_* & & \downarrow \varepsilon_* \\ \pi_n(\Omega Y, *) & \xrightarrow{\quad} & \pi_{n+1}(Y, *) \\ \downarrow \psi & \nearrow \cong & \\ A & & \end{array}$$

Commutates because ε is adjoint to ρ .

2. Equivalently $\Sigma^{-1}(\varepsilon^*(\iota_{A,n+1})) = \iota_{A,n}$. We show that $\Sigma^{-1}(\varepsilon^*(\iota_{A,n+1}))$ has the property that defines $\iota_{A,n}$:

$$\pi_n(X, *) \xrightarrow[\downarrow f]{\mapsto} f_*[(S^n)] \text{Hurewicz} H_n(X, \mathbb{Z}) \xrightarrow{\Sigma^{-1}(\varepsilon^*(\iota_{A,n+1})) \cap _} A$$

equals ϕ . We make our choices $[S^n] \in H_n(S^n, \mathbb{Z})$ consistent with suspension.

Then

$$\begin{aligned} \Sigma^{-1}(\varepsilon^*(\iota_{A,n+1})) \cap f_*[S^n] &= f^*(\Sigma^{-1}(\varepsilon^*(\iota_{A,n+1}))) \cap [S^n] \\ &= \Sigma^{-1}((\Sigma f)^*(\varepsilon^*(\iota_{A,n+1}))) \cap [S^n] \\ &= (\Sigma f)^*(\varepsilon^*(\iota_{A,n+1})) \cap \Sigma[S^n] \\ &= (\varepsilon \circ (\Sigma f))^*(\iota_{A,n+1}) \cap [S^{n+1}] \\ &= \iota_{A,n+1} \cap (\varepsilon \circ (\Sigma f))_*[S^{n+1}] \\ &= \iota_{A,n+1} \cap \text{Hur}(\varepsilon \circ (\Sigma f)) \\ &= \psi[e \circ (\Sigma f)] = \phi(f) \end{aligned}$$

□

Appendix

List of definitions

1.1	Relative Homotopy Groups	3
1.2	n -Connectedness	3
1.7	8
1.12	9
1.13	10
1.25	Relative Hurewicz map	16
1.31	18
1.33	19
1.47	Cohomology operation	26
1.50	Reduced cohomology operation	29

List of statements

1.3	Blakers Massey	4
1.4	1.14	5
1.8	Freudenthal suspension theorem	8
1.14	11
1.15	Hurewicz	11
1.21	Hurewicz	13
1.26	relative Hurewicz, (simply connected case)	16
1.28	Whitehead theorem	17
1.29	17
1.30	12.16 for Lück	17
1.32	Relative Hurewicz with π_1	18
1.34	19
1.35	20
1.40	Fundamental class (non-standard-notation)	23
1.42	Group homomorphism	24
1.43	24
1.44	25
1.48	26
1.52	29