

UNIVERSITÄT BONN

Notes for the lecture

Topology II

held by

Stefan Schwede

T_EXed by

Jan Malmström

SoSe 2025

Corrections and improvements

If you have corrections or improvements, contact me via (s94jmalm@uni-bonn.de).

Contents

Lecture	1
1 Cohomology	2
1.1 Last Term	2
1.2 Cup-product	2
1.3 Commutativity of the cup-product	6
1.4 Künneth theorem	10
1.4.1 The Eilenberg-Zilber-theorem	10
1.4.2 Commutativity of the cup-product revisited	19
1.4.3 Relation between Homs and Tensors	26
Appendix	A
List of definitions	B
List of statements	C
Index	D

Lecture

Chapter 1

Cohomology


[07.04.2025, Lecture 1]

1.1 Last Term

In last term, we discussed

- CW-complexes
- higher homotopy groups
- Whitehead theorem
- Singular homology
- cellular homology

In the very end, cohomology was started. Remember



1.2 Cup-product

Let X be a simplicial set, and R^1 a ring.

$$C^n(X, R) = \text{maps}(X_n, R)$$

is an abelian group under pointwise addition. There is a differential

$$d^n : C^n(X, R) \rightarrow C^{n+1}(X, R)$$

given by

$$d^n(f)(y) = \sum_{i=0}^{n+1} (-1)^i f(d_i^*(y))$$

with $f : X_n \rightarrow R, y \in X_{n+1}$

¹A ring is not necessarily commutative, but has a unit

Construction 1.1 (Cup product/Alexander Whitney map). The cup product/Alexander Whitney map

$$\cup: C^n(X, R) \times C^m(X, R) \rightarrow C^{m+n}(X, R)$$

with $n, m \geq 0$ is defined by

$$(f \cup g)(x) := f(d_{front}^*(x)) \cdot g(d_{back}^*(x))$$

with $f: X_n \rightarrow R, g: X_m \rightarrow R, x \in X_{n+m}$.

Where we use $[n+m] = \{0, 1, \dots, n+m\}$ and $d_{front}: [n] \rightarrow [n+m], d_{back}: [m] \rightarrow [n+m]$ are given by $d_{front}(i) = i, d_{back}(i) = n+i$. Note, that d_{front} and d_{back} respectively suppress in their notation n and m .

Theorem 1.2: fundamental properties of cup product

The cup-product satisfies the following properties.

1. The AW-map is biadditive and satisfies a boundary formula:

$$d(f \cup g) = (df) \cup g + (-1)^n f \cup (dg) \in C^{m+n+1}(X, R)$$

2. Associativity: For $h \in C^k(X, R), (f \cup g) \cup h = f \cup (g \cup h) \in C^{n+m+k}(X, R)$.

Let $1 \in C^0(X, R)$ be the constant function $1: X_0 \rightarrow R$ with value 1. Then $1 \cup f = f \cup 1 = f$.

3. Naturality: Let $\alpha: Y \rightarrow X$ be a morphism of simplicial sets. Then

$$\alpha^*(f \cup g) = \alpha^*(f) \cup \alpha^*(g), \quad \alpha^*(1) = 1.$$

where $\alpha^*: C^n(X, R) \rightarrow C^n(Y, R), f \mapsto f \circ \alpha_n$.

Proof.

1. We check some properties: Let $d_{front}: [n] \rightarrow [n+m], d_{back}: [m] \rightarrow [n+m]$ be as in the definition of \cup . Then

$$d_i \circ d_{front} = \begin{cases} d_{front} \circ d_i & 0 \leq i \leq n+1 \\ d_{front} & n+1 \leq i \leq n+m+1 \end{cases}$$

and

$$d_i \circ d_{back} = \begin{cases} d_{back} \circ d_i & 0 \leq i \leq n \\ d_{back} \circ d_{i-n} & n \leq i \leq n+m+1 \end{cases}$$

Note, that for $n+1$ and n respectively the cases are the same.

Now we calculate

$$\begin{aligned}
d(f \cup g)(x) &= \sum_{i=0}^{n+m+1} (-1)^i (f \cup g)(d_i^*(x)) \\
&= \sum_{i=0}^{n+m+1} (-1)^i \cdot f(d_{front}^*(x)) \cdot g(d_{back}^*(d_i^*(x))) \\
&= \sum_{i=0}^n (-1)^i \cdot f(d_{front}^*(d_i^*(x))) \cdot g(d_{back}^*(d_i^*(x))) + \sum_{j=1}^{m+1} (-1)^{n+j} \cdot f(d_{front}^*(d_{j+n}^*(x))) \cdot g(d_{back}^*(d_{j+n}^*(x))) \\
&= \sum_{i=0}^{n+1} (-1)^i \cdot f(d_i^*(d_{front}^*(x))) \cdot g(d_{back}^*(x)) + \sum_{j=0}^{m+1} (-1)^{n+j} f(d_{front}^*(x)) \cdot g(d_j^*(d_{back}^*(x))) \\
&= d(f)(d_{front}^*(x)) \cdot g(d_{back}^*(x)) + (-1)^n \cdot f(d_{front}^*(x)) \cdot d(g)(d_{back}^*(x)) \\
&= ((df) \cup g)(x) + (-1)^n \cdot (f \cup dg)(x) \\
&= ((df) \cup g + (-1)^n \cdot f \cup (dg))(x)
\end{aligned}$$

2. For $x \in X_{n+m+k}$ we see

$$\begin{aligned}
((f \cup g) \cup h)(x) &= (f \cup g)(d_{front}^*(x)) \cdot h(d_{back}^*(x)) \\
&= f(d_{front}^*(d_{front}^*(x))) \cdot g(d_{back}^*(d_{front}^*(x))) \cdot h(d_{back}^*(x)) \\
&= f(d_{front}^*(x)) \cdot g(d_{middle}^*(x)) \cdot h(d_{back}^*(x))
\end{aligned}$$

Note that we abuse that d_{front} suppresses the indices for which the map is the front map. We have in the last line

$$d_{front}: [n] \rightarrow [n+m+k], d_{middle}: [m] \rightarrow [n+m+k], d_{back}: [k] \rightarrow [n+m+k]$$

defined by

$$d_{front}(i) = i, d_{middle}(i) = n+i, d_{back}(i) = n+m+i$$

this is obviously associative in the inputs²

3. Naturality for $\alpha: Y \rightarrow X$ we see

$$\begin{aligned}
(\alpha^*(f \cup g))(y) &= (f \cup g)(\alpha_{n+m}(y)) \\
&= f(d_{front}^*(\alpha_{n+m}(y))) \cdot g(d_{back}^*(\alpha_{n+m}(y))) = f(\alpha_n(d_{front}^*(y))) \cdot g(\alpha_m(d_{back}^*(y))) \\
&= \alpha^*(f)(d_{front}^*(y)) \cdot \alpha^*(g)(d_{back}^*(y)) \\
&= (\alpha^*(f) \cup \alpha^*(g))(y).
\end{aligned}$$

□

²for Schwede at least.

Definition 1.3: Differential graded ring

A differential graded ring (dg-ring) is a cochain-complex $A = \{A^n, d^n\}_{n \in \mathbb{Z}}$ equipped with biadditive maps

$$\cdot : A^n \times A^m \rightarrow A^{n+m}, \quad n, m \in \mathbb{Z}$$

and a unit $1 \in A^0$, such that;

- \cdot is associative and has 1 as a unit element.
- the Leibniz rule holds:

$$d(a \cdot b) = (da) \cdot b + (-1)^n \cdot a \cdot (db)$$

with $a \in A^n, b \in A^m$.¹

¹The sign is somehow connected to a sign-rule I couldn't follow. The d moved past the a or something.

Example 1.4. Some Differential graded rings are:

- $C^*(X, R)$ for a simplicial set X and a ring R .
- De Rham complex of a smooth manifold.

Construction 1.5 (Cup-Product on cohomology). Let $A = (A^n, d, \cdot)$ be a dg-ring. We define a map

$$\cdot : H^n(A) \times H^m(A) \rightarrow H^{n+m}(A), \quad [a] \cdot [b] = [a \cdot b]$$

This is well defined:

$$d(a \cdot b) = \underset{=0}{(da)} \cdot b + (-1)^n \underset{=0}{a} \cdot (db) = 0$$

so $a \cdot b$ is a cycle and we can take its homology class. Let $x \in A^{n-1}$.

$$(a + dx) \cdot b = a \cdot b + (dx) \cdot b = a \cdot b + d(x \cdot b) = [(a + dx) \cdot b] = [a \cdot b]$$

so it only depends on the cohomology class of a , analogous for b .

The product on cohomology inherits associativity and unity with $1 = [1] \in H^0(A)$. We need to see 1 is a cocycle:

$$d(1) = d(1 \cdot 1) = (d1) \cdot 1 + (-1)^0 1 \cdot (d1) = 2 \cdot d(1)$$

and so $d(1) = 0$.

The cup product on the R -cohomology of a simplicial set X is the product induced by the cup product on $C^*(X, R)$ in $H^*(C(X, R)) = H^*(X, R)$.

Theorem 1.6: Properties of the cup-product on homology

Let X be a simplicial set and R a ring. Then

- The cup product on $H^*(X, R)$ is associative and unital, with unit the cohomology class of the constant function $1: X_0 \rightarrow R$.
- For a morphism of simplicial sets $\alpha: Y \rightarrow X$, the relation

$$\alpha^*([x] \cup [y]) = \alpha^*[X] \cup \alpha^*[y]$$

holds for all $[x] \in H^n(X, R), [y] \in H^m(X, R)$.

Remark 1.7. The cup product generalizes to relative cohomology: For A, B simplicial subsets of X . We have

$$C^m(X, A; R) = \{f: X_n \rightarrow R \mid f(A_n) = \{0\}\}$$

The relative cup product is the restriction of \cup on $C^*(X, R)$ to

$$C^m(X, A; R) \times C^m(X, B; R) \xrightarrow{\cup} C^{n+m}(X, A \cup B; R).$$

Let $x \in (A \cup B)_{n+m}$, then

$$(f \cup g)(x) = f(d_{front}^*(x)) \cdot g(d_{back}^*(x))$$

if $x \in A_{n+m}$ then $f(d_{front}^*(x)) = 0$ and analogous with B_{n+m} , anyways the product is 0.

This gives us biadditive well defined maps

$$\cup: H^n(X, A; R) \times H^n(X, B; R) \rightarrow H^{n+m}(X, A \cup B; R)$$

In particular for $A = B$ we get

$$\cup: H^n(X, A; R) \times H^n(X, A; R) \rightarrow H^{n+m}(X, A; R)$$

which is well defined and associative, but not unital anymore.

1.3 Commutativity of the cup-product

Theorem 1.8: Commutativity of the cup-product

Let X be a simplicial set and R a commutative ring. Then for all $[x] \in H^n(X, R); [y] \in H^m(X, R)$ the relation

$$[x] \cup [y] = (-1)^{n \cdot m} \cdot [y] \cup [x]$$

holds.

Schwede points out, that the easy way doesn't work. **Warning.** For $f \in C^n(X, R), g \in C^m(Y, R)$, then in general $f \cup g \neq (-1)^{n+m}(g \cup f)$ in $C^{n+m}(X, R)$. The commutativity is a property we only get on homology.

Construction 1.9. The \cup_1 -product (spoken Cup-one)

$$\cup_1: C^m(X, R) \times C^m(X, R) \rightarrow C^{m+m-1}(X, R)$$

is defined by

$$(f \cup_1 g)(x) = \sum_{i=0}^{n-1} (-1)^{(n-1) \cdot (m+1)} f((d_i^{out})^*(x)) \cdot g((d_i^{inner})^*(x))$$

for $f \in C^n, g \in C^m$ and $x \in X_{n+m-1}$.³ where $d_i^{out}: [n] \rightarrow [n+m-1], d_i^{inner}: [m] \rightarrow [n+m-1]$ are the unique monotone injective maps with images $\text{Im}(d_i^{out}) = \{0, \dots, i\} \cup \{i+m, \dots, n+m-1\}$ and $\text{Im}(d_i^{inner}) = \{i, \dots, i+m\}$.

³There are also \cup_i for $i \in \mathbb{N}$. However, they are quite messy and combinatorial.

Theorem 1.10: \cup_1 -Product

The \cup_1 -product satisfies the following formula

$$d(f \cup_1 g) = (df) \cup_1 g + (-1)^n \cdot f \cup_1 (dg) - (-1)^{n+m} (f \cup g) - (-1)^{n+1} m + 1 (g \cup f)$$

for $f \in C^n(X, R)$ and $g \in C^m(X, R)$.

Remark 1.11. What we want to see, is that $f \cup g$ and $g \cup f$ are not the same but rather homotopic, and \cup_1 witnesses that homotopy.

Proof. This theorem will not be proven, because it is quite messy. You should find a lecture-video for that. \square

Now suppose that f and g are cocycles, i.e. $df = 0$, $dg = 0$. Then

$$d(f \cup_1 g) = -(-1)^{n+m} (f \cup g) - (-1)^{(n+1)(m+1)} (g \cup f)$$

and we get

$$(-1)^{n+m+1} \cdot d(f \cup_1 g) = f \cup g - (-1)^{n \cdot m} (g \cup f)$$

and as such

$$0 = [(-1)^{n+m-1}] = [f] \cup [g] - (-1)^{n \cdot m} [g] \cup [f]$$

Remark 1.12. Last term we discussed the tensor product of two chain complexes (in an exercise):

$$(C \otimes D)_n = \bigoplus_{p+q=n} C_p \otimes D_q$$

and differential

$$d(x \otimes y) = (dx) \otimes y + (-1)^{|x|} \cdot x \otimes (dy)$$

Remark 1.13. Reinterpretation of $d(f \cup_1 g)$. The cup product yields a morphism of cochain complexes

$$C^*(X, R) \otimes C^*(X, R) \rightarrow C^*(X, R)$$

and we get a diagram

$$\begin{array}{ccc} x \otimes y & C^*(X, R) \otimes C^*(X, R) & \xrightarrow{\cup} C^*(X, R) \\ \downarrow & \downarrow & \searrow \cup \\ y \otimes x & C^*(X, R) \otimes C^*(X, R) & \end{array}$$

that does not commute, however it does so up to cochain homotopy and \cup_1 is exactly a cochain homotopy between the two maps.

[07.04.2025, Lecture 1]
[09.04.2025, Lecture 2]

Only with the definition of the cup-product we cannot calculate a lot yet. Some methods to compute cup-products are:

- directly from the definition
- cellular approximation of the diagonal (whatever that means, he gives a little intuition I failed to record.) (this might be used later)
- Group homology (one example later today, something for AT I)

- Poincaré duality (later this term)
- Analysis on smooth manifolds together with De Rahm Cohomology

The first two methods are not very practical.

Example 1.14. Let X be a discrete space, Then $\mathcal{S}(X)$ is a constant simplicial set. The chain complex has the form

$$\xrightarrow{0} \mathbb{Z}[X] \xrightarrow{=} \mathbb{Z}[X] \xrightarrow{0} \mathbb{Z}[X]$$

And so $H^n(X, R) = 0$ for $n \geq 0$. And only for $n = m = 0$ something nontrivial happens. for $f: X_0 \rightarrow R, g: X_0 \rightarrow R$, we have $(f \cup g)(x) = f(d_{front}^*(x)) \cdot g(d_{back}^*(x)) = f(x) \cdot g(x)$ and so the cup product is just pointwise multiplication in dimension 0.

More generally: $H^0(X, R) = \text{maps}(\pi_0(X), R)$ with \cup -prodcut pointwise multiplication

Example 1.15. Let G be a group: Define a category \underline{G}^4 wit one object $*$ and $\text{Hom}_{\underline{G}}(*, *) = G$. We then define

$$BG = N(\underline{G})$$

Where N is the Nerve-Functor $\mathbf{CAT} \rightarrow \mathbf{Sset}$. Then

$$(BG)_n = G^n, \quad d_i^*: G^n \rightarrow G^{n-1}(g_1, \dots, g_n) \mapsto \begin{cases} (g_2, \dots, g_n) & i = 0 \\ (g_1, \dots, g_i \circ g_{i+1}, \dots, g_n) & 1 \leq i \leq n-1 \\ (g_1, \dots, g_{n-1}) & i = n \end{cases}$$

And $s_i(g_1, \dots, g_n) = (g_1, \dots, g_i, 1, g_{i+1}, \dots, g_n)$.

The general case of this is too hard to calculate. We take $G = (\mathbb{F}_2, +)$ and $R = \mathbb{F}_2$ and we calculate $H^*(B\mathbb{F}_2, \mathbb{F}_2)$. We see

$$\begin{array}{ccccccc} C^0(BG, A) & \xrightarrow{d} & C^1(BG, A) & \xrightarrow{d} & C^2(BG, A) & \longrightarrow & \dots \\ \parallel & & \parallel & & \parallel & & \\ \text{maps}(\{1\}, A) & \xrightarrow{0} & \text{maps}(G, A) & \longrightarrow & \text{maps}(G^2, A) & & \\ \parallel & & & & & & \\ A & & (f: G \rightarrow A) & \longrightarrow & (df)(g, h) & & \end{array}$$

And the map is defined by

$$f(d_0^*(g, h)) - f(d_1^*(g, h)) + f(d_2^*(g, h)) = f(h) - f(g \cdot h) + f(g)$$

and

$$df = 0 \Leftrightarrow f(g, h) = f(g) + f(h)$$

\Rightarrow 1-cocycles are the group homomorphisms from G to A

$$H^1(BG, A) \cong \text{Hom}(G, A)$$

and for $G = (\mathbb{F}_2, +)$, $A = \mathbb{F}_2$

We define

$$0 \neq x := [\text{Id}_{\mathbb{F}_2}] \in H^1(B\mathbb{F}_2, \mathbb{F}_2).$$

⁴via geometric realization, these define interesting spaces, namely some (missed word)-MacLane spaces $M(G, 1)$, didn't catch it all

We will show that $x^n = x \cup \dots \cup x$ (n -times) $\in H^n(B\mathbb{F}_2, \mathbb{F}_2)$ is nonzero.

Proposition. $x^n \in H^n(B\mathbb{F}_2, \mathbb{F}_2)$ is represented by

$$f_n: (\mathbb{F}_2)^n \rightarrow \mathbb{F}_2, f_n(\lambda_1, \dots, \lambda_n) = \lambda_1 \cdot \dots \cdot \lambda_n = \begin{cases} 1 & \text{if } \lambda_1 = \lambda_2 = \dots = \lambda_n = 1 \\ 0 & \text{else} \end{cases}$$

Proof. By induction on n . We checked for $n = 1$. For $n \geq 2$ we have

$$\begin{aligned} x^n &= x^{n-1} \cup x = [f_{n-1}] \cup [\text{Id}_{\mathbb{F}_2}] \\ &= [f_{n-1} \cup \text{Id}] \end{aligned}$$

Then

$$\begin{aligned} (f_{n-1} \cup \text{Id})(\lambda_1, \dots, \lambda_n) &= f_{n-1}(d_{\text{front}}^*(\lambda_1, \dots, \lambda_n)) \cdot \text{Id}(d_{\text{back}}^*(\lambda_1, \dots, \lambda_n)) \\ &= f_{n-1}(\lambda_1, \dots, \lambda_{n-1}) \cdot \text{Id}(\lambda_n) \\ &= (\lambda_1 \cdot \dots \cdot \lambda_{n-1}) \cdot \lambda_n \end{aligned}$$

□

Claim: $x^n \neq 0$. In the UCT for cohomology we used the evaluation pair

$$\Phi: H^n(X, A) \rightarrow \text{Hom}(H_n(X; \mathbb{Z}); A), \quad [f_n: X_n \rightarrow A] \mapsto \left\{ \left[\sum b_i x_i \right] \mapsto \sum b_i f(x_i) \right\}$$

for $b_i \in \mathbb{Z}, x_i \in X_n$. We can slightly vary that for ring coefficients:

$$\Phi: H^n(X, R) \rightarrow \text{Hom}(H_n(X, R), R)$$

and $[f: X_n \rightarrow R] \mapsto \left\{ \left[\sum r_i \cdot x_i \right] \mapsto \sum r_i \cdot f(x_i) \right\}$ with $r_i \in R, x_i \in X_n$.

With $X = B\mathbb{F}_2, R = \mathbb{F}_2$, we consider

$$y := \sum_{(\lambda_1, \dots, \lambda_n) \in (\mathbb{F}_2)^n} 1(\lambda_1, \dots, \lambda_n) \in \mathbb{F}_2[(\mathbb{F}_2)^n] = \mathbb{F}_2[(B\mathbb{F}_2)_n]$$

Claim: y is an n -cycle in $C_*(B\mathbb{F}_2, \mathbb{F}_2)$.

$$\begin{aligned} dy &= \sum_{i=0, \dots, n} (-1)^i \cdot d_i^* \left(\sum_1 \cdot (\lambda_1, \dots, \lambda_n) \right) \\ &= \sum_{i=0, \dots, n} \underbrace{\sum_{(\lambda_1, \dots, \lambda_n) \in \mathbb{F}_2^n} (-1)^i \cdot d_i^*(\lambda_1, \dots, \lambda_n)}_{\text{cancel in pairs}} \\ &= 0 \end{aligned}$$

Now

$$d_0^*(0, \lambda_2, \dots, \lambda_n) = (\lambda_2, \dots, \lambda_n) = d_0^*(1, \lambda_2, \dots, \lambda_n)$$

So

$$\Phi(x^n): H_n(B\mathbb{F}_2, \mathbb{F}_2) \rightarrow \mathbb{F}_2$$

$$\Phi(x^n)[y] = \Phi[f_n] \left[\sum_{(\lambda_1, \dots, \lambda_n) \in \mathbb{F}_2^n} (\lambda_1, \dots, \lambda_n) \right] = \sum_{(\lambda_1, \dots, \lambda_n)} f_n(\lambda_1, \dots, \lambda_n) = \sum_{(\lambda_1, \dots, \lambda_n)} \lambda_1 \cdot \dots \cdot \lambda_n = 1 \neq 0$$

and $[y] \neq 0$ in $H_n(B\mathbb{F}_2, \mathbb{F}_2)$.

We will later see, that in fact $H^*(B\mathbb{F}_2; \mathbb{F}_2) = \mathbb{F}_2[X]$.

Remark. Let p be an odd prime. $H^*(B\mathbb{F}_p, \mathbb{F}_p) = ?$.

$$0 \neq x = [\text{Id}_{\mathbb{F}_p} \in H^1(B\mathbb{F}_p; \mathbb{F}_p)]$$

still makes sense, but now there are more scalars and

$$x^n = 0$$

for $n \geq 2$. The graded commutativity says:

$$x \cup x = (-1)^{1 \cdot 1} x \cup x = -x \cup x$$

so if R is commutative, $x \in H^n(X, R)$ and n is odd, then $2 \cdot (x \cup x) = 0$ in $H^{2n}(X, R)$. And then $2 \cdot x^2 = 0 \Rightarrow x^2 = 0$.

Define $h: \mathbb{F}_p \times \mathbb{F}_p \rightarrow \mathbb{F}_p$ by

$$h(i, j) = \begin{cases} 0 & \text{if } i + j < p \\ 1 & \text{if } i + j \geq p \end{cases}$$

where we write $\mathbb{F}_p = \{0, \dots, p-1\}$. Now $h \in C^2(B\mathbb{F}_p, \mathbb{F}_p)$. Fact: $dh = 0$ and $0 \neq y := [h] \in H^2(B\mathbb{F}_p, \mathbb{F}_p)$.

We then get (but do not prove)

$$H^*(B\mathbb{F}_p, \mathbb{F}_p) = \Lambda(x) \otimes \mathbb{F}_p[y]$$

and

$$H^{2n}(B\mathbb{F}_p, \mathbb{F}_p) = \mathbb{F}_p\{y^n\}, \quad H^{2n+1}(B\mathbb{F}_p, \mathbb{F}_p) = \mathbb{F}_p\{xy^n\}$$

1.4 Künneth theorem

The Künneth theorem is an algebraic relationship between $H_*(X, R)$, $H_*(Y, R)$ and $H_*(X \times Y, R)$ ⁵.

Here is a simplest version in homology with field coefficients:

Theorem 1.16: Künneth, simple version

Let X and Y be spaces and k a field. Then

$$H_n(X \times Y, k)$$

is natural isomorphic to

$$\bigoplus_{p+q=n} H_p(X, k) \otimes_k H_q(Y, k)$$

1.4.1 The Eilenberg-Zilber-theorem

Let A, B be simplicial abelian groups. Then we get two natural chain homotopy equivalences

$$\begin{array}{ccc} & \text{Eilenberg-Zilber-Map} & \\ & \curvearrowright & \\ C_*(A) \otimes C_*(B) & & C_*(A \otimes B) \\ & \curvearrowleft & \\ & \text{Alexander Whitney map} & \end{array}$$

⁵ H_*^* denotes, that Schwede was too lazy to write the statement for homology and cohomology separately

Definition 1.17: Simplicial abelian group

A *simplicial abelian group* is a functor $A: \Delta^{op} \rightarrow \mathbf{Ab.Groups}$.

Remark 1.18. Equivalently a simplicial abelian group is a collection of abelian groups A_n , and homomorphisms $\alpha^*: A_m \rightarrow A_n$ for all $\alpha: [n] \rightarrow [m]$ in Δ , s.t. $(\alpha \circ \beta)^* = \beta^* \circ \alpha^*$.

Equivalently a simplicial abelian group is a simplicial set endowed with abelian group structure on the sets of n -simplices, such that all α^* are homomorphisms.

Example 1.19. Let X be a simplicial set and A an abelian group. Then the composite

$$\Delta^{op} \xrightarrow{X} (\mathbf{Sets}) \xrightarrow{A[_]} (\mathbf{ab.grps})$$

$A[X]$

is a simplicial abelian group.

Construction 1.20. Let $A: \Delta^{op} \rightarrow (\mathbf{ab.grps})$ be a simplicial abelian group. Its *chain complex* $C_*(A)$ is the chain complex with $C_n(A) = A_n$ with differential

$$d: C_n(A) = A_n \rightarrow A_{n-1} = C_{n-1}(A), \quad d(a) = \sum_{i=0, \dots, n} (-1)^i d_i^*(a)$$

And one can easily check $d \circ d = 0$.

Note. The following commutes

$$\begin{array}{ccc} (\mathbf{Ssets}) & \xrightarrow{X \mapsto C_*(X,A)} & (\mathbf{Chains}) \\ & \searrow A[_] \quad \nearrow C_* & \\ & (\mathbf{s.ab.grps}) & \end{array}$$

Remark 1.21. The tensor product of chain complexes C, D is

$$(C \otimes D)_n := \bigoplus_{p+q=n} C_p \otimes D_q$$

with differential

$$d(x \otimes y) = (dx \otimes y) + (-1)^p x \otimes (dy)$$

for $x \in C_p, y \in D_q$.

We can also form the tensor product of simplicial abelian groups:

Definition 1.22: Tensor product of simplicial abelian groups

$A, B: \Delta^{op} \rightarrow (\mathbf{ab.grps})$ by

$$(A \otimes B)_n = A_n \otimes B_n, \quad \alpha^*: (A \otimes B)_n \rightarrow (A \otimes B)_m$$

for $\alpha: [m] \rightarrow [n]$ is defined as $\alpha^*(a \otimes b) = \alpha^*(a) \otimes \alpha^*(b)$ and we write $\alpha_{A \otimes B}^* := \alpha_A^* \otimes \alpha_B^*$. This can be equally described as the composite

$$\Delta^{op} \xrightarrow{(A,B)} (\mathbf{ab.grps}) \times (\mathbf{ab.grps}) \xrightarrow{\otimes} (\mathbf{ab.grps})$$

Warning. For $A, B \in (\mathbf{SAB}) = \text{simplicial abelian groups}$

$$C_*(A \otimes B) \neq C_*(A) \otimes C_*(B)$$

Also he did this in dimension n , but I lacked time to copy.

The Eilenberg-Zilber theorem is a natural pair of chain homotopy equivalences between these two.

Construction 1.23. Let A, B be simplicial chain groups. The *Alexander-Whitney map* is the chain map

$$AW : C_*(A \otimes B) \rightarrow C_*(A) \otimes C_*(B)$$

defined by

$$\begin{array}{ccc} C_n(A \otimes B) & \longrightarrow & \bigoplus_{p+q=n, p,q \geq 0} A_p \otimes B_q \\ \parallel & & \parallel \\ A_n \otimes B_n & & C_*(A) \otimes C_*(B) \\ AW_n(a \otimes b) = \sum_{p+q=n} d_{front}^*(a) \otimes d_{back}^*(b) \end{array}$$

Where $[p] \xrightarrow{d_{front}} [p+q] = [n] \xleftarrow{d_{back}} [q]$.

You may check for yourself, that this is a chain map, however Schwede didn't do that.

[09.04.2025, Lecture 2]
[14.04.2025, Lecture 3]

Remark. An example for a simplicial abelian group, that is not of the form

$$\Delta^{op} \xrightarrow{X} \mathbf{sets} \xrightarrow{A[-]} (\mathbf{ab.grps.})$$

is for any abelian group G the simplicial set BG , that also admits structure of a simplicial abelian group.

Remark 1.24 (Relation between AW-map and cup-product). For a simplicial set X and ring R ,

$$C^*(X, R) = \text{Hom}(C_*(X, \mathbb{Z}), R) = \text{Hom}(C_*(\mathbb{Z}[X]), R)$$

and $C^n(X, R) = \text{Hom}(C_n(X, \mathbb{Z}), R)$. If $\psi \in C^n(X, R)$ is a cocycle, i.e. $d(\psi) = 0$, then it extends to a chain map

$$\tilde{\psi} : C_*(\mathbb{Z}[X]) \rightarrow R[n]$$

where $R[n]$ is the complex with R in dimension n and 0 otherwise. and $\tilde{\psi}$ is ψ in dimension n and 0 otherwise.

For $f \in C^n(X, R), g \in C^m(X, R)$ cocycles, we have $f \cup g \in C^{n+m}(X, R)$. Then $f \tilde{\cup} g$ is the following composite

$$\begin{array}{ccccc} C_*(\mathbb{Z}[X]) & \xrightarrow{C_*(\mathbb{Z}[\text{diagonal}])} & C_*(\mathbb{Z}[X \times X]) & \cong & C_*(\mathbb{Z}[X] \otimes \mathbb{Z}[X]) \\ & & \searrow \text{AW} & & \\ C_*(\mathbb{Z}[X]) \otimes C_*(\mathbb{Z}[X]) & \xleftarrow{\tilde{f} \otimes \tilde{g}} R[n] \otimes R[m] & \xrightarrow{\text{mult}} & R[n+m] \end{array}$$

Definition 1.25: (p,q)-shuffle

A (p, q) -shuffle for $p, q \geq 0$ is a permutation σ of $\{0, 1, \dots, p+q-1\}$, such that the restriction of σ to $\{0, 1, \dots, p-1\}$ is monotone, and the restriction of σ to $\{p, \dots, p+q-1\}$ is monotone.

Remark. „Shuffles leave the first p elements in order and the last q elements in order.“

Example 1.26. The only $(p, 0)$ -shuffle or $(0, q)$ -shuffles are the identity.

There are precisely two $(1, 1)$ -shuffles, namely both permutations of $\{0, 1\}$.

$\sigma \in S_3$ given by $\sigma(0) = 0, \sigma(1) = 2, \sigma(2) = 1$ is not a $(2, 1)$ -shuffle, but it is a $(1, 2)$ -shuffle.

Remark 1.27. (p, q) -shuffles biject with p -element subsets of $\{0, 1, \dots, p+q-1\}$ by $\sigma \mapsto \{\sigma(0), \dots, \sigma(p)\}$ and also with q -element subsets of $\{0, 1, \dots, p+q-1\}$ by $\sigma \mapsto \{\sigma(p), \dots, \sigma(p+q-1)\}$.

This means $|(p, q)\text{-shuffles}| = \binom{p+q}{p} = \binom{p+q}{q}$.

Notation 1.28. Let σ be a (p, q) -shuffle. We write $\mu_i := \sigma(i-1)$ for $1 \leq i \leq p$ and $\nu_i := \sigma(p+i-1)$ for $1 \leq i \leq q$.

This means $0 \leq \mu_1 \leq \dots \leq \mu_p$ and $0 \leq \nu_1 \leq \dots \leq \nu_q \leq p+q-1$.

Definition 1.29: Eilenberg-Zilber map

Let A, B be simplicial abelian groups. The Eilenberg-Zilber map /shuffle map is

$$EZ: C_*(A) \otimes C_*(B) \rightarrow C_*(A \otimes B)$$

is the direct sum of the homomorphisms

$$\nabla_{p,q}: A_p \otimes B_q \rightarrow A_{p+q} \otimes B_{p+q}$$

given by

$$a \otimes b \mapsto a \nabla b := \sum_{\sigma: (p,q)\text{-shuffle}} \text{sgn}(\sigma) \cdot (s_{\nu_i} \circ \dots \circ s_{\nu_q})^*(a) \otimes (s_{\mu_1} \circ \dots \circ s_{\mu_p})^*(b)$$

Example 1.30. There is only one $(p, 0)$ -shuffle, the identity of $\{0, \dots, p-1\}$. Then $\mu_i = i-1$.

$$\nabla_{p,0}: A_p \otimes B_0 \rightarrow A_p \otimes B_p$$

is defined by

$$a \otimes b \mapsto a \nabla b = a \otimes (s_0 \circ \dots \circ s_{p-1})^*(b).$$

For $p = q = 1$ i didn't have the time to copy.

Schwede claims, that the Eilenberg-Zilber map is a chain map and he can't believe he actually did those calculations 4 years ago. He will not torture us, but you may watch the videos.

Theorem 1.31: Shuffle maps form a chain map

The shuffle maps $\nabla_{p,q}$ for varying $p, q \geq 0$ assemble into a chain map. Furthermore, for $a \in A_p, b \in B_q$

$$d(a \nabla b) = (da) \nabla b + (-1)^p a \nabla (db)$$

He specifies, that the calculation takes up 8 pages of his notes.

Theorem 1.32: Eilenberg-Zilber

Let A, B be simplicial abelian groups. Then the morphisms

$$\begin{array}{ccc} & \xrightarrow{\text{Eilenberg-Zilber}} & \\ C_*(A) \otimes C_*(B) & & C_*(A \otimes B) \\ & \xleftarrow{AW} & \end{array}$$

are mutually inverse natural chain homotopy equivalences.

Proof. A first method of proof would be explicit formulas for the chain homotopies $AW \circ EZ \sim \text{Id}$ and $EZ \circ AW \sim \text{Id}$. That is however infinitely annoying and we will not do this.

For the special case, where $A = \mathbb{Z}[X], B = \mathbb{Z}[Y]$ for simplicial sets X, Y we prove this via acyclic models. For that we need some category-theory:

Proposition 1.33 (Yoneda lemma). Let \mathcal{C} be a category and c an object of \mathcal{C} . Let $F: \mathcal{C} \rightarrow (\mathbf{sets})$ be a functor: Then the evaluation map

$$\text{Nat}_{\mathcal{C} \rightarrow \mathbf{sets}}(\mathcal{C}(c, _), F) \rightarrow F(c)$$

given by

$$(\tau: \mathcal{C}(c, _) \rightarrow F) \mapsto (\tau_c: \mathcal{C}(c, c) \rightarrow F(c))(\text{id}_c)$$

is bijective.

Equally: for every $x \in F(c)$, there is a unique natural transformation $\tau: (\mathcal{C}(c, _) \rightarrow F)$, such that $\tau_c(\text{id}_c) = x$.

Remark. A special case of this is

$$\text{Hom}_{\mathbf{sset}}(\Delta^n, X) \cong X_n, \quad (f: \Delta^n \rightarrow X) \mapsto f_n(\text{id}_{[n]}).$$

where $\Delta^n = \Delta(_, [n])$.

Proof. We show injectivity and surjectivity.

Injectivity Let $\tau: \mathcal{C}(c, _) \rightarrow F$ be any natural transformation. Let d be another object of \mathcal{C} , $f: c \rightarrow d$ any morphism. Then we have

$$\tau_d: \mathcal{C}(c, d) \rightarrow F(d)$$

and

$$\tau_d(f: c \rightarrow d) = \tau_d(\mathcal{C}(c, f)(\text{id}_c)) = F(f)(\tau_c(\text{id}_c))$$

where we use naturality of τ :

$$\begin{array}{ccc} \mathcal{C}(c, d) & \xrightarrow{\tau_d} & F(d) \\ \downarrow \mathcal{C}(c, g) & & \downarrow F(g) \\ \mathcal{C}(c, e) & \xrightarrow{\tau_e} & F(e) \end{array}$$

which implies the value of τ at $d, f: c \rightarrow d$ is determined by its value of (c, id_c) and the functoriality of F .

Surjectivity Let $y \in F(c)$ be given. For an object d of \mathcal{C} and morphism $f: c \rightarrow d$, we define

$$\tau_d: \mathcal{C}(c, d) \rightarrow F(d) \quad \tau_d(f) := F(f)(y).$$

We check $\tau_c(\text{id}_c) = F(\text{id}_c)(y) = y$. We need to check for naturality. Let $g: d \rightarrow e$ be another morphism. Then

$$\begin{aligned} F(g)(\tau_d(f)) &= F(g)(F(f)(y)) = F(g \circ f)(y) \\ &= \tau_e(g \circ f) = \tau_e(\mathcal{C}(c, g)(f)) \end{aligned}$$

□

Let \mathcal{C} be a category, c an object of \mathcal{C} . We define the functor $\mathbb{Z}[\mathcal{C}(c, _)]: \mathcal{C} \rightarrow (\mathbf{ab.grps.})$ as the composite

$$\mathcal{C} \xrightarrow{\mathcal{C}(c, _)} (\mathbf{sets}) \xrightarrow{\mathbb{Z}[_]} (\mathbf{ab.grps.}).$$

In particular, $\mathbb{Z}[\mathcal{C}(c, _)](d) = \mathbb{Z}[\mathcal{C}(c, d)]$.

Proposition (Additive Yoneda lemma). Let $c \in \text{ob}(\mathcal{C})$, $F: \mathcal{C} \rightarrow (\mathbf{ab.grps.})$ any functor. Then the evaluation map

$$\text{Nat}_{\mathcal{C} \rightarrow (\mathbf{ab.grps.})}(\mathbb{Z}[\mathcal{C}(c, _)], F) \rightarrow F(c)$$

is bijective. ($\tau: \mathbb{Z}[\mathcal{C}(c, _)] \rightarrow F) \mapsto \tau_c(1 \cdot \text{id}_c)$.

Proof. For varying objects d of \mathcal{C} , the bijections

$$\text{Hom}_{AB}(\mathbb{Z}[\mathcal{C}(c, d)], F(d)) \cong \text{Hom}_{\mathbf{sets}}(\mathcal{C}(c, d), F(d))$$

assemble into a bijection⁶

$$\text{Nat}_{\mathcal{C} \rightarrow \mathbf{Ab}}(\mathbb{Z}[\mathcal{C}(c, _)], F) \cong \text{Nat}_{\mathcal{C} \rightarrow \mathbf{sets}}(\mathcal{C}(c, _), F) \xrightarrow{\text{Yoneda}} F(c)$$

□

Definition 1.34: Representable functor

A functor $F: \mathcal{C} \rightarrow \mathbf{Ab}$ is representable if there is an object $c \in \mathcal{C}$ and a natural isomorphism $F \cong \mathbb{Z}[\mathcal{C}(c, _)]$

Note. Any isomorphism $F \cong \mathbb{Z}[\mathcal{C}(c, _)]$ is determined by the „universal element“ in $F(c)$.

Example 1.35. Let $\mathcal{C} = (\mathbf{ssets}) \times (\mathbf{ssets})$ be the product of two copies of the category of simplicial sets. Define $f: (\mathbf{ssets}) \times (\mathbf{ssets}) \rightarrow \mathbf{Ab}$ given by $F(X, Y) = \mathbb{Z}[X_p \times Y_q]$ for some $p, q \geq 0$. **Claim.** This functor is representable by (Δ^p, Δ^q) with natural isomorphisms.

$$(\mathbf{ssets} \times \mathbf{ssets})((\Delta^p, \Delta^q), (X, Y)) = \mathbf{sets}(\Delta^p, X) \times \mathbf{sets}(\Delta^q, Y) \cong X_p \times Y_q$$

Apply free abelian groups to get

$$\mathbb{Z}[(\mathbf{ssets} \times \mathbf{ssets})((\Delta^p, \Delta^q)(X, Y))] \cong \mathbb{Z}[X_p \times Y_q]$$

Notation 1.36. For $F: \mathcal{C} \rightarrow \mathbf{Chains}$ we write $F_n = (_)_n \circ F: \mathcal{C} \rightarrow \mathbf{Ab}$ as the composite.

$$\mathcal{C} \xrightarrow{F} \mathbf{Chains} \xrightarrow{(_)_n} \mathbf{Ab}$$

⁶I don't know why though.

and the second map sends $C = C(n, d_n)_{n \in \mathbb{Z}} \mapsto C_n$.

Theorem 1.37: Acyclic models

Let \mathcal{C} be a category, $F, G: \mathcal{C} \rightarrow \mathbf{Chains}_+$ = non-negative grade chain complexes. Let $\psi: F \rightarrow G$ be a natural transformation of functors. Suppose;

1. The transformation $\psi_0: F_0 \rightarrow G_0: \mathcal{C} \rightarrow \mathbf{Ab}$ is the zero natural transformation
2. For every $n \geq 1$, the functor $F_n: \mathcal{C} \rightarrow \mathbf{Ab}$ is isomorphic to a direct sum of representable functors, $\bigoplus_{i \in I} \mathbb{Z}[\mathcal{C}(c_i, _)]$ for some family $\{c_i\}_{i \in I}$ of \mathcal{C} -objects such that $H_n(G(c_i)) = 0$.

Then ψ is naturally chain nullhomotopic.

[14.04.2025, Lecture 3]
[16.04.2025, Lecture 4]

Proof. For $n \geq 0$, we will construct natural transformations

$$s_n: F_n \rightarrow G_{n+1}$$

of functors $\mathcal{C} \rightarrow \mathbf{Ab}$, such that

$$d_{n+1} \circ s_n + s_{n-1} \circ d_n = \psi_n \quad (*)$$

as natural transformations (i.e. they have the chain homotopy property).

The construction is by induction on n . We begin with $s_0 = 0$ and $s_{-1} = 0$. Suppose $n \geq 1$ and that s_0, \dots, s_{n-1} have been constructed satisfying (*). Then

$$d_n^G \circ (\psi_n - s_{n-1} \circ d_n^F) = d_n^G \circ \psi_n - d_n^G \circ s_{n-1} \circ d_n^F$$

as ψ is a chain map,

$$= \psi_{n-1} \circ d_n^F - d_n^G \circ s_{n-1} \circ d_n^F = (\psi_{n-1} - d_n^G \circ s_{n-1}) \circ d_n^F \stackrel{(*)}{=} s_{n-2} \circ d_{n-1}^F \circ d_n^F = 0.$$

So $\psi_n - s_{n-1} \circ d_n^F: F_n \rightarrow G_n$ takes values in cycles. By 2.,

$$f_n = \bigoplus_{i \in I} \mathbb{Z}[\mathcal{C}(c_i, _)]$$

for some set $\{c_i\}_{i \in I}$ of \mathcal{C} -objects, such that $H_n(G(c_i)) = 0$. Let $j \in I$, write

$$x_j \in F(c_j) = \bigoplus_{i \in I} \mathbb{Z}[\mathcal{C}(c_i, c_j)]$$

be the element $1 \cdot \text{id}_j$ in the j -th summand. Then

$$\psi_n^{c_j}(x_j) - s_{n-1}^{c_j}(d_n^{F, c_j}(x_j)) \in G_n(c_j)$$

is a cycle. Since $H_n(G(c_j)) = 0$, the class is a boundary in the complex $G(c_j)$.

Let $y_j \in G(c_j)_{n+1}$ be a element such that

$$d_{n+1}^{c_j}(y_j) = \psi_n^{c_j}(x_j) - s_{n-1}^{c_j}(d_n^{F, c_j}(x_j))$$

The additive Yoneda lemma provides a unique natural transformation

$$s_{n,j}: \mathbb{Z}[\mathcal{C}(c_j, _)] \rightarrow G_{n+1}$$

such that $s_{n,j}(x_j) = s_{n,j}^{c_j}(1 \cdot \text{id}_{c_j}) = y_j \in G_{n+1}(c_j)$.

We define the natural transformation

$$s_n: F_n = \bigoplus_{i \in I} \mathbb{Z}[\mathcal{C}(c_i, _)] \rightarrow G_{n+1}$$

as $s_n = \bigoplus_{j \in I} s_{n,j}$.

It suffices now to show, that $(*)$ holds on each summand $\mathbb{Z}[\mathcal{C}(c_j, _)]$. By the additive Yoneda lemma, there it suffices to check the relation on $1 \cdot \text{id}_{c_j}$, which holds by definition. \square

Remark. We only proved „half“ of the acyclic models theorem. The other half states:

Let \mathcal{C} and $F, G: \mathcal{C} \rightarrow \mathbf{Chains}_+$ be as before, satisfying 2.. Then any natural transformation $\psi_0: F_0 \rightarrow G_0$ can be extended to a natural transformation $\psi: F \rightarrow G$.

Now to actually prove the Eilenberg-Zilber-Theorem ?? (at least in a special case.) Let A, B be simplicial abelian groups. We assume $A = \mathbb{Z}[X]$, $B = \mathbb{Z}[Y]$ for some simplicial sets X, Y . We write $C_*(X), C_*(Y)$. For sets S, T ,

$$\begin{array}{ccc} & \curvearrowright & \\ \mathbb{Z}[S] \otimes \mathbb{Z}[T] & & \mathbb{Z}[S \times T] \\ & \curvearrowleft & \\ s \otimes t & \longrightarrow & (s, t) \end{array}$$

is naturally isomorphic. Dimensionwise this gives $\mathbb{Z}[X] \otimes \mathbb{Z}[Y] \cong \mathbb{Z}[X \times Y]$.

We want to move this further to $C_*(X) \otimes C_*(Y) \cong C_*(X \times Y)$.

Proposition 1.38.

1. For all $p \geq 0$, the simplicial set Δ^p is simplicially contractible.
2. For all $p \geq 0$, the complex $C_*(\Delta^p)$ is chain homotopy equivalent to the complex $\mathbb{Z}[0]$, the complex consisting of \mathbb{Z} in dimension 0.
3. For $p, q \geq 0$, the chain complex $C_*(\Delta^p) \otimes C_*(\Delta^q)$ is chain homotopy equivalent to $\mathbb{Z}[0]$. In particular,

$$H_n(C_*(\Delta^p) \otimes C_*(\Delta^q)) = 0$$

for $n > 0$.

Proof.

1. We define a morphism of simplicial sets $H: \Delta^p \times \Delta^1 \rightarrow \Delta^p$ that contracts Δ^p to the last vertex.⁷ In dimension n ,

$$H_n: \Delta([n], [p]) \times \Delta([n], [1]) \rightarrow \Delta([n], [p])$$

is given by

$$H_n(\alpha, \beta)(i) = \begin{cases} \alpha(i) & \text{if } \beta(i) = 0 \\ p & \text{if } \beta(i) = 1 \end{cases}$$

for $0 \leq i \leq n$. Let $\gamma: [m] \rightarrow [n]$ be any morphism in Δ . Then

$$H_m(\gamma^*(\alpha, \beta))(j) = H_m(\alpha \circ \gamma, \beta \circ \gamma)(j) = \begin{cases} \alpha(\gamma(j)) & \text{if } \beta(\gamma(j)) = 0 \\ p & \text{if } \beta(\gamma(j)) = 1 \end{cases} = H_n(\alpha, \beta)(\gamma(j)) = \gamma^*(H_n(\alpha, \beta))(j)$$

⁷remember, that Homotopy is not symmetric in Simplicial sets. This is such an example.

This means H is a homotopy from Id_{Δ^p} to the composite

$$\Delta^p \rightarrow \Delta^0 \xrightarrow{p\text{-th vertex}} \Delta^p$$

2. $C_*: \mathbf{ssets} \rightarrow \mathbf{chains}$ takes simplicial homotopies to chain homotopies. So we know $C_*(\Delta^p)$ is chain homotopy equivalent to

$$C_*(\Delta^0) = (\dots \mathbb{Z} \xrightarrow{\text{Id}} \mathbb{Z} \xrightarrow{0} \mathbb{Z})$$

which is chain homotopy equivalent to

$$(\dots 0 \rightarrow 0 \rightarrow 0 \rightarrow 0 \rightarrow \mathbb{Z}) = \mathbb{Z}[0]$$

3. The tensor product of chain complexes preserves chain homotopy equivalences in each variable separately. So

$$C_*(\Delta^p) \otimes C_*(\Delta^q) \sim \mathbb{Z}[0] \otimes C_*(\Delta^1) \sim \mathbb{Z}[0] \otimes \mathbb{Z}[0] \cong \mathbb{Z}[0].$$

□

We now must produce natural chain homotopies from

$$\mathbf{AW} \circ \mathbf{EZ}: C_*(X) \otimes C_*(Y) \rightarrow C_*(X) \otimes C_*(Y)$$

and

$$\mathbf{EZ} \circ \mathbf{AW}: C_*(X \times Y) \rightarrow C_*(X \times Y)$$

to the respective identities.

Claim. $\mathbf{AW} \circ \mathbf{EZ} - \text{Id}_{C_*(X) \otimes C_*(Y)}: C_*(X) \otimes C_*(Y) \rightarrow C_*(X) \otimes C_*(Y)$ satisfies the hypothesis of acyclic models.

Proof.

$$\begin{array}{ccc} C_0(X) \otimes C_0(Y) & \cong & \mathbb{Z}[X_0] \otimes \mathbb{Z}[Y_0] \\ \parallel & & \uparrow \cong \\ (C_*(X) \otimes C_*(Y))_0 & & \mathbb{Z}[X_0 \times Y_0] \\ & & \downarrow \cong \\ & & C_0(X \times Y) \end{array}$$

Which means $(\mathbf{AW} \circ \mathbf{EZ})_0 = \text{Id}$ and $(\mathbf{EZ} \circ \mathbf{AW})_0 = \text{Id}$. which means $\psi_0 = \text{zero natural transformation}$.

$$(C_*(X) \otimes C_*(Y))_n = \bigoplus_{p+q=n} C_p(X) \otimes C_q(Y) = \bigoplus_{p+q=n} \mathbb{Z}[X_p] \otimes \mathbb{Z}[Y_q] \cong \bigoplus_{p+q=n} \mathbb{Z}[X_p \times Y_q]$$

which is represented by (Δ^p, Δ^q) . Then $H_n(C_*(\Delta^p \otimes \Delta^q)) = 0$ (I think, he erased before I could copy.)

We consider $\phi: \mathbf{EZ} \circ \mathbf{AW} - \text{Id}_{C_*(X \times Y)}: C_*(X \times Y) \rightarrow C_*(X \times Y)$. We know, $\phi_0 = 0$. We need to show, that ϕ satisfies the hypothesis of acyclic models.

$$C_n(X \times Y) = \mathbb{Z}[X_n \times Y_n]$$

is representable by (Δ^n, Δ^n) .

$$H_n(C_*(\Delta^n \times \Delta^n)) \cong H_n(\Delta^0 \times \Delta^0) = H_n(\Delta^0) = 0$$

for $n > 0$, where we used $\Delta^n \sim \Delta^0$ and so $\Delta^n \times \Delta^n \sim \Delta^0 \times \Delta^0$. So acyclic models produces a natural chain nullhomotopy of ϕ . \square

This concludes the proof of the Künneth theorem. \square

1.4.2 Commutativity of the cup-product revisited

The symmetry isomorphism of chain complexes C, D is the morphism.

$$\tau_{C,D}: C \otimes D \xrightarrow{\cong} D \otimes C$$

is given by

$$\begin{aligned} \tau_{C,D_n} &: (C \otimes D)_n && (D \otimes C)_n \\ &\oplus_{p+q=n} C_p \otimes D_q && \oplus_{q+p=n} D_q \otimes C_p \\ &c \otimes d && (-1)^{pq} \cdot d \otimes c \end{aligned}$$

Fact.

$$\begin{array}{ccc} C_*(X) \otimes C_*(Y) & \xrightarrow{\mathbf{EZ}} & C_*(X, Y) \\ \downarrow \tau & & \downarrow C_*(flip) \\ C_*(Y) \otimes C_*(X) & \xrightarrow{\mathbf{EZ}} & C_*(Y \otimes X) \end{array}$$

commutes. where $flip: X \times Y \rightarrow Y \times X$, $(x, y) \mapsto (y, x)$. Hence, „The Eilenberg-Zilber map is symmetric“.

But however for AW the same diagram does NOT commute.

However it does so up to natural chain homotopy by applying the acyclic models to the difference of the two composites. He explains, why we can apply acyclic models.

Let X be a simplicial set. The diagonal $\Delta: X \rightarrow X \times X$ is flip-invariant, i.e.

$$\begin{array}{ccc} X & \xrightarrow{\Delta} & X \times X \\ & \searrow \Delta & \downarrow flip \\ & & X \times X \end{array}$$

We draw a diagram:

$$\begin{array}{ccccc} C_*(X) & \xrightarrow{C_*(\Delta)} & C_*(X \times X) & \xrightarrow{\mathbf{AW}} & C_*(X) \otimes C_*(X) \\ & \searrow C_*(\Delta) & \downarrow C_*(flip) & & \downarrow \tau \\ & & C_*(X \times X) & \xrightarrow{\mathbf{AW}} & C_*(X) \otimes C_*(X) \end{array}$$

that commutes up to homotopy. We apply the functor $\text{Hom}(_, R)$ to get a new diagram and my speed at copying was not capable of keeping up. You may want to have a look at the videos for this.

[14.04.2025, Lecture 4]
[23.04.2025, Lecture 5]

The Plan for today is to show the Künneth theorem for homology. The rough approximation is, that product of spaces goes to Tensorproducts of abelian groups.

If X, Y are simplicial sets, then by EZ we have $H_*(X \times Y; R) = H_*(C_*(X \times Y; R)) \cong H_*((C_*(X, R)) \otimes_R C_*(Y; R))$ and we want to see how that relates to $H_*(X, R) \otimes_R H_*(Y; R)$.

In the following R is a commutative ring (have integers and fields in mind).

Definition 1.39: Tensor Product of R -chains

Let C, D be chain complexes of R -modules. We define a new complex of R -modules $C \otimes_R D$:

$$(C \otimes_R D)_n = \bigoplus_{p+q=n} C_p \otimes_R D_q$$

with differential

$$d(x \otimes y) = dx \otimes y + (-1)^{pq} x \otimes dy.$$

Note that $R \otimes \mathbb{Z}[S] \cong R[S]$ for S a simplicial set. And $R[S] \otimes_R R[T] \cong R[S \times T]$ for S, T simplicial sets.

For X, Y simplicial sets, we have

$$R \otimes C_*(X, \mathbb{Z}) \otimes C_*(Y, \mathbb{Z}) \xrightarrow{R \otimes \mathbf{EZ}} R \otimes C_*(X \times Y; \mathbb{Z}) \cong C_*(X \otimes Y; R)$$

and for $R \otimes C_*(X; \mathbb{Z}) \otimes C_*(Y; \mathbb{Z}) \cong (R \otimes C_*(X; \mathbb{Z})) \otimes_R (R \otimes C_*(Y; \mathbb{Z})) = C_*(X, R) \otimes_R C_*(Y, R)$, so we get a Eilenberg-Zilber map

$$C_*(X, R) \otimes_R C_*(Y, R) \xrightarrow{\mathbf{EZ}} C_*(X \times Y; R)$$

Aim. relate $H_*(C \otimes_R D)$ to $H_*(C), H_*(D)$. Our hope is to have a map

$$\bigoplus_{p+q=n} H_p(C) \otimes_R H_q(D) \xrightarrow{???} H_n(C \otimes_R D)$$

For example taking $R = \mathbb{Z}$ and $C = D = (\dots, \rightarrow 0 \rightarrow \mathbb{Z} \xrightarrow{2} \mathbb{Z} \rightarrow 0)$. Then

$$H_n(C) = H_n(D) = \begin{cases} \mathbb{Z}/2 & n = 0 \\ 0 & n \neq 0 \end{cases}$$

but $C \otimes D = (0 \rightarrow 0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Z}^2 \rightarrow \mathbb{Z} \rightarrow 0)$. And

$$H_1(C \otimes D) = \{(x, -x) \in \mathbb{Z}\} / \{(2y, -2y) \mid y \in \mathbb{Z}\} \cong \mathbb{Z}/2 \neq 0$$

Definition 1.40: Projective R -modules

An R -module P is *projective* if for every epimorphism $\varepsilon: M \rightarrow N$ of R -modules, the map

$$\mathrm{Hom}(P, \varepsilon): \mathrm{Hom}(P, M) \rightarrow \mathrm{Hom}(P, N)$$

is surjective.

$$\begin{array}{ccc} & & M \\ & \nearrow & \downarrow \varepsilon \\ P & \xrightarrow{f} & N \end{array}$$

Fact. P is projective iff P is a direct summand of a free module iff there exists a R -module Q and a set S , such that

$$P \oplus Q \cong R[S].$$

Proof. Free modules are projective:

$$\begin{array}{ccc} & & M \\ & \nearrow g & \downarrow \varepsilon \\ R[S] & \xrightarrow{f} & N \end{array}$$

for every $s \in S$ choose $m_s \in M$ $\varepsilon(m_s) = f(s)$. Then there is a unique homomorphism $g: R[S] \rightarrow M$ such that $g(s) = m_s$.

Let P be projective and Q a summand of P . For reasons I couldn't copy, then Q is also projective.

Let P be a projective R -module. Consider the epimorphism

$$\begin{array}{ccc} R[P] & \rightarrow & P \\ p & \mapsto & p \end{array}$$

Then we have

$$\begin{array}{ccc} & & R[P] \\ & \nearrow g & \downarrow \\ p & \xrightarrow{\text{id}} & P \end{array}$$

So P is a direct summand of $R[P]$.

□

- If R is a field, then all modules are free, hence projective.
- $R = \mathbb{Z}/6$, $P = \mathbb{Z}/2$, $Q = \mathbb{Z}/3$. Then $\mathbb{Z}/6 \cong \mathbb{Z}/2 \oplus \mathbb{Z}/3$, so, as $\mathbb{Z}/6$ is free, $\mathbb{Z}/2$ and $\mathbb{Z}/3$ are projective, but not free.

Proposition 1.41. Let R be a commutative ring, and

$$0 \rightarrow I \xrightarrow{\alpha} M \xrightarrow{\beta} N \rightarrow 0$$

be a short exact sequence of R -modules.

Then for every R -module P , the sequence

$$P \otimes_R I \xrightarrow{P \otimes_R \alpha} P \otimes_R M \xrightarrow{P \otimes_R \beta} P \otimes_R N \rightarrow 0$$

is exact. („ $P \otimes_R _$ is right exact“). If moreover P is projective, then it is also exact with a 0 on the left, i.e. $P \otimes_R \alpha$ is injective. („projective modules are flat“).

Proof.

$$p \otimes_R \beta \circ (p \otimes_R \alpha) = P \otimes_R (\beta \circ \alpha) = P \otimes_R 0 = 0$$

so $\text{Im}(P \otimes_R \alpha) \subseteq \ker(P \otimes_R \beta)$ so we get an induced homomorphism

$$\gamma \frac{P \otimes_R M}{\text{Im}(P \otimes_R \alpha)} \rightarrow P \otimes_R N$$

exactness is equivalent to δ being an isomorphism. We define a homomorphism $\delta: P \otimes_R N \rightarrow \frac{P \otimes_R M}{\text{Im}(P \otimes_R \alpha)}$ given by $(p, n) \in P \otimes N$ choose $\tilde{n} \in M$, such that $\beta(\tilde{n}) = n$.

Claim. $\delta(p \otimes n) = p \otimes \tilde{n} + \text{Im}(P \otimes_R \alpha)$ is independent of choice of \tilde{n}

Proof. Let $\tilde{\tilde{n}} \in M$ also satisfy $\beta(\tilde{\tilde{n}}) = n$. Then $\beta(\tilde{\tilde{n}} - \tilde{n}) = 0$, so there is $i \in I$ s.t. $\alpha(i) = \tilde{\tilde{n}} - \tilde{n}$.
 $p \otimes \tilde{\tilde{n}} - p \otimes \tilde{n} = p \otimes (\tilde{\tilde{n}} - \tilde{n}) = p \otimes \alpha(i) \in \text{Im}(P \otimes_R \alpha)$. \square

Claim. The assignment of δ is biadditive and sends (rp, n) and (p, rn) to the same element.

Then this extends to a well defined R -linear map

$$P \otimes_R N \rightarrow \frac{P \otimes_R M}{\text{Im}(P \otimes_R \alpha)}$$

which is isomorphic.

Now let P be projective. We show that then $P \otimes_R \alpha$ is injective.

Case 1 $P = R[S]$ free, S some set. Then

$$P \otimes_R M = R[S] \otimes_R M \cong \bigoplus_{s \in S} s \in SM$$

we have a natural isomorphism of R -modules in M .

From this we get a commutative square of R -modules:

$$\begin{array}{ccc} P \otimes_R I & \xrightarrow{P \otimes_R \alpha} & P \otimes_R M \\ \parallel & & \parallel \\ \bigoplus_{s \in S} I & \xrightarrow{\bigoplus_{s \in S} \alpha} & \bigoplus_{s \in S} M \end{array}$$

where the bottom map is injective.

General case P projective is a summand of a free module F , i.e. there are homomorphisms

$$P \xrightarrow{\lambda} F \xrightarrow{\mu} P$$

s.t. $\mu \circ \lambda = \text{Id}_P$. We consider the commutative square

$$\begin{array}{ccc} P \otimes_R I & \xrightarrow{P \otimes_R \alpha} & P \otimes_R N \\ \downarrow \lambda \otimes_R I & & \downarrow \lambda \otimes_R N \\ F \otimes_R I & \xrightarrow{F \otimes_R \alpha} & F \otimes_R N \end{array}$$

where the bottom map is injective by Case 1 and $\lambda \otimes_R I$ is injective, as it admits a retraction.

\square

Definition 1.42: Global dimension of rings

A commutative ring R has global dimension ≤ 1 if every submodule of a projective module is projective.

Example 1.43. Some rings with global dimension ≤ 1 are

- fields
- the ring of integers \mathbb{Z} (subgroups of free abelian groups are free).
- every PID⁸ is of this form. See for example $k[x]$ for k a field or $\mathbb{Z}[i]$ the gaussian integers
- \mathbb{Z}_p the p -adic integers.

Definition 1.44: Tor of nice rings

Let R be a commutative ring of global dimension ≤ 1 . Let M, N be R -modules. Choose an epimorphism $p: P \rightarrow N$ of R -modules with P projective. Define

$$\mathrm{Tor}^R(M, N) = \mathrm{Ker}(M \otimes_R N \xrightarrow{M \otimes_R \mathrm{incl}} M \otimes_R P)$$

Facts. This is independent up to preferred isomorphism of the choice of $p: P \rightarrow N$.

It is symmetric, i.e. we can resolve M instead of N .

If P is projective, then $\mathrm{Tor}^R(P, N) = 0 = \mathrm{Tor}^R(M, P)$.

Construction 1.45. For R a commutative ring, C, D complexes of R -modules. We define a natural homomorphism

$$\Phi: H_p(C) \otimes_R H_q(D) \rightarrow H_{p+q}(C \otimes_R D)$$

via $[x] \otimes [y] \mapsto [x \otimes y]$

We can check this is well defined.

Theorem 1.46: Algebraic Künneth theorem

Let R be a commutative ring of global dimension ≤ 1 . Let C, D be complexes of projective R -modules. Then the following map is R -linearly split injective

$$\bigoplus \Phi: \bigoplus_{p+q=n} H_p(C) \otimes_R H_q(D) \rightarrow H_n(C \otimes_R D)$$

Moreover the cokernel is naturally isomorphic to

$$\bigoplus_{p+q=n-1} \mathrm{Tor}^R(H_p(C), H_q(D)).$$

Equivalently, there is a natural and split short exact sequence

$$0 \rightarrow \bigoplus_{p+q=n} H_p(C) \otimes_R H_q(D) \xrightarrow{\Phi} H_n(C \otimes_R D) \rightarrow \bigoplus_{p+q=n-1} \mathrm{Tor}^R(H_p(C), H_q(D)) \rightarrow 0$$

Proof. We let $Z = \{Z_q\}_{q \in \mathbb{Z}}$ be the complex of R modules with $d = 0$ where $Z_q = \mathrm{Ker}(d: D_q \rightarrow D_{q-1})$, let $B = \{B_q\}$ be the complex with $d = 0$ where $B_q = \mathrm{Im}(D: D_{q+1} \rightarrow D_q)$. We have a short exact sequence of complexes of R -modules

$$0 \rightarrow Z \xrightarrow{\mathrm{incl}} D \xrightarrow{d} B[1] \rightarrow 0$$

where $B[1]$ is the complex B shifted up by 1.

We have $B_q \subseteq Z_q \subseteq D_q$ projective by hypothesis. Since R has global dimension ≤ 1 , B_q and Z_q are also projective.

$$0 \rightarrow Z_q \rightarrow D_q \xrightarrow{d} B_{q-1} \rightarrow 0$$

⁸no zero divisors and every ideal is generated by a single element.

is short exact, B_{q-1} is projective, so the sequence splits.

For every R -module N , the sequence

$$0 \rightarrow N \otimes_R Z_p \rightarrow N \otimes_R D_q \rightarrow N \otimes_R B_{q-1} \rightarrow 0$$

is exact.

This means we get a short exact sequence of complexes

$$0 \rightarrow C \otimes_R Z \rightarrow C \otimes_R D \rightarrow C \otimes_R B[1] \rightarrow 0$$

This means we get a long exact homology sequence

$$\rightarrow H_n(C \otimes_R Z) \xrightarrow{H_n(C \otimes_R \text{incl})} H_n(C \otimes D) \xrightarrow{H_n(C \otimes d)} H_{n-1}(C \otimes_R B) \xrightarrow{\partial} H_{n-1}(C \otimes_R Z) \rightarrow \dots$$

Since Z has trivial differential:

$$H_n(C \otimes_R Z) = H_n\left(\bigoplus_{q \in \mathbb{Z}} C[q] \otimes Z_q\right) \cong \bigoplus_{q \in \mathbb{Z}} H_n(C[q] \otimes Z_q) \cong \bigoplus_{q \in \mathbb{Z}} H_n(C[q]) \otimes_R Z_q = \bigoplus_{p \in \mathbb{Z}} H_{n-q}(C) \otimes_R Z_q$$

where we use that Z_q is projective.

Similarly $H_n(C \otimes_R B) \cong \bigoplus_{q \in \mathbb{Z}} H_{n-q}(C) \otimes B_q$.

This gives us a long exact sequence

$$\dots \rightarrow \bigoplus_{p+q=n} H_p(C) \otimes_R Z_q \rightarrow H_n(C \otimes_R D) \rightarrow \bigoplus_{p+q=n-1} H_p(C) \otimes B_q \rightarrow \bigoplus_{p+q=n-1} H_p(C) \otimes Z_q$$

This splits up into short exact sequences

$$0 \rightarrow \bigoplus_{p+q=n} \text{Coker}(H_p(C) \otimes B_q \rightarrow H_p(C) \otimes_R Z_p) \rightarrow H_n(C \otimes_R D) \rightarrow \bigoplus_{p+q=n-1} \text{Ker}(H_p(C) \otimes_R B_q \rightarrow H_p(C) \otimes Z_q) \rightarrow 0$$

We know $0 \rightarrow B_q \rightarrow Z_q \rightarrow H_q(D)$ is a projective resolution of $H_q(D)$.

This means for all R -modules N ,

$$\text{Tor}^R(N; H_q(D)) = \text{Ker}(N \otimes_R B_q \rightarrow N \otimes_R Z_q)$$

$$N \otimes_R H_q(D) \cong \text{Coker}(N \otimes_R B_q \rightarrow N \otimes_R Z_q)$$

So we get:

$$0 \rightarrow \bigoplus_{p+q=n} H_p(C) \otimes H_q(D) \xrightarrow{\Phi} H_n(C \otimes_R D) \rightarrow \bigoplus_{p+q=n-1} \text{Tor}^R(H_p(C), H_q(D)) \rightarrow 0$$

for next lecture remains, that Φ has a R -linear retraction!

[23.04.2025, Lecture 5]
[28.04.2025, Lecture 6]

For the R -linear splitting.

Because B_q is projective, the following s.e.s. splits:

$$0 \rightarrow Z_q \xrightarrow{\text{incl}} D_q \xrightarrow{d} B_q \rightarrow 0$$

and the map Z_q to D_q admits a retraction. We choose a retraction $r_q: D_q \rightarrow Z_q$ to the inclusion.

Then

$$\begin{array}{ccccc}
 D_q + 1 & & & & \\
 \downarrow d & \searrow 0 & & & \\
 B_q & & & & \\
 \downarrow \cap & \searrow 0 & & & \\
 D_q & \xrightarrow{r_q} & Z_q & \longrightarrow & H_q(D)
 \end{array}$$

the retraction $\{r_q\}_{q \in \mathbb{Z}}$ for a morphism of chain complexes

$$r: D \rightarrow \{H_q(D), d = 0\}_q$$

that induces the identity on homology.

$H_q(r) \cong H_q(D) \rightarrow H_q(H_*(D), d = 0) = H_q(D)$. Similalry, there is a chain map $\rho: C \rightarrow \{H_p(C), d = 0\}$ that is the identity on homology. This gives a chain mpa $\rho \otimes_R r: C \otimes_R D \rightarrow (H_*(C) \otimes_R H_*(D), d = 0)$ which on homology

$$H_n(\rho \otimes r): H_n(C \otimes_R D) \rightarrow H_n(H_*(C) \otimes_R H_*(D), d = 0) = \bigoplus_{p+q=n} H_n(C) \otimes_R H_n(D)$$

which is a retraction to

$$\Psi: \bigoplus_{p+q=n} H_p(C) \otimes_R H_q(D) \rightarrow H_n(C \otimes_R D)$$

□

Example 1.47. Let R be a field. Then every module is free, hence projective, and

$$\text{Tor}^R(M, N) = 0$$

for all R -modules M, N . For all complexes of R -modules C, D ;

$$\psi: \bigoplus_{p+q=n} H_p(C) \otimes_R H_q(D) \xrightarrow{\cong} H_n(C \otimes_R D).$$

is an isomorphism.

If $R = \mathbb{Z}$. Let C, D be a complex of free abelian groups. Then there is a split s.e.s.

$$0 \rightarrow \bigoplus_{p+q=n} H_p(C) \otimes H_q(D) \rightarrow H_n(C \otimes D) \rightarrow \bigoplus_{p+q=n-1} \text{Tor}(H_p(C), H_q(D)) \rightarrow 0$$

Construction 1.48 (Homology exterior pairing). Let X, Y be simplicial sets. Let R be a commutative ring. We define

$$\times: H_p(X, R) \otimes_R H_q(Y, R) \rightarrow H_{p+q}(X \times Y, R)$$

as the composite

$$H_p(C_*(X, R)) \otimes_R H_q(C_*(Y, R)) \xrightarrow{\Phi} H_{p+q}(C_*(X, R) \otimes C_*(Y, R)) \xrightarrow{H_{p+q}(\text{EZ})} H_{p+q}(C_*(X \times Y, R))$$

For topological spaces A, B we Define

$$\times: H_p(A; R) \otimes_R H_q(B, R) \rightarrow H_{p+q}(A \times B, R)$$

as the composite

$$H_p(\mathcal{S}(A), R) \otimes_R H_q(\mathcal{S}(B), R) \xrightarrow{\times} H_{p+q}(\mathcal{S}(A) \otimes \mathcal{S}(B), R) \cong H_{p+q}(\mathcal{S}(A \times B); R)$$

where the isomorphism is given by the fact, that simplicial complex commutes with products. The isomorphism is the canonical map

$$\mathcal{S}(A) \times \mathcal{S}(B) \xleftarrow{(\mathcal{S}(p_A), \mathcal{S}(p_B))} \mathcal{S}(A \times B)$$

Theorem 1.49: Künneth theorem for homology with field coefficients

Let R be a field. Let X, Y be simplicial sets or spaces. Then the homology external product

$$\times: \bigoplus_{p+q=n} H_p(X, R) \otimes_R H_q(Y, R) \rightarrow H_n(X \times Y; R)$$

is an isomorphism.

Proof. Follows directly from algebraic Künneth + Eilenberg-Zilber □

Theorem 1.50: Künneth theorem for homology

Let X, Y be spaces or simplicial sets. Then there is a natural and split s.e.s.

$$0 \rightarrow \bigoplus_{p+q=n} H_p(X, \mathbb{Z}) \otimes H_q(Y, \mathbb{Z}) \rightarrow H_n(X \times Y; \mathbb{Z}) \rightarrow \bigoplus_{p+q=n-1} \text{Tor}(H_p(X, \mathbb{Z}), H_q(Y, \mathbb{Z})) \rightarrow 0$$

Special Case. Let X, Y be spaces or simplicial sets. Suppose that $H_n(X, \mathbb{Z})$ is free for all $n \geq 0$. Then

$$\bigoplus_{p+q=n} H_p(X, \mathbb{Z}) \otimes H_q(Y, \mathbb{Z}) \xrightarrow{\Phi} H_n(X \times Y; \mathbb{Z})$$

is an isomorphism.

Next we want to show the Künneth theorem for cohomology. The strategy:

- EZ provides a chain homotopy equivalence $C_*(X, \mathbb{Z}) \otimes C_*(Y, \mathbb{Z}) \rightarrow C_*(X \times Y, \mathbb{Z})$.
- $\text{Hom}(_, R): \mathbf{Chains} \rightarrow \mathbf{coChains}_R$ preserves chain homotopies, so

$$\text{Hom}(C_*(X, \mathbb{Z}), R) \otimes \text{Hom}(C_*(Y, \mathbb{Z}), R) \cong \text{Hom}((C_*(X \times Y), \mathbb{Z}), R)$$

- in favorable cases we can relate

$$H^*(\text{Hom}(C, R) \otimes_R \text{Hom}(D, R)) \rightarrow H^*(\text{Hom}(C, R)) \otimes_R H^*(\text{Hom}(D, R))$$

- apply the algebraic Künneth theorem.

Step 3 is the hard step.

1.4.3 Relation between Homs and Tensors

Let A be an abelian group and R a commutative ring. We make the set $\text{Hom}(A, R)$ of group homomorphisms into an R module by pointwise addition and skalar multiplication. So $f, g \in \text{Hom}(A, R)$, $r \in R$. then

$$(f + g)(a) = f(a) + g(a), \quad ((r \cdot f)(a) = r \cdot f(a))$$

Let B be another abelian group. Then

$$\bullet: \text{Hom}(A, R) \times \text{Hom}(B, R) \rightarrow \text{Hom}(A \otimes B, R)$$

by $(f \bullet g)(a \otimes b) = f(a) \cdot g(b)$. This is additive in f and g .

$$(f + f') \bullet g = (f \bullet g) + (f' \bullet g)$$

and

$$(rf) \bullet g = r \cdot (f \bullet g) = f \bullet (r \cdot g)$$

for all $r \in R$. This means this extends to a well-defined R -linear map

$$\text{Hom}(A, R) \otimes_R \text{Hom}(B, R) \rightarrow \text{Hom}(A \otimes B, R)$$

Proposition 1.51. Let A, B be abelian groups and R a commutative ring. If A is finitely generated and free, then

$$\text{Hom}(A, R) \otimes_R \text{Hom}(B, R) \rightarrow \text{Hom}(A \otimes B, R)$$

is an isomorphism of R -modules.

Proof. For $A = \mathbb{Z}$:

$$\begin{array}{ccc} \text{Hom}(\mathbb{Z}, R) \otimes_R \text{Hom}(B, R) & \xrightarrow{\bullet} & \text{Hom}(\mathbb{Z} \otimes B, R) \\ \downarrow \text{ev} \otimes_R \text{Hom}(B, R) & & \downarrow \cong \text{Hom}(k, R) \\ R \otimes_R \text{Hom}(B, R) & \xrightarrow[r \otimes g \mapsto r \cdot g]{\cong} & \text{Hom}(B, R) \end{array}$$

where we have $k: B \rightarrow \mathbb{Z} \otimes B$ with $b \mapsto 1 \otimes b$.

Suppose the claim holds for A and A' . Then it holds for $A \oplus A'$.

$$\begin{array}{ccc} \text{Hom}(A \oplus A', R) \otimes_R \text{Hom}(B, R) & \xrightarrow{\bullet} & \text{Hom}((A \oplus A') \otimes B, R) \\ \parallel & & \parallel \\ (\text{Hom}(A, R) \oplus \text{Hom}(A', R)) \otimes_R \text{Hom}(B, R) & & \text{Hom}((A \otimes B) \oplus (A' \otimes B), R) \\ \parallel & & \parallel \\ (\text{Hom}(A, R) \otimes_R \text{Hom}(B, R)) \oplus (\text{Hom}(A', R) \otimes_R \text{Hom}(B, R)) & \xrightarrow[\text{by assumption}]{\cong} & \text{Hom}(A \otimes B, R) \oplus \text{Hom}(A' \otimes B, R) \end{array}$$

The claim holds for $A = \mathbb{Z}^k$, $k \in \mathbb{N}$. any finitely generated free abelian group is isomorphic to \mathbb{Z}^k . \square

Example 1.52. $R = \mathbb{F}_2$ $A = B = \mathbb{Z}[\mathbb{N}]$. Then $\text{Hom}(\mathbb{Z}[\mathbb{N}], R) \cong \text{maps}(\mathbb{N}, R)$ by evaluation of generators. This is R -linear by the R -module structure on $\text{maps}(\mathbb{N}, R)$.

$$\text{Hom}(A, R) \otimes_R \text{Hom}(B, R) \xrightarrow{\bullet} \text{Hom}(A \otimes B, R)$$

$$\text{maps}(\mathbb{N}, R) \otimes_R \text{maps}(\mathbb{N}, R) \quad \text{Hom}(\mathbb{Z}[\mathbb{N} \times \mathbb{N}], R)$$

$$\text{maps}(\mathbb{N} \times \mathbb{N}, R)$$

This is however not an isomorphism.

$A = B = \mathbb{Z}/2$ and $R = \mathbb{Z}/4$. Then $\text{Hom}(A, R) = \text{Hom}(B, R) = \text{Hom}(\mathbb{Z}/2, \mathbb{Z}/4)$ is cyclic of order two generated by $i: \mathbb{Z}/2 \rightarrow \mathbb{Z}/4, n + 2\mathbb{Z} \mapsto 2n + 4\mathbb{Z}$.

$$\begin{array}{ccc}
 \text{Hom}(A, R) \otimes_R \text{Hom}(B, R) & \xrightarrow{\bullet} & \text{Hom}(A \otimes B, R) \\
 \parallel & & \\
 \text{Hom}(\mathbb{Z}/2, \mathbb{Z}/4) \otimes_{\mathbb{Z}/4} \text{Hom}(\mathbb{Z}/2, \mathbb{Z}/4) & & \text{Hom}(\mathbb{Z}/2 \otimes \mathbb{Z}/2, \mathbb{Z}/4) \\
 \parallel & & \parallel \\
 \mathbb{Z}/2 & \xrightarrow{0} & \mathbb{Z}/2
 \end{array}$$

This shows, that both assumptions are strictly necessary.

Now let C, D be complexes of abelian groups. Then $\text{Hom}(C, R), \text{Hom}(D, R)$ are cochain complexes of R -modules.

$$\text{Hom}(C, R)^n = \text{Hom}(C_n, R)$$

and

$$d^n: \text{Hom}(C, R)^n \rightarrow \text{Hom}(C, R^{n+1}) = \text{Hom}(D_{n+1}, R)$$

The sum of the \oplus homomorphism gives a cochain map

$$\bigoplus: \text{Hom}(C, R) \otimes_R \text{Hom}(D, R) \rightarrow \text{Hom}(C \otimes D, R)$$

which is in dimension n :

$$\bigoplus_{p+q=n} \text{Hom}(C_p, R) \otimes_R \text{Hom}(D_q, R) \xrightarrow{\text{sum of } \bigoplus} \text{Hom}\left(\bigoplus_{p+q=n} C_p \otimes D_q, R\right)$$

Proposition 1.53. Let C and D be chain complexes of abelian groups, such that $C_n = 0 = D_n$ for $n < 0$ and that C_n is finitely generated and free for all $n \geq 0$. Then \bigoplus is an isomorphism.

$$\bigoplus: \text{Hom}(C, R) \otimes_R \text{Hom}(D, R) \rightarrow \text{Hom}(C \otimes D, R)$$

is an isomorphism of cochain complexes.

Proof. The vanishing hypothesis makes the potentially infinite sums

$$\bigoplus_{p+q=n} \text{Hom}(C_p, R) \otimes_R \text{Hom}(D_q, R)$$

finite.

Then $\text{Hom}(_, R)$ preserves sums. And

$$\text{Hom}(C_p, R) \otimes_R \text{Hom}(D_q, R) \xrightarrow{\bigoplus} \text{Hom}(C_p \otimes D_q, R)$$

is an isomorphism by the previous proposition. \square

This is not yet good enough to apply to topological spaces, as they are very not finitely generated.

Proposition 1.54. Let C be a chain complex of free abelian groups, such that $C_n = 0$ for $n < 0$. Suppose that $H_n(C)$ is finitely generated for all $n > 0$.

Then there is a subcomplex B of C , such that

- B_n is finitely generated and free for all $n \geq 0$.
- The inclusion $B \rightarrow C$ is a chain homotopy equivalence.

Proof. We construct subgroups B_n of C_n by induction on $n \geq 0$, such that

- $d(B_n) \subseteq B_{n-1}$
- the inclusions of $0 \rightarrow B_n \xrightarrow{d} B_{n-1} \xrightarrow{d} \cdots \rightarrow B_0 \rightarrow 0$
- into C induces an isomorphism on H_i for all $0 \leq i \leq n-1$ and an epimorphism on H_n .

Induction start: Let x_1, \dots, x_m be elements of C_0 , that generate $H_0(C)$. Select B_0 to be the subgroups of C_0 generated by x_1, \dots, x_m .

Induction step: Suppose B_0, \dots, B_{n-1} have been constructed fulfilling the conditions. Let x_1, \dots, x_m be cycles in C_n whose homology classes generate $H_n(C)$, which is possible because $H_n(C)$ is finitely generated. Set

$$Z = \text{Ker}(d: B_{n-1} \rightarrow B_{n-2}) \cap \text{Im}(d: C_n \rightarrow C_{n-1})$$

which is finitely generated because B_{n-1} is. Let z_1, \dots, z_k generate this intersection. Choose $y_1, \dots, y_k \in C_n$, such that $d(y_i) = z_i$ for $1 \leq i \leq k$.

Let B_n be the subgroup generated by $x_1, \dots, x_m, y_1, \dots, y_k$. Then $d(B_n) \subseteq B_{n-1}$.

Let $B_{\leq n}$ and $B_{< n}$ be the subcomplexes of C generated by B_0, \dots, B_n and B_0, \dots, B_{n-1}

Then $B_{< n} \subseteq B_{\leq n} \subseteq C$ where $B_{< n}$ induces isomorphism on H_i for $0 \leq i \leq n-2$ and epi on H_{n-1} . Similarly $B_{< n} \rightarrow B_{\leq n}$ is iso in dimension $\leq n-1$.

Then $B_{\leq n}$ is an Isomorphism on H_i for $0 \leq i \leq n-2$ and surjective on H_n because we include x_1, \dots, x_m that generate $H_n(C)$.

Let $x \in B_{n-1}$ be any cycle whose class is in the kernel of $H_{n-1}(B_{< n}) \rightarrow H_{n-1}(C)$. Then $x \in Z$ so x is a linear combination of the classes z_1, \dots, z_k and hence a boundary of a linear combination of y_1, \dots, y_k . So $x = d(w)$ for some $w \in B_n$. Then

$$\begin{array}{ccc} & H_{n-1}(B_n) & \\ \nearrow & & \searrow \\ H_{n-1}(B_{< n}) & \xrightarrow{\quad\quad\quad} & H_{n-1}(C) \end{array}$$

the class of x maps to 0 and the map becomes injective and hence an isomorphism.

We let B be the subcomplex of C generated by all B_i for all $i \geq 0$. Then the inclusion $B \rightarrow C$ induces an isomorphism on H_i for all $i \geq 0$, so it is a quasi-isomorphism.

By the end of last term we proved, it is already a chain homotopy equivalence! \square

[23.04.2025, Lecture 6]
[30.04.2025, Lecture 7]

Korollar 1.55 (Algebraic Künneth theorem, cohomology). *Let R be a commutative ring of global dimension ≤ 1 . Let C, D be chain complexes of abelian groups such that $C_n = 0 = D_n$ for $n < 0$ and all C_n are free and $H_n(C)$ is finitely generated free.*

Then for all $n \geq 0$:

$$\bigoplus_{p+q=n} H^p(\text{Hom}(C, R)) \otimes_R H^q(\text{Hom}(D, R)) \xrightarrow{\Phi} H^n(\text{Hom}(C \otimes D, R))$$

is injective and its cokernel is isomorphic to

$$\bigoplus_{p+q=n+1} \text{Tor}^R(H^p(\text{hom}(C, R)), H^q(\text{Hom}(D, R)))$$

Warning. We do not assume, that there is a splitting.

Proof. „Basically just putting all the hard stuff we*ve already done together in the right way.“

Case 1 Suppose that also C_n is finitely generated for all $n \geq 0$. Then $\bullet: \text{Hom}(C, R) \otimes_R \text{Hom}(D, R) \rightarrow \text{Hom}(C \otimes D, R)$ is an isomorphism of cochain complexes. Applying the homological algebraic Künneth theorem to

$$H^n(\text{Hom}(C \otimes D, R)) \cong H^n(\text{Hom}(C, R) \otimes_R \text{Hom}(D, R))$$

since C_n is finitely generated and free, it is isomorphic to \mathbb{Z}^k for some $k \geq 0$, so $\text{Hom}(C, R)^n = \text{Hom}(C_n, R) \cong \text{Hom}(\mathbb{Z}^k, R) = R^k$ which is free hence projective as an R -module for all $n \geq 0$.

Caveat 1. we make cochain complexes into chain complexes, then apply Künneth, then come back. This turns $n - 1$ in the \oplus for $\text{Tor}R$ into $n + 1$.

Caveat 2. The proof of the homological Künneth theorem (without the splitting) used only that one complex is dimensionwise projective. Hence it is no problem, that D is not projective.

General case We choose a subcomplex B of C such that B_n is finitely generated for all $n \geq 0$ and $B \hookrightarrow C$ is a chain homotopy equivalence. Then

$$\text{Hom}(i, R): \text{Hom}(B, R) \rightarrow \text{Hom}(C, R)$$

is a chain homotopy equivalence of R -module complexes.⁹

Note Additive functors preserve chain homotopy equivalences, however not quasi-isomorphisms. Because of that, quasi-Isomorphisms and chain homotopy equivalences are quite different.

Similarly we see

$$\text{Hom}(i \otimes D, R): \text{Hom}(C \otimes D, R) \rightarrow \text{Hom}(B \otimes D, R)$$

is a chain homotopy equivalence.

This gives a commutative square in **coChains** $_R$:

$$\begin{array}{ccc} \bigoplus_{p+q=n} H^p(\text{Hom}(C, R)) \otimes H^q(\text{Hom}(D, R)) & \xleftarrow{\Phi} & H^n(\text{Hom}(C \otimes D, R)) \\ \downarrow \cong & & \\ \bigoplus_{p+q=n} H^p(\text{Hom}(B, R)) \otimes_R H^q(\text{Hom}(D, R)) & \xleftarrow[\text{by special case}]{\Phi} & H^n(\text{Hom}(B \otimes D, R)) \end{array}$$

□

Construction 1.56. Let X, Y be spaces or simplicial sets. R a commutative ring. The *exterior cup product*

$$\times: H^p(X, R) \times H^q(Y, R) \rightarrow H^{p+q}(X \otimes Y, R)$$

is defined by $(x, y) \mapsto p_1^*(x) \cup p_2^*(y)$, where $p_1: X \times Y \rightarrow X$ and $p_2: X \times Y \rightarrow Y$.

Recall. The AW-map is

$$\text{AW}: C_*(X \times Y) \rightarrow C_*(X) \otimes C_*(Y)$$

⁹This is due to the Hom-functor being additive. Unfortunately I don't know what that means.

Proposition 1.57. Let X, Y be simplicial sets, R commutative ring. Then the composite

$$H^p(X, R) \otimes_R H^q(Y, R) \xrightarrow{[f]} [g] \mapsto [f \otimes g] \Phi H^{p+q}(\text{Hom}(C_*(X), R) \otimes_R \text{Hom}(C_*(Y), R)) \xrightarrow{H^{p+q}(\bullet)} H^{p+q}(\text{Hom}(C_*(X) \otimes C_*(Y), R))$$

equals the external cup product.

Proof. In the notes. □

Theorem 1.58: Künneth theorem in cohomology

Let R be a commutative ring of global dimension ≤ 1 . Let X, Y be spaces such that $H_n(X, \mathbb{Z})$ is finitely generated for all $n \geq 0$. Then the total exterior cup product map

$$\bigoplus_{p+q=n} H^p(X, R) \otimes_R H^q(Y, R) \rightarrow H^n(X \times Y, R)$$

is injective, and its cokernel is naturally isomorphic to

$$\bigoplus_{p+q=n+1} \text{Tor}^R(H^p(X, R), H^q(Y, R))$$

Proof. Similar to the homological one. Use the cohomological algebraic Künneth theorem and the Eilenberg-Zilber theorem. You can read it up somewhere. □

Remark 1.59. Let X be a CW-complex of finite type i.e. such that it has only finitely many cells in every dimension. (ex. $\mathbb{R}P^\infty$). Then

$$C_A^{Cell}(X, \mathbb{Z})$$

is finitely generated free in every dimension, hence $H_n^{cell}(X, \mathbb{Z}) \cong H_n(X, \mathbb{Z})$ if finitely generated, so Künneth theorem applies.

Construction 1.60. Let A, B be graded-commutative¹⁰ rings. Then $A \otimes B$ is another graded-commutative ring by

$$(A \otimes B)_n = \bigoplus_{p+q=n} A_p \otimes B_q$$

with multiplication for $a \in A_p, b \in B_q, a' \in A_{p'}, b' \in B_{q'}$.

$$(a \otimes b) \cdot (a' \otimes b') = (-1)^{p' \cdot q} (aa') \otimes (bb')$$

Check for well-definedness yourself.

Korollar 1.61. Let R be a field, X, Y spaces and suppose, that $H_n(X, \mathbb{Z})$ is finitely generated for all $n \geq 0$. Then

$$\times: H^*(X, R) \otimes_R H^*(Y, R) \rightarrow H^*(X \times Y, R)$$

is an isomorphism of graded-commutative R -algebras.

Note. We already knew that this is a isomorphism of abelian groups. The new information is, that this is compatible with ring structure.

¹⁰ $a \cdot b = (-1)^{\deg(A) \cdot \deg(B)} b \cdot a$

Proof. We take $x \in H^p(X, R)$, $x' \in H^{p'}(X, R)$, $y \in H^q(Y, R)$, $y' \in H^{q'}(Y, R)$ and then

$$\begin{aligned} (x \cup x') \times (y \cup y') &= p_1^*(x \cup x') \cup p_2^*(y \cup y') \\ &= (p_1^*(x) \cup p_1^*(x')) \cup (p_2^*(y) \cup p_2^*(y')) \\ &= (-1)^{p' \cdot q} (p_1^*(x) \cup p_2^*(y)) \cup (p_1^*(x') \cup p_2^*(y')) \\ &= (-1)^{p \cdot q'} (x \times y) \cup (x' \times y') \end{aligned}$$

□

Korollar 1.62. *Let X, Y be spaces such that $H_n(X, \mathbb{Z})$ is finitely generated and free for all $n \geq 0$. Then*

$$H^*(X, \mathbb{Z}) \otimes H^*(Y, \mathbb{Z}) \rightarrow H^*(X \times Y, \mathbb{Z})$$

is an isomorphism of graded-commutative rings.

Now we are actually calculating some cohomology rings. Namely for $S^k \times S^l$, $S^1 \times \cdots \times S^1$ and $\mathbb{C}P^2$.

Remember

$$H^n(S^k) = \begin{cases} \mathbb{Z} & n = 0, k \\ 0 & n \neq 0, k \end{cases}$$

and assume $k \geq 1$. For dimensional reasons, the cup product on $H^*(S^k, \mathbb{Z})$ is trivial. $H^*(S^k, \mathbb{Z})$ is dimensionwise finitely generated free, and hence for every space Y the exterior cup product

$$H^*(S^k, \mathbb{Z}) \otimes H^*(Y, \mathbb{Z}) \rightarrow H^*(S^k \times Y, \mathbb{Z})$$

is an isomorphism of graded-commutative rings. Take $Y = S^l$ for $l \geq 1$.

Let $e_k \in H^n(S^k; \mathbb{Z})$ be one of the two generators. Then $H^+(S^k, \mathbb{Z}) = \Lambda(e_k)$ where Λ denotes an exterior product. This includes $e_k^2 = 0$. We define $a := p_1^*(e_k) \in H^k(S^k \times S^l; \mathbb{Z})$ and $b := p_2^*(e_l) \in H^l(S^k \times S^l; \mathbb{Z})$. Then

$$H^*(S^k, \mathbb{Z}) \otimes H^*(S^l, \mathbb{Z}) \xrightarrow{x} H^*(S^k \times S^l; \mathbb{Z}) = \mathbb{Z}\{1 \times 1, 1 \times e_l, e_k \times 1, e_k \cdot e_l\}$$

where we have $1 \times 1 = 1$, $1 \times e_l = b$, $e_k \times 1 = a$, $e_k \cdot e_l = a \cup b$.

We look at multiplicative relations:

$$a^2 = 0, b^2 = 0$$

and so

$$a^2 = (p_1^*(e_k))^2 = p_1^*(e_k^2) = p_1^*(0) = 0$$

If k or l is even, then $a \cup b = b \cup a$ and if both are odd, then $a \cup b = -b \cup a$.

We summarize, if k and l are even, then

$$H^*(S^k \times S^l; \mathbb{Z}) = \mathbb{Z}[a, b]/(a^2 = 0, b^2 = 0)$$

and if one is odd

$$H^*(S^k \times S^l; \mathbb{Z}) = \Lambda(a, b)$$

where Λ again denotes exterior products.

We give an inductive description of $H^*(S^1 \times \cdots \times S^1; \mathbb{Z})$ n -times. We use, that

$$\times : H^*(S^1; \mathbb{Z}) \otimes H^*\left(\underbrace{S^1 \times \cdots \times S^1}_{n-1 \text{ times}}\right) \cong H^*(S^1 \times \cdots \times S^1, \mathbb{Z})$$

we define $a_i = p_i^*(e_1) \in H^1(\underbrace{S^1 \times \cdots \times S^1}_n; \mathbb{Z})$, where $p_i: (S^1)^n \rightarrow S^1$ is projection to the i -th factor for $1 \leq i \leq n$. We get $a_i^2 = 0$ and $a_i \cup a_j = -a_j \cup a_i$ for $i \neq j$. This gives us, that an additive basis of $H^*(S^1)^n; \mathbb{Z}$ is given by

$$a_{i_1} \cup \cdots \cup a_{i_k} \text{ for all tuples } 1 \leq a_i < a_2 < \cdots < a_k \leq n$$

This gives us $\text{rank}(H^*((Sq)^n \mathbb{Z})) = 2^n$. The multiplicative structure is given by $H^*((S^1)^n, \mathbb{Z}) = \Lambda(a_1, \dots, a_n)$.

Later we will compute $H^*(\mathbb{C}P^n; \mathbb{Z})$ via Poincaré-duality to get $\cong \mathbb{Z}[X]/(X^{n+1})$ for $x \in H^2(\mathbb{C}P^n, 2)$.

We will now use a trick to at least calculate $H^*(\mathbb{C}P^2; \mathbb{Z})$. We know, that

$$H^n(\mathbb{C}P^2, \mathbb{Z}) = \begin{cases} \mathbb{Z} & n = 0, 2, 4 \\ 0 & \text{else} \end{cases}$$

we take $x \in H^2(\mathbb{C}P^2; \mathbb{Z})$ a generator. The multiplicative structure is completely defined by which multiple of the generator of $H^4(\mathbb{C}P^2, \mathbb{Z})$ x^2 is.

We use homogenous coordinate notation for $\mathbb{C}P^2$. For $0 \neq (x, y, z) \in \mathbb{C}^3$ we write $[x, y, z] := \mathbb{C} \cdot (x, y, z) \in \mathbb{C}P^2$. We define a continuous map

$$\mu: \mathbb{C}P^1 \times \mathbb{C}P^1 \rightarrow \mathbb{C}P^2$$

given by $([v, w], [x, y]) \mapsto [vx, vy + wx, wy]$. We let $e = [1, 0]$ a basepoint in $\mathbb{C}P^1$. Then $\mu(e, _), \mu(_, e): \mathbb{C}P^1 \rightarrow \mathbb{C}P^2$. are both the „standard inclusions“ $[x, y] \mapsto [x, y, 0]$.

Proposition 1.63. The map $\mu^*: H^4(\mathbb{C}P^2, \mathbb{Z}) \rightarrow H^4(\mathbb{C}P^1 \times \mathbb{C}P^1, \mathbb{Z})$ is injective and its image has index 2.

proof next time.

[30.04.2025, Lecture 7]

[5.05.2025, Lecture 8] Rather sleepy today, quality may be accordingly.

Note. Remember $\mathbb{C}P^2 \cong S^2$.

Proof. We will drop coefficients from the notation. $H^*(X) := H^*(X; \mathbb{Z})$. The continuous map $\mathbb{C}^2 \rightarrow \mathbb{C}P^2$, $\pi(a, b) = (a^2 - b, 2a, 1)$ is an open embedding and a homoeomorphism onto the open 4-cell $\mathbb{C}P^2 \setminus \mathbb{C}P^1$. That is just the set $[x, y, 1]$ for $(x, y) \in \mathbb{C}^2$. Then

$$(x, y) = (a^2 - b, 2a) \implies a = y/2, b = (a - x = y^2/4 - x)$$

This gives an isomorphism of relative cohomology groups

$$\pi^*: H^4(\mathbb{C}P^2 \setminus \mathbb{C}P^1, \mathbb{C}P^2 \setminus (\mathbb{C}P^1 \cup [0, 0, 1])) \rightarrow H^4(\mathbb{C}^2, \mathbb{C}^2 \setminus (0, 0))$$

Then we have AN EXCISION isomorphism:

$$H^4(\mathbb{C}P^2, \mathbb{C}P^2 \setminus [0, 0, 1]) \cong H^4(\mathbb{C}P^2 \setminus \mathbb{C}P^1, \mathbb{C}P^2 \setminus \mathbb{C}P^1 \cup [0, 0, 1])$$

The long exact sequence of the pair gives an isomorphism

$$H^4(\mathbb{C}P^2, \mathbb{C}P^2 \setminus [0, 0, 1]) \rightarrow H^4(\mathbb{C}P^2)$$

We also Define

$$\pi': \mathbb{C}^2 \rightarrow \mathbb{C}P^1 \times \mathbb{C}P^1, (a, b) \mapsto ([a + b, 1], [a - b, 1])$$

A similar calculation gives

$$H^4(\mathbb{C}P^1 \times \mathbb{C}P^1, \mathbb{C}P^1 \times \mathbb{C}P^1 \setminus ([0, 1], [0, 1]))$$

as isomorphic to $H^4(\mathbb{C}P^1 \times \mathbb{C}P^1)$.

We now also define ν .

$$\nu: \mathbb{C}^2 \rightarrow \mathbb{C}^2; \quad (a, b) \mapsto (a, b^2)$$

Now a diagram I didn't copy commutes.

The problem now reduces to show that

$$\nu^*: H^4(\mathbb{C}^2, \mathbb{C}^2 \setminus (0, 0)) \rightarrow H^4(\mathbb{C}^2, \mathbb{C}^2 \times 0), 0$$

is multiplication by 2.

A diagram I didn't copy. He applied Künneth and found out some map is multiplication by 2. \square

Proposition 1.64. Let $x \in H^2(\mathbb{C}P^2, \mathbb{Z})$ be an additive generator. Then x^2 is an additive generator of $H^4(\mathbb{C}P^2, \mathbb{Z})$. So $H^*(\mathbb{C}P^2, \mathbb{Z})$ is a truncated polynomial algebra i.e.

$$H^*(\mathbb{C}P^2, \mathbb{Z}) = \mathbb{Z}[X]/(x^3)$$

Outlook. $H^*(\mathbb{C}P^m; \mathbb{Z}) = \mathbb{Z}[X]/(x^{m+1})$ This will be proven later using Poincaré-Duality.

Proof. We write $i: \mathbb{C}P^1 \rightarrow \mathbb{C}P^2$ for „the inclusion“, $i[x, y] = [x, y, 0]$. Then

$$H^*(\mathbb{C}P^1, \mathbb{Z}) = \mathbb{Z}\{1, i^2(x)\}$$

$$\times: H^*(\mathbb{C}P^2) \otimes H^*(\mathbb{C}P^1) \cong H^*(\mathbb{C}P^1 \times \mathbb{C}P^1)$$

we write $a := p_1^*(i^*(x)), b := p_2^*(i^*(x))$. Then

$$H^*(\mathbb{C}P^1 \times \mathbb{C}P^1) = \mathbb{Z}\{1, a, b, a \cdot b\}$$

with $a^2 = b^2 = 0, ab = ba$.

Claim. We have

$$\begin{aligned} \mu^*(x) = a + b & \in H^2(\mathbb{C}P^1 \times \mathbb{C}P^1) \\ & \downarrow \cong \\ & H^2(\mathbb{C}P^1 \vee \mathbb{C}P^1) \\ & \downarrow \cong \\ & H^2(\mathbb{C}P^1) \times H^2(\mathbb{C}P^1) \end{aligned}$$

where we use that the wedge is an isomorphism on the 2-skeleton of $\mathbb{C}P^1 \times \mathbb{C}P^1$. The composite map is given by

$$z \mapsto ((e, _)^*(z)), (_ , e)^*(z))$$

We note

$$\begin{aligned} (e, _)^*(a + b) &= (e, _)^*(p_1^*(i^*(x))) + (e, _)^*(p_2^*(i^*(x))) \\ &= \underbrace{(i \circ p_1 \circ (e, _))^*(x)}_{\text{constant}} + \underbrace{(i \circ p_2 \circ (e, _))^*(x)}_{\text{identity}} \\ &= i^*(x) \end{aligned}$$

and also

$$(e, _)^*(\mu^*(x)) = \underbrace{(\mu \circ (e, _))}_{=1}^* = i^*(x)$$

This gives $\mu^*(x) = a + b$. Now let $y \in H^4(\mathbb{C}P^2)$ be a generator and let $n \in \mathbb{Z}$ be such, that $x^2 = n \cdot y$. Now

$$2ab = (a + b)^2 = (\mu^*(x))^2 = \mu^*(x^2) = \mu^*(ny) = n \cdot \mu^*(y) = n \cdot 2 \cdot ab$$

where the last equality uses degree 2 of μ . This holds in the free abelian group $H^4(\mathbb{C}P^1 \times \mathbb{C}P^1) = \mathbb{Z}\{a, b\}$. This means $2 = 2n$ and hence $n = 1$ and so $x^2 = y$. \square

Application to the Hopf map.

The Hopf map $\eta: S^3 \rightarrow S^2$ is defined as

$$S^3 = S(\mathbb{C}^2) \rightarrow \mathbb{C}P^1 \cong S^2$$

given by $(x, y) \mapsto [x, y]$.

Then $0 \neq [y] \in \pi_3(S^2, *) \cong \mathbb{Z}\{y\}$.

Proposition 1.65. Attaching a 4-cell to $\mathbb{C}P^1$ yields a space homeomorphic to $\mathbb{C}P^2$. Informally: „ η is the attaching map of the 4-cell in $\mathbb{C}P^2$.“

Proof. Consider the map $\alpha: D(\mathbb{C}^2) \rightarrow \mathbb{C}P^2$, $(x, y) \mapsto [x, y, 1 - |x|^2 - |y|^2]$.

This restricts to a homeomorphism from $D(\mathbb{C}^4) \setminus S(\mathbb{C}^2)$ onto $\mathbb{C}P^2 \setminus \mathbb{C}P^1$ and the following commutes:

$$\begin{array}{ccc} S(\mathbb{C}^2) & \xrightarrow{\eta} & \mathbb{C}P^1 \\ \downarrow & & \downarrow i \\ D(\mathbb{C}^2) & \xrightarrow{\alpha} & \mathbb{C}P^2 \end{array} \quad \begin{array}{c} [x, y] \\ \downarrow \\ [x, y, 0] \end{array}$$

this gives a well-defined continuous map $D(\mathbb{C}^2) \cup_{S(\mathbb{C}^2), \eta} \mathbb{C}P^1 \rightarrow \mathbb{C}P^2$. This is a continuous bijection between compact Hausdorff spaces, hence a homeomorphism. \square

Theorem 1.66: Hopf map is not constant

The Hopf map η is not homotopic to a constant map.

Proof. By contradiction. If η was homotopic to the constant map $c: S^3 \rightarrow S^2$, then $D^4 \cup_{S^3, \eta} \mathbb{C}P^1$ would be homotopy-equivalent to $D^4 \cup_{S^3, \text{const}} \mathbb{C}P^1 = \mathbb{C}P^1 \vee (D^4/S^3) \cong S^2 \vee S^4$.

These spaces have the same additive cohomology. However, their cup-product differs. Namely in $H^*(\mathbb{C}P^1 \vee S^4, \mathbb{Z})$ the square of every 2-dimensional class is 0.

As such, $\mathbb{C}P^1 \vee S^4 \not\cong \mathbb{C}P^2$. \square

Outlook. The Hopf map is sometimes presented as the map

$$S(\mathbb{C}^2) \rightarrow \mathbb{C} \cup \{\infty\} = \text{one point compactification of } \mathbb{C} \cong S^2$$

given by $(x, y) \mapsto x/y$. For \mathbb{H} = the quaternions = \mathbb{R}^4 with the skew-field multiplication = $\mathbb{R}\{1, i, j, k\}$ and $i^2 = j^2 = k^2 = ijk = -1$. And then we get

$$\nu: S^7 = S(\mathbb{H}^2) \mapsto \mathbb{H} \cup \{\infty\} = S^4$$

given by $(x, y) \mapsto x/y = xy^{-1} \vee y^{-1}x$. This map is also called the second Hopf-map. Using that most of linear algebra still applies to skew-fields, we can define $\mathbb{H}P^n$ and see by a similar argument, that ν is not nullhomotopic. Then $[\nu] \in \pi_7(S^4, *) \cong \mathbb{Z}\{\nu\} \oplus \mathbb{Z}/?$ Schwede doesn't remember what exactly π_7 is.

Then we also have $\mathbb{O} = \text{Cayley octonians} = \mathbb{R}^8$ with a nonassociative, noncommutative division algebra structure $\cdot: \mathbb{O} \times \mathbb{O} \rightarrow \mathbb{O}$. Then there is still an $\mathbb{O}P^2$ but no general $\mathbb{O}P^n$.

However this is enough to still calculate that $H^*(\mathbb{O}P^2, \mathbb{Z}) = \mathbb{Z}[w]/w^3$ where $w \in H^8(\mathbb{O}P^2, \mathbb{Z})$. And you can show

$$\sigma: S(\mathbb{O}^2) \rightarrow \mathbb{O}P^1 = \mathbb{O} \cup \{\infty\}$$

given by $(x, y) \mapsto x/y$ is non zero-homotopic. And $[\sigma] \in \pi_{15}(S^8) = \mathbb{Z} \oplus \mathbb{Z}/120$.

He also talks about a theorem, that these are all the Hopf-Maps that exist. No more in higher dimensions.

Appendix

List of definitions

1.3	Differential graded ring	5
1.17	Simplicial abelian group	11
1.22	Tensor product of simplicial abelian groups	11
1.25	(p,q) -shuffle	13
1.29	Eilenberg-Zilber map	13
1.34	Representable functor	15
1.39	Tensor Product of R -chains	20
1.40	Projective R -modules	20
1.42	Global dimension of rings	22
1.44	Tor of nice rings	23

List of statements

1.2	fundamental properties of cup product	3
1.6	Properties of the cup-product on homology	5
1.8	Commutativity of the cup-product	6
1.10	\cup_1 -Product	7
1.16	Künneth, simple version	10
1.31	Shuffle maps form a chain map	13
1.32	Eilenberg-Zilber	14
1.37	Acyclic models	16
1.46	Algebraic Künneth theorem	23
1.49	Künneth theorem for homology with field coefficients	26
1.50	Künneth theorem for homology	26
1.58	Künneth theorem in cohomology	31
1.66	Hopf map is not constant	35

Index

(p,q) -shuffle, 13

Alexander Whitney map, 3

cup-product, 3, 5

differential graded ring, 5

Eilenberg-Zilber map, 13

exterior cup product, 30

finite type CW-complex, 31

Global dimension, 22

graded ring, 5

Künneth theorem, 10

Projective Module, 20

representable functor, 15

ring, 2

shuffle map, 13

simplicial abelian group, 11

Tensor product of simplicial abelian groups,
11