

UNIVERSITÄT BONN

Notes for the lecture

Topology II

held by

Stefan Schwede

T_EXed by

Jan Malmström

SoSe 2025

Corrections and improvements

If you have corrections or improvements, contact me via (s94jmalm@uni-bonn.de).

Contents

Lecture	1
I Cup product and Künneth theorem	2
1 Cup-product	4
1.1 Commutativity of the cup-product	7
2 Künneth theorem	12
2.1 The Eilenberg-Zilber-theorem	12
2.2 Yoneda Lemma & Acyclic models	16
2.3 Revisiting Commutativity of the cup-product	21
2.4 Algebraic Künneth theorem.	22
2.5 Homological Künneth theorem	28
2.6 Künneth theorem for cohomology	29
2.7 Example calculations for Cohomology-rings	34
II Manifolds and Poincaré Duality	39
3 Topological manifolds	40
3.1 Orientations	45
3.1.1 Orientability of $\mathbb{R}P^n$	48
3.2 The fundamental class	52
4 Cap Product	55
5 Poincaré Duality	57
5.1 Cohomology with compact support	57
5.1.1 Covariant functoriality for open embeddings	59
5.2 The duality map	63
5.2.1 Mayer-Vietoris sequences for compactly supported cohomology	66
Appendix	A
List of definitions	B
List of statements	C
Index	D

Lecture


Part I

Cup product and Künneth theorem

In last term, we discussed

- CW-complexes
- higher homotopy groups
- Whitehead theorem
- Singular homology
- cellular homology

In the very end, cohomology was started. Remember



Chapter 1

Cup-product

Let X be a simplicial set, and R^1 a ring.

$$C^n(X, R) = \text{maps}(X_n, R)$$

is an abelian group under pointwise addition. There is a differential

$$d^n : C^n(X, R) \rightarrow C^{n+1}(X, R)$$

given by

$$d^n(f)(y) = \sum_{i=0}^{n+1} (-1)^i f(d_i^*(y))$$

with $f : X_n \rightarrow R, y \in X_{n+1}$

Construction 1.1 (Cup product/Alexander Whitney map). The cup product/Alexander Whitney map

$$\cup : C^n(X, R) \times C^m(X, R) \rightarrow C^{m+n}(X, R)$$

with $n, m \geq 0$ is defined by

$$(f \cup g)(x) := f(d_{front}^*(x)) \cdot g(d_{back}^*(x))$$

with $f : X_n \rightarrow R, g : X_m \rightarrow R, x \in X_{n+m}$.

Where we use $[n+m] = \{0, 1, \dots, n+m\}$ and $d_{front} : [n] \rightarrow [n+m], d_{back} : [m] \rightarrow [n+m]$ are given by $d_{front}(i) = i, d_{back}(i) = n+i$. Note, that d_{front} and d_{back} respectively suppress in their notation n and m .

Theorem 1.2: fundamental properties of cup product

The cup-product satisfies the following properties.

1. The AW-map is biadditive and satisfies a boundary formula:

$$d(f \cup g) = (df) \cup g + (-1)^n f \cup (dg) \in C^{m+n+1}(X, R)$$

2. Associativity: For $h \in C^k(X, R), (f \cup g) \cup h = f \cup (g \cup h) \in C^{n+m+k}(X, R)$.

Let $1 \in C^0(X, R)$ be the constant function $1 : X_0 \rightarrow R$ with value 1. Then $1 \cup f = f \cup 1 = f$.

3. Naturality: Let $\alpha : Y \rightarrow X$ be a morphism of simplicial sets. Then

$$\alpha^*(f \cup g) = \alpha^*(f) \cup \alpha^*(g), \quad \alpha^*(1) = 1.$$

where $\alpha^* : C^n(X, R) \rightarrow C^n(Y, R), f \mapsto f \circ \alpha_n$.

¹A ring is not necessarily commutative, but has a unit

Proof.

1. We check some properties: Let $d_{front}: [n] \rightarrow [n+m]$, $d_{back}: [m] \rightarrow [n+m]$ be as in the definition of \cup . Then

$$d_i \circ d_{front} = \begin{cases} d_{front} \circ d_i & 0 \leq i \leq n+1 \\ d_{front} & n+1 \leq i \leq n+m+1 \end{cases}$$

and

$$d_i \circ d_{back} = \begin{cases} d_{back} \circ d_i & 0 \leq i \leq n \\ d_{back} \circ d_{i-n} & n \leq i \leq n+m+1 \end{cases}$$

Note, that for $n+1$ and n respectively the cases are the same.

Now we calculate

$$\begin{aligned} d(f \cup g)(x) &= \sum_{i=0}^{n+m+1} (-1)^i (f \cup g)(d_i^*(x)) \\ &= \sum_{i=0}^{n+m+1} (-1)^i \cdot f(d_{front}^*(x)) \cdot g(d_{back}^*(d_i^*(x))) \\ &= \sum_{i=0}^n (-1)^i \cdot f(d_{front}^*(d_i^*(x))) \cdot g(d_{back}^*(d_i^*(x))) + \sum_{j=1}^{m+1} (-1)^{n+j} \cdot f(d_{front}^*(d_{j+n}^*(x))) \cdot g(d_{back}^*(d_{j+n}^*(x))) \\ &= \sum_{i=0}^{n+1} (-1)^i \cdot f(d_i^*(d_{front}^*(x))) \cdot g(d_{back}^*(x)) + \sum_{j=0}^{m+1} (-1)^{n+j} f(d_{front}^*(x)) \cdot g(d_j^*(d_{back}^*(x))) \\ &= d(f)(d_{front}^*(x)) \cdot g(d_{back}^*(x)) + (-1)^n \cdot f(d_{front}^*(x)) \cdot d(g)(d_{back}^*(x)) \\ &= ((df) \cup g)(x) + (-1)^n \cdot (f \cup dg)(x) \\ &= ((df) \cup g + (-1)^n \cdot f \cup (dg))(x) \end{aligned}$$

2. For $x \in X_{n+m+k}$ we see

$$\begin{aligned} ((f \cup g) \cup h)(x) &= (f \cup g)(d_{front}^*(x)) \cdot h(d_{back}^*(x)) \\ &= f(d_{front}^*(d_{front}^*(x))) \cdot g(d_{back}^*(d_{front}^*(x))) \cdot h(d_{back}^*(x)) \\ &= f(d_{front}^*(x)) \cdot g(d_{middle}^*(x)) \cdot h(d_{back}^*(x)) \end{aligned}$$

Note that we abuse that d_{front} suppresses the indices for which the map is the front map. We have in the last line

$$d_{front}: [n] \rightarrow [n+m+k], d_{middle}: [m] \rightarrow [n+m+k], d_{back}: [k] \rightarrow [n+m+k]$$

defined by

$$d_{front}(i) = i, d_{middle}(i) = n+i, d_{back}(i) = n+m+i$$

this is obviously associative in the inputs²

3. Naturality for $\alpha: Y \rightarrow X$ we see

$$\begin{aligned} (\alpha^*(f \cup g))(y) &= (f \cup g)(\alpha_{n+m}(y)) \\ &= f(d_{front}^*(\alpha_{n+m}(y))) \cdot g(d_{back}^*(\alpha_{n+m}(y))) = f(\alpha_n(d_{front}^*(y))) \cdot g(\alpha_m(d_{back}^*(y))) \\ &= \alpha^*(f)(d_{front}^*(y)) \cdot \alpha^*(g)(d_{back}^*(y)) \\ &= (\alpha^*(f) \cup \alpha^*(g))(y). \end{aligned}$$

□

²for Schwede at least.

Definition 1.3: Differential graded ring

A differential graded ring (dg-ring) is a cochain-complex $A = \{A^n, d^n\}_{n \in \mathbb{Z}}$ equipped with biadditive maps

$$\cdot : A^n \times A^m \rightarrow A^{n+m}, \quad n, m \in \mathbb{Z}$$

and a unit $1 \in A^0$, such that;

- \cdot is associative and has 1 as a unit element.
- the Leibniz rule holds:

$$d(a \cdot b) = (da) \cdot b + (-1)^n \cdot a \cdot (db)$$

with $a \in A^n, b \in A^m$.¹

¹The sign is somehow connected to a sign-rule I couldn't follow. The d moved past the a or something.

Example 1.4. Some Differential graded rings are:

- $C^*(X, R)$ for a simplicial set X and a ring R .
- De Rham complex of a smooth manifold.

Construction 1.5 (Cup-Product on cohomology). Let $A = (A^n, d, \cdot)$ be a dg-ring. We define a map

$$\cdot : H^n(A) \times H^m(A) \rightarrow H^{n+m}(A), \quad [a] \cdot [b] = [a \cdot b]$$

This is well defined:

$$d(a \cdot b) = \underset{=0}{(da) \cdot b} + (-1)^n \underset{=0}{a \cdot (db)} = 0$$

so $a \cdot b$ is a cycle and we can take its homology class. Let $x \in A^{n-1}$.

$$(a + dx) \cdot b = a \cdot b + (dx) \cdot b = a \cdot b + d(x \cdot b) = [(a + dx) \cdot b] = [a \cdot b]$$

so it only depends on the cohomology class of a , analogous for b .

The product on cohomology inherits associativity and unity with $1 = [1] \in H^0(A)$. We need to see 1 is a cocycle:

$$d(1) = d(1 \cdot 1) = (d1) \cdot 1 + (-1)^0 1 \cdot (d1) = 2 \cdot d(1)$$

and so $d(1) = 0$.

The cup product on the R -cohomology of a simplicial set X is the product induced by the cup product on $C^*(X, R)$ in $H^*(C(X, R)) = H^*(X, R)$.

Theorem 1.6: Properties of the cup-product on homology

Let X be a simplicial set and R a ring. Then

- The cup product on $H^*(X, R)$ is associative and unital, with unit the cohomology class of the constant function $1: X_0 \rightarrow R$.
- For a morphism of simplicial sets $\alpha: Y \rightarrow X$, the relation

$$\alpha^*([x] \cup [y]) = \alpha^*[X] \cup \alpha^*[y]$$

holds for all $[x] \in H^n(X, R), [y] \in H^m(X, R)$.

Remark 1.7. The cup product generalizes to relative cohomology: For A, B simplicial subsets of X . We have

$$C^m(X, A; R) = \{f: X_n \rightarrow R \mid f(A_n) = \{0\}\}$$

The relative cup product is the restriction of \cup on $C^*(X, R)$ to

$$C^m(X, A; R) \times C^m(X, B; R) \xrightarrow{\cup} C^{n+m}(X, A \cup B; R).$$

Let $x \in (A \cup B)_{n+m}$, then

$$(f \cup g)(x) = f(d_{front}^*(x)) \cdot g(d_{back}^*(x))$$

if $x \in A_{n+m}$ then $f(d_{front}^*(x)) = 0$ and analogous with B_{n+m} , anyways the product is 0.

This gives us biadditive well defined maps

$$\cup: H^n(X, A; R) \times H^n(X, B; R) \rightarrow H^{n+m}(X, A \cup B; R)$$

In particular for $A = B$ we get

$$\cup: H^n(X, A; R) \times H^n(X, A; R) \rightarrow H^{n+m}(X, A; R)$$

which is well defined and associative, but not unital anymore.

1.1 Commutativity of the cup-product

Theorem 1.8: Commutativity of the cup-product

Let X be a simplicial set and R a commutative ring. Then for all $[x] \in H^n(X, R); [y] \in H^m(X, R)$ the relation

$$[x] \cup [y] = (-1)^{n \cdot m} \cdot [y] \cup [x]$$

holds.

Schwede points out, that the easy way doesn't work. **Warning.** For $f \in C^n(X, R), g \in C^m(Y, R)$, then in general $f \cup g \neq (-1)^{n+m}(g \cup f)$ in $C^{n+m}(X, R)$. The commutativity is a property we only get on homology.

Construction 1.9. The \cup_1 -product (spoken Cup-one)

$$\cup_1: C^m(X, R) \times C^m(X, R) \rightarrow C^{m+m-1}(X, R)$$

is defined by

$$(f \cup_1 g)(x) = \sum_{i=0}^{n-1} (-1)^{(n-1) \cdot (m+1)} f((d_i^{out})^*(x)) \cdot g((d_i^{inner})^*(x))$$

for $f \in C^n, g \in C^m$ and $x \in X_{n+m-1}$.³ where $d_i^{out}: [n] \rightarrow [n+m-1], d_i^{inner}: [m] \rightarrow [n+m-1]$ are the unique monotone injective maps with images $\text{Im}(d_i^{out}) = \{0, \dots, i\} \cup \{i+m, \dots, n+m-1\}$ and $\text{Im}(d_i^{inner}) = \{i, \dots, i+m\}$.

³There are also \cup_i for $i \in \mathbb{N}$. However, they are quite messy and combinatorial.

Theorem 1.10: \cup_1 -Product

The \cup_1 -product satisfies the following formula

$$d(f \cup_1 g) = (df) \cup_1 g + (-1)^n \cdot f \cup_1 (dg) - (-1)^{n+m} (f \cup g) - (-1)^{n+1} m + 1 (g \cup f)$$

for $f \in C^n(X, R)$ and $g \in C^m(X, R)$.

Remark 1.11. What we want to see, is that $f \cup g$ and $g \cup f$ are not the same but rather homotopic, and \cup_1 witnesses that homotopy.

Proof. This theorem will not be proven, because it is quite messy. You should find a lecture-video for that. \square

Now suppose that f and g are cocycles, i.e. $df = 0$, $dg = 0$. Then

$$d(f \cup_1 g) = -(-1)^{n+m} (f \cup g) - (-1)^{(n+1)(m+1)} (g \cup f)$$

and we get

$$(-1)^{n+m+1} \cdot d(f \cup_1 g) = f \cup g - (-1)^{n \cdot m} (g \cup f)$$

and as such

$$0 = [(-1)^{n+m-1}] = [f] \cup [g] - (-1)^{n \cdot m} [g] \cup [f]$$

Remark 1.12. Last term we discussed the tensor product of two chain complexes (in an exercise):

$$(C \otimes D)_n = \bigoplus_{p+q=n} C_p \otimes D_q$$

and differential

$$d(x \otimes y) = (dx) \otimes y + (-1)^{|x|} \cdot x \otimes (dy)$$

Remark 1.13. Reinterpretation of $d(f \cup_1 g)$. The cup product yields a morphism of cochain complexes

$$C^*(X, R) \otimes C^*(X, R) \rightarrow C^*(X, R)$$

and we get a diagram

$$\begin{array}{ccc} x \otimes y & C^*(X, R) \otimes C^*(X, R) & \xrightarrow{\cup} C^*(X, R) \\ \downarrow & \downarrow & \searrow \cup \\ y \otimes x & C^*(X, R) \otimes C^*(X, R) & \end{array}$$

that does not commute, however it does so up to cochain homotopy and \cup_1 is exactly a cochain homotopy between the two maps.

[07.04.2025, Lecture 1]
[09.04.2025, Lecture 2]

Only with the definition of the cup-product we cannot calculate a lot yet. Some methods to compute cup-products are:

- directly from the definition
- cellular approximation of the diagonal (whatever that means, he gives a little intuition I failed to record.) (this might be used later)
- Group homology (one example later today, something for AT I)

- Poincaré duality (later this term)
- Analysis on smooth manifolds together with De Rahm Cohomology

The first two methods are not very practical.

Example 1.14. Let X be a discrete space, Then $\mathcal{S}(X)$ is a constant simplicial set. The chain complex has the form

$$\xrightarrow{0} \mathbb{Z}[X] \xrightarrow{=} \mathbb{Z}[X] \xrightarrow{0} \mathbb{Z}[X]$$

And so $H^n(X, R) = 0$ for $n \geq 0$. And only for $n = m = 0$ something nontrivial happens. for $f: X_0 \rightarrow R, g: X_0 \rightarrow R$, we have $(f \cup g)(x) = f(d_{front}^*(x)) \cdot g(d_{back}^*(x)) = f(x) \cdot g(x)$ and so the cup product is just pointwise multiplication in dimension 0.

More generally: $H^0(X, R) = \text{maps}(\pi_0(X), R)$ with \cup -prodcut pointwise multiplication

Example 1.15. Let G be a group: Define a category \underline{G}^4 wit one object $*$ and $\text{Hom}_{\underline{G}}(*, *) = G$. We then define

$$BG = N(\underline{G})$$

Where N is the Nerve-Functor $\mathbf{CAT} \rightarrow \mathbf{Sset}$. Then

$$(BG)_n = G^n, \quad d_i^*: G^n \rightarrow G^{n-1}(g_1, \dots, g_n) \mapsto \begin{cases} (g_2, \dots, g_n) & i = 0 \\ (g_1, \dots, g_i \circ g_{i+1}, \dots, g_n) & 1 \leq i \leq n-1 \\ (g_1, \dots, g_{n-1}) & i = n \end{cases}$$

And $s_i(g_1, \dots, g_n) = (g_1, \dots, g_i, 1, g_{i+1}, \dots, g_n)$.

The general case of this is too hard to calculate. We take $G = (\mathbb{F}_2, +)$ and $R = \mathbb{F}_2$ and we calculate $H^*(B\mathbb{F}_2, \mathbb{F}_2)$. We see

$$\begin{array}{ccccccc} C^0(BG, A) & \xrightarrow{d} & C^1(BG, A) & \xrightarrow{d} & C^2(BG, A) & \longrightarrow & \dots \\ \parallel & & \parallel & & \parallel & & \\ \text{maps}(\{1\}, A) & \xrightarrow{0} & \text{maps}(G, A) & \longrightarrow & \text{maps}(G^2, A) & & \\ \parallel & & & & & & \\ A & & (f: G \rightarrow A) & \longrightarrow & (df)(g, h) & & \end{array}$$

And the map is defined by

$$f(d_0^*(g, h)) - f(d_1^*(g, h)) + f(d_2^*(g, h)) = f(h) - f(g \cdot h) + f(g)$$

and

$$df = 0 \Leftrightarrow f(g, h) = f(g) + f(h)$$

\Rightarrow 1-cocycles are the group homomorphisms from G to A

$$H^1(BG, A) \cong \text{Hom}(G, A)$$

and for $G = (\mathbb{F}_2, +)$, $A = \mathbb{F}_2$

We define

$$0 \neq x := [\text{Id}_{\mathbb{F}_2}] \in H^1(B\mathbb{F}_2, \mathbb{F}_2).$$

⁴via geometric realization, these define interesting spaces, namely some (missed word)-MacLane spaces $M(G, 1)$, didn't catch it all

We will show that $x^n = x \cup \dots \cup x$ (n -times) $\in H^n(B\mathbb{F}_2, \mathbb{F}_2)$ is nonzero.

Proposition. $x^n \in H^n(B\mathbb{F}_2, \mathbb{F}_2)$ is represented by

$$f_n: (\mathbb{F}_2)^n \rightarrow \mathbb{F}_2, f_n(\lambda_1, \dots, \lambda_n) = \lambda_1 \cdot \dots \cdot \lambda_n = \begin{cases} 1 & \text{if } \lambda_1 = \lambda_2 = \dots = \lambda_n = 1 \\ 0 & \text{else} \end{cases}$$

Proof. By induction on n . We checked for $n = 1$. For $n \geq 2$ we have

$$\begin{aligned} x^n &= x^{n-1} \cup x = [f_{n-1}] \cup [\text{Id}_{\mathbb{F}_2}] \\ &= [f_{n-1} \cup \text{Id}] \end{aligned}$$

Then

$$\begin{aligned} (f_{n-1} \cup \text{Id})(\lambda_1, \dots, \lambda_n) &= f_{n-1}(d_{\text{front}}^*(\lambda_1, \dots, \lambda_n)) \cdot \text{Id}(d_{\text{back}}^*(\lambda_1, \dots, \lambda_n)) \\ &= f_{n-1}(\lambda_1, \dots, \lambda_{n-1}) \cdot \text{Id}(\lambda_n) \\ &= (\lambda_1 \cdot \dots \cdot \lambda_{n-1}) \cdot \lambda_n \end{aligned}$$

□

Claim: $x^n \neq 0$. In the UCT for cohomology we used the evaluation pair

$$\Phi: H^n(X, A) \rightarrow \text{Hom}(H_n(X; \mathbb{Z}); A), \quad [f_n: X_n \rightarrow A] \mapsto \left\{ \left[\sum b_i x_i \right] \mapsto \sum b_i f(x_i) \right\}$$

for $b_i \in \mathbb{Z}, x_i \in X_n$. We can slightly vary that for ring coefficients:

$$\Phi: H^n(X, R) \rightarrow \text{Hom}(H_n(X, R), R)$$

and $[f: X_n \rightarrow R] \mapsto \left\{ \left[\sum r_i \cdot x_i \right] \mapsto \sum r_i \cdot f(x_i) \right\}$ with $r_i \in R, x_i \in X_n$.

With $X = B\mathbb{F}_2, R = \mathbb{F}_2$, we consider

$$y := \sum_{(\lambda_1, \dots, \lambda_n) \in (\mathbb{F}_2)^n} 1(\lambda_1, \dots, \lambda_n) \in \mathbb{F}_2[(\mathbb{F}_2)^n] = \mathbb{F}_2[(B\mathbb{F}_2)_n]$$

Claim: y is an n -cycle in $C_*(B\mathbb{F}_2, \mathbb{F}_2)$.

$$\begin{aligned} dy &= \sum_{i=0, \dots, n} (-1)^i \cdot d_i^* \left(\sum_1 \cdot (\lambda_1, \dots, \lambda_n) \right) \\ &= \sum_{i=0, \dots, n} \underbrace{\sum_{(\lambda_1, \dots, \lambda_n) \in \mathbb{F}_2^n} (-1)^i \cdot d_i^*(\lambda_1, \dots, \lambda_n)}_{\text{cancel in pairs}} \\ &= 0 \end{aligned}$$

Now

$$d_0^*(0, \lambda_2, \dots, \lambda_n) = (\lambda_2, \dots, \lambda_n) = d_0^*(1, \lambda_2, \dots, \lambda_n)$$

So

$$\Phi(x^n): H_n(B\mathbb{F}_2, \mathbb{F}_2) \rightarrow \mathbb{F}_2$$

$$\Phi(x^n)[y] = \Phi[f_n] \left[\sum_{(\lambda_1, \dots, \lambda_n) \in \mathbb{F}_2^n} (\lambda_1, \dots, \lambda_n) \right] = \sum_{(\lambda_1, \dots, \lambda_n)} f_n(\lambda_1, \dots, \lambda_n) = \sum_{(\lambda_1, \dots, \lambda_n)} \lambda_1 \cdot \dots \cdot \lambda_n = 1 \neq 0$$

and $[y] \neq 0$ in $H_n(B\mathbb{F}_2, \mathbb{F}_2)$.

We will later see, that in fact $H^*(B\mathbb{F}_2; \mathbb{F}_2) = \mathbb{F}_2[X]$.

Remark. Let p be an odd prime. $H^*(B\mathbb{F}_p, \mathbb{F}_p) = ?$.

$$0 \neq x = [\text{Id}_{\mathbb{F}_p} \in H^1(B\mathbb{F}_p; \mathbb{F}_p)]$$

still makes sense, but now there are more scalars and

$$x^n = 0$$

for $n \geq 2$. The graded commutativity says:

$$x \cup x = (-1)^{1 \cdot 1} x \cup x = -x \cup x$$

so if R is commutative, $x \in H^n(X, R)$ and n is odd, then $2 \cdot (x \cup x) = 0$ in $H^{2n}(X, R)$. And then $2 \cdot x^2 = 0 \Rightarrow x^2 = 0$.

Define $h: \mathbb{F}_p \times \mathbb{F}_p \rightarrow \mathbb{F}_p$ by

$$h(i, j) = \begin{cases} 0 & \text{if } i + j < p \\ 1 & \text{if } i + j \geq p \end{cases}$$

where we write $\mathbb{F}_p = \{0, \dots, p-1\}$. Now $h \in C^2(B\mathbb{F}_p, \mathbb{F}_p)$. Fact: $dh = 0$ and $0 \neq y := [h] \in H^2(B\mathbb{F}_p, \mathbb{F}_p)$.

We then get (but do not prove)

$$H^*(B\mathbb{F}_p, \mathbb{F}_p) = \Lambda(x) \otimes \mathbb{F}_p[y]$$

and

$$H^{2n}(B\mathbb{F}_p, \mathbb{F}_p) = \mathbb{F}_p\{y^n\}, \quad H^{2n+1}(B\mathbb{F}_p, \mathbb{F}_p) = \mathbb{F}_p\{xy^n\}$$

Chapter 2

Künneth theorem

The Künneth theorem is an algebraic relationship between $H_*^*(X, R)$, $H_*^*(Y, R)$ and $H_*^*(X \times Y, R)$ ¹.

Here is a simplest version in homology with field coefficients:

Theorem 2.1: Künneth, simple version

Let X and Y be spaces and k a field. Then

$$H_n(X \times Y, k)$$

is natural isomorphic to

$$\bigoplus_{p+q=n} H_p(X, k) \otimes_k H_q(Y, k)$$

2.1 The Eilenberg-Zilber-theorem

Let A, B be simplicial abelian groups. Then we get two natural chain homotopy equivalences

$$\begin{array}{ccc} & \text{Eilenberg-Zilber-Map} & \\ & \curvearrowright & \\ C_*(A) \otimes C_*(B) & & C_*(A \otimes B) \\ & \curvearrowleft & \\ & \text{Alexander Whitney map} & \end{array}$$

Definition 2.2: Simplicial abelian group

A *simplicial abelian group* is a functor $A: \Delta^{op} \rightarrow \mathbf{Ab.Groups}$.

Remark. Equivalently a simplicial abelian group is a collection of abelian groups A_n , and homomorphisms $\alpha^*: A_m \rightarrow A_n$ for all $\alpha: [n] \rightarrow [m]$ in Δ , s.t. $(\alpha \circ \beta)^* = \beta^* \circ \alpha^*$.

Equivalently a simplicial abelian group is a simplicial set endowed with abelian group structure on the sets of n -simplices, such that all α^* are homomorphisms.

Example 2.3. Let X be a simplicial set and A an abelian group. Then the composite

$$\Delta^{op} \xrightarrow{X} (\mathbf{Sets}) \xrightarrow{A[_]} (\mathbf{ab.grps})$$

$A[X]$

¹ H_*^* denotes, that Schwede was too lazy to write the statement for homology and cohomology separately

is a simplicial abelian group.

Construction 2.4. Let $A: \Delta^{op} \rightarrow (\mathbf{ab.grps})$ be a simplicial abelian groups. Its *chain complex* $C_*(A)$ is the chain complex with $C_n(A) = A_n$ with differential

$$d: C_n(A) = A_n \rightarrow A_{n-1} = C_{n-1}(A), \quad d(a) = \sum_{i=0, \dots, n} (-1)^i d_i^*(a)$$

And one can easily check $d \circ d = 0$.

Note. The following commutes

$$\begin{array}{ccc} (\mathbf{Ssets}) & \xrightarrow{X \mapsto C_*(X, A)} & (\mathbf{Chains}) \\ & \searrow A[_] \quad \nearrow C_* & \\ & (\mathbf{s.ab.grps}) & \end{array}$$

Remark 2.5. The tensor product of chain complexes C, D is

$$(C \otimes D)_n := \bigoplus_{p+q=n} C_p \otimes D_q$$

with differential

$$d(x \otimes y) = (dx \otimes y) + (-1)^p x \otimes (dy)$$

for $x \in C_p, y \in D_q$.

We can also form the tensor product of simplicial abelian groups:

Definition 2.6: Tensor product of simplicial abelian groups

$A, B: \Delta^{op} \rightarrow (\mathbf{ab.grps})$ by

$$(A \otimes B)_n = A_n \otimes B_n, \quad \alpha^*: (A \otimes B)_n \rightarrow (A \otimes B)_m$$

for $\alpha: [m] \rightarrow [n]$ is defined as $\alpha^*(a \otimes b) = \alpha^*(a) \otimes \alpha^*(b)$ and we write $\alpha_{A \otimes B}^* := \alpha_A^* \otimes \alpha_B^*$. This can be equally described as the composite

$$\Delta^{op} \xrightarrow{(A, B)} (\mathbf{ab.grps}) \times (\mathbf{ab.grps}) \xrightarrow{\otimes} (\mathbf{ab.grps})$$

Warning. For $A, B \in (\mathbf{SAB}) = \text{simplicial abelian groups}$

$$C_*(A \otimes B) \neq C_*(A) \otimes C_*(B)$$

Also he did this in dimension n , but I lacked time to copy.

The Eilenberg-Zilber theorem is a natural pair of chain homotopy equivalences between these two.

Construction 2.7. Let A, B be simplicial chain groups. The *Alexander-Whitney map* is the chain map

$$AW: C_*(A \otimes B) \rightarrow C_*(A) \otimes C_*(B)$$

defined by

$$\begin{array}{ccc}
 C_n(A \otimes B) & \longrightarrow & \bigoplus_{p+q=n, p, q \geq 0} A_p \otimes B_q \\
 \parallel & & \parallel \\
 A_n \otimes B_n & & C_*(A) \otimes C_*(B) \\
 AW_n(a \otimes b) = \sum_{p+q=n} d_{front}^*(a) \otimes d_{back}^*(b)
 \end{array}$$

Where $[p] \xrightarrow{d_{front}} [p+q] = [n] \xleftarrow{d_{back}} [q]$.

You may check for yourself, that this is a chain map, however Schwede didn't do that.

[09.04.2025, Lecture 2]

[14.04.2025, Lecture 3]

Remark. An example for a simplicial abelian group, that is not of the form

$$\Delta^{op} \xrightarrow{X} \mathbf{sets} \xrightarrow{A[-]} (\mathbf{ab.grps.})$$

is for any abelian group G the simplicial set BG , that also admits structure of a simplicial abelian group.

Remark 2.8 (Relation between AW-map and cup-product). For a simplicial set X and ring R ,

$$C^*(X, R) = \text{Hom}(C_*(X, \mathbb{Z}), R) = \text{Hom}(C_*(\mathbb{Z}[X]), R)$$

and $C^n(X, R) = \text{Hom}(C_n(X, \mathbb{Z}), R)$. If $\psi \in C^n(X, R)$ is a cocycle, i.e. $d(\psi) = 0$, then it extends to a chain map

$$\tilde{\psi}: C_*(\mathbb{Z}[X]) \rightarrow R[n]$$

where $R[n]$ is the complex with R in dimension n and 0 otherwise. and $\tilde{\psi}$ is ψ in dimension n and 0 otherwise.

For $f \in C^n(X, R), g \in C^m(X, R)$ cocycles, we have $f \cup g \in C^{n+m}(X, R)$. Then $f \tilde{\cup} g$ is the following composite

$$\begin{array}{ccccc}
 C_*(\mathbb{Z}[X]) & \xrightarrow{C_*(\mathbb{Z}[\text{diagonal}])} & C_*(\mathbb{Z}[X \times X]) & \cong & C_*(\mathbb{Z}[X] \otimes \mathbb{Z}[X]) \\
 & & \searrow \text{AW} & & \\
 C_*(\mathbb{Z}[X]) \otimes C_*(\mathbb{Z}[X]) & \xleftarrow{\tilde{f} \otimes g} & R[n] \otimes R[m] & \xrightarrow{\text{mult}} & R[n+m]
 \end{array}$$

Definition 2.9: (p,q)-shuffle

A (p, q) -shuffle for $p, q \geq 0$ is a permutation σ of $\{0, 1, \dots, p+q-1\}$, such that the restriction of σ to $\{0, 1, \dots, p-1\}$ is monotone, and the restriction of σ to $\{p, \dots, p+q-1\}$ is monotone.

Remark. „Shuffles leave the first p elements in order and the last q elements in order.“

Example 2.10. The only $(p, 0)$ -shuffle or $(0, q)$ -shuffles are the identity.

There are precisely two $(1, 1)$ -shuffles, namely both permutations of $\{0, 1\}$.

$\sigma \in S_3$ given by $\sigma(0) = 0, \sigma(1) = 2, \sigma(2) = 1$ is not a $(2, 1)$ -shuffle, but it is a $(1, 2)$ -shuffle.

Remark 2.11. (p, q) -shuffles biject with p -element subsets of $\{0, 1, \dots, p + q - 1\}$ by $\sigma \mapsto \{\sigma(0), \dots, \sigma(p)\}$ and also with q -element subsets of $\{0, 1, \dots, p + q - 1\}$ by $\sigma \mapsto \{\sigma(p), \dots, \sigma(p + q - 1)\}$.

This means $|(p, q)\text{-shuffles}| = \binom{p+q}{p} = \binom{p+q}{q}$.

Notation 2.12. Let σ be a (p, q) -shuffle. We write $\mu_i := \sigma(i - 1)$ for $1 \leq i \leq p$ and $\nu_i := \sigma(p + i - 1)$ for $1 \leq i \leq q$.

This means $0 \leq \mu_1 \leq \dots \leq \mu_p$ and $0 \leq \nu_1 \leq \dots \leq \nu_q \leq p + q - 1$.

Definition 2.13: Eilenberg-Zilber map

Let A, B be simplicial abelian groups. The Eilenberg-Zilber map /shuffle map is

$$EZ: C_*(A) \otimes C_*(B) \rightarrow C_*(A \otimes B)$$

is the direct sum of the homomorphisms

$$\nabla_{p,q}: A_p \otimes B_q \rightarrow A_{p+q} \otimes B_{p+q}$$

given by

$$a \otimes b \mapsto a \nabla b := \sum_{\sigma: (p,q)\text{-shuffle}} \text{sgn}(\sigma) \cdot (s_{\nu_i} \circ \dots \circ s_{\nu_q})^*(a) \otimes (s_{\mu_1} \circ \dots \circ s_{\mu_p})^*(b)$$

Example 2.14. There is only one $(p, 0)$ -shuffle, the identity of $\{0, \dots, p - 1\}$. Then $\mu_i = i - 1$.

$$\nabla_{p,0}: A_p \otimes B_0 \rightarrow A_p \otimes B_p$$

is defined by

$$a \otimes b \mapsto a \nabla b = a \otimes (s_0 \circ \dots \circ s_{p-1})^*(b).$$

For $p = q = 1$ i didn't have the time to copy.

Schwede claims, that the Eilenberg-Zilber map is a chain map and he can't believe he actually did those calculations 4 years ago. He will not torture us, but you may watch the videos.

Theorem 2.15: Shuffle maps form a chain map

The shuffle maps $\nabla_{p,q}$ for varying $p, q \geq 0$ assemble into a chain map. Furthermore, for $a \in A_p, b \in B_q$

$$d(a \nabla b) = (da) \nabla b + (-1)^p a \nabla (db)$$

He specifies, that the calculation takes up 8 pages of his notes.

Theorem 2.16: Eilenberg-Zilber

Let A, B be simplicial abelian groups. Then the morphisms

$$\begin{array}{ccc} & \xrightarrow{\text{Eilenberg-Zilber}} & \\ C_*(A) \otimes C_*(B) & & C_*(A \otimes B) \\ & \xleftarrow{AW} & \end{array}$$

are mutually inverse natural chain homotopy equivalences.

A first method of proof would be explicit formulas for the chain homotopies $AW \circ EZ \sim \text{Id}$ and $EZ \circ AW \sim \text{Id}$. That is however infinitely annoying and we will not do this.

For the special case, where $A = \mathbb{Z}[X], B = \mathbb{Z}[Y]$ for simplicial sets X, Y we prove this via acyclic models. For that we need some category-theory:

2.2 Yoneda Lemma & Acyclic models

Theorem 2.17: Yoneda lemma

Let \mathcal{C} be a category and c an object of \mathcal{C} . Let $F: \mathcal{C} \rightarrow (\mathbf{sets})$ be a functor: Then the evaluation map

$$\text{Nat}_{\mathcal{C} \rightarrow \mathbf{sets}}(\mathcal{C}(c, _), F) \rightarrow F(c)$$

given by

$$(\tau: \mathcal{C}(c, _) \rightarrow F) \mapsto (\tau_c: \mathcal{C}(c, c) \rightarrow F(c))(\text{id}_c)$$

is bijective.

Equally: for every $x \in F(c)$, there is a unique natural transformation $\tau: (\mathcal{C}(c, _) \rightarrow F)$, such that $\tau_c(\text{id}_c) = x$.

Remark. A special case of this is

$$\text{Hom}_{\mathbf{sset}}(\Delta^n, X) \cong X_n, \quad (f: \Delta^n \rightarrow X) \mapsto f_n(\text{id}_{[n]}).$$

where $\Delta^n = \Delta(_, [n])$.

Proof. We show injectivity and surjectivity.

Injectivity Let $\tau: \mathcal{C}(c, _) \rightarrow F$ be any natural transformation. Let d be another object of \mathcal{C} , $f: c \rightarrow d$ any morphism. Then we have

$$\tau_d: \mathcal{C}(c, d) \rightarrow F(d)$$

and

$$\tau_d(f: c \rightarrow d) = \tau_d(\mathcal{C}(c, f)(\text{id}_c)) = F(f)(\tau_c(\text{id}_c))$$

where we use naturality of τ :

$$\begin{array}{ccc} \mathcal{C}(c, d) & \xrightarrow{\tau_d} & F(d) \\ \downarrow \mathcal{C}(c, g) & & \downarrow F(g) \\ \mathcal{C}(c, e) & \xrightarrow{\tau_e} & F(e) \end{array}$$

which implies the value of τ at $d, f: c \rightarrow d$ is determined by its value of (c, id_c) and the functoriality of F .

Surjectivity Let $y \in F(c)$ be given. For an object d of \mathcal{C} and morphism $f: c \rightarrow d$, we define

$$\tau_d: \mathcal{C}(c, d) \rightarrow F(d) \quad \tau_d(f) := F(f)(y).$$

We check $\tau_c(\text{id}_c) = F(\text{id}_c)(y) = y$. We need to check for naturality. Let $g: d \rightarrow e$ be another morphism. Then

$$\begin{aligned} F(g)(\tau_d(f)) &= F(g)(F(f)(y)) = F(g \circ f)(y) \\ &= \tau_e(g \circ f) = \tau_e(\mathcal{C}(c, g)(f)) \end{aligned}$$

□

Let \mathcal{C} be a category, c an object of \mathcal{C} . We define the functor $\mathbb{Z}[\mathcal{C}(c, _)] : \mathcal{C} \rightarrow (\mathbf{ab.grps.})$ as the composite

$$\mathcal{C} \xrightarrow{\mathcal{C}(c, _)} (\mathbf{sets}) \xrightarrow{\mathbb{Z}[_]} (\mathbf{ab.grps.}).$$

In particular, $\mathbb{Z}[\mathcal{C}(c, _)](d) = \mathbb{Z}[\mathcal{C}(c, d)]$.

Proposition (Additive Yoneda lemma). Let $c \in \text{ob}(\mathcal{C})$, $F : \mathcal{C} \rightarrow (\mathbf{ab.grps.})$ any functor. Then the evaluation map

$$\text{Nat}_{\mathcal{C} \rightarrow (\mathbf{ab.grps.})}(\mathbb{Z}[\mathcal{C}(c, _)], F) \rightarrow F(c)$$

is bijective. $(\tau : \mathbb{Z}[\mathcal{C}(c, _)] \rightarrow F) \mapsto \tau_c(1 \cdot \text{id}_c)$.

Proof. For varying objects d of \mathcal{C} , the bijections

$$\text{Hom}_{AB}(\mathbb{Z}[\mathcal{C}(c, d)], F(d)) \cong \text{Hom}_{\mathbf{sets}}(\mathcal{C}(c, d), F(d))$$

assemble into a bijection²

$$\text{Nat}_{\mathcal{C} \rightarrow \mathbf{Ab}}(\mathbb{Z}[\mathcal{C}(c, _)], F) \cong \text{Nat}_{\mathcal{C} \rightarrow \mathbf{sets}}(\mathcal{C}(c, _), F) \xrightarrow{\text{Yoneda}} F(c)$$

□

Definition 2.18: Representable functor

A functor $F : \mathcal{C} \rightarrow \mathbf{Ab}$ is representable if there is an object $c \in \mathcal{C}$ and a natural isomorphism $F \cong \mathbb{Z}[\mathcal{C}(c, _)]$

Note. Any isomorphism $F \cong \mathbb{Z}[\mathcal{C}(c, _)]$ is determined by the „universal element“ in $F(c)$.

Example 2.19. Let $\mathcal{C} = (\mathbf{ssets}) \times (\mathbf{ssets})$ be the product of two copies of the category of simplicial sets. Define $f : (\mathbf{ssets}) \times (\mathbf{ssets}) \rightarrow \mathbf{Ab}$ given by $F(X, Y) = \mathbb{Z}[X_p \times Y_q]$ for some $p, q \geq 0$. **Claim.** This functor is representable by (Δ^p, Δ^q) with natural isomorphisms.

$$(\mathbf{ssets} \times \mathbf{ssets})((\Delta^p, \Delta^q), (X, Y)) = \mathbf{sets}(\Delta^p, X) \times \mathbf{sets}(\Delta^q, Y) \cong X_p \times Y_q$$

Apply free abelian groups to get

$$\mathbb{Z}[(\mathbf{ssets} \times \mathbf{ssets})((\Delta^p, \Delta^q), (X, Y))] \cong \mathbb{Z}[X_p \times Y_q]$$

Notation 2.20. For $F : \mathcal{C} \rightarrow \mathbf{Chains}$ we write $F_n = (_)_n \circ F : \mathcal{C} \rightarrow \mathbf{Ab}$ as the composite.

$$\mathcal{C} \xrightarrow{F} \mathbf{Chains} \xrightarrow{(_)_n} \mathbf{Ab}$$

and the second map sends $C = C(n, d_n)_{n \in \mathbb{Z}} \mapsto C_n$.

²I don't know why though.

Theorem 2.21: Acyclic models

Let \mathcal{C} be a category, $F, G: \mathcal{C} \rightarrow \mathbf{Chains}_+$ = non-negative grade chain complexes. Let $\psi: F \rightarrow G$ be a natural transformation of functors. Suppose;

1. The transformation $\psi_0: F_0 \rightarrow G_0: \mathcal{C} \rightarrow \mathbf{Ab}$ is the zero natural transformation
2. For every $n \geq 1$, the functor $F_n: \mathcal{C} \rightarrow \mathbf{Ab}$ is isomorphic to a direct sum of representable functors, $\bigoplus_{i \in I} \mathbb{Z}[\mathcal{C}(c_i, _)]$ for some family $\{c_i\}_{i \in I}$ of \mathcal{C} -objects such that $H_n(G(c_i)) = 0$.

Then ψ is naturally chain nullhomotopic.

[14.04.2025, Lecture 3]
[16.04.2025, Lecture 4]

Proof. For $n \geq 0$, we will construct natural transformations

$$s_n: F_n \rightarrow G_{n+1}$$

of functors $\mathcal{C} \rightarrow \mathbf{Ab}$, such that

$$d_{n+1} \circ s_n + s_{n-1} \circ d_n = \psi_n \quad (*)$$

as natural transformations (i.e. they have the chain homotopy property).

The construction is by induction on n . We begin with $s_0 = 0$ and $s_{-1} = 0$. Suppose $n \geq 1$ and that s_0, \dots, s_{n-1} have been constructed satisfying (*). Then

$$d_n^G \circ (\psi_n - s_{n-1} \circ d_n^F) = d_n^G \circ \psi_n - d_n^G \circ s_{n-1} \circ d_n^F$$

as ψ is a chain map,

$$= \psi_{n-1} \circ d_n^F - d_n^G \circ s_{n-1} \circ d_n^F = (\psi_{n-1} - d_n^G \circ s_{n-1}) \circ d_n^F \stackrel{(*)}{=} s_{n-2} \circ d_{n-1}^F \circ d_n^F = 0.$$

So $\psi_n - s_{n-1} \circ d_n^F: F_n \rightarrow G_n$ takes values in cycles. By 2.,

$$f_n = \bigoplus_{i \in I} \mathbb{Z}[\mathcal{C}(c_i, _)]$$

for some set $\{c_i\}_{i \in I}$ of \mathcal{C} -objects, such that $H_n(G(c_i)) = 0$. Let $j \in I$, write

$$x_j \in F(c_j) = \bigoplus_{i \in I} \mathbb{Z}[\mathcal{C}(c_i, c_j)]$$

be the element $1 \cdot \text{id}_j$ in the j -th summand. Then

$$\psi_n^{c_j}(x_j) - s_{n-1}^{c_j}(d_n^{F, c_j}(x_j)) \in G_n(c_j)$$

is a cycle. Since $H_n(G(c_j)) = 0$, the class is a boundary in the complex $G(c_j)$.

Let $y_j \in G(c_j)_{n+1}$ be a element such that

$$d_{n+1}^{c_j}(y_j) = \psi_n^{c_j}(x_j) - s_{n-1}^{c_j}(d_n^{F, c_j}(x_j))$$

The additive Yoneda lemma provides a unique natural transformation

$$s_{n,j}: \mathbb{Z}[\mathcal{C}(c_j, _)] \rightarrow G_{n+1}$$

such that $s_{n,j}(x_j) = s_{n,j}^{c_j}(1 \cdot \text{id}_{c_j}) = y_j \in G_{n+1}(c_j)$.

We define the natural transformation

$$s_n: F_n = \bigoplus_{i \in I} \mathbb{Z}[\mathcal{C}(c_i, _)] \rightarrow G_{n+1}$$

as $s_n = \bigoplus_{j \in I} s_{n,j}$.

It suffices now to show, that $(*)$ holds on each summand $\mathbb{Z}[\mathcal{C}(c_j, _)]$. By the additive Yoneda lemma, there it suffices to check the relation on $1 \cdot \text{id}_{c_j}$, which holds by definition. \square

Remark. We only proved „half“ of the acyclic models theorem. The other half states:

Let \mathcal{C} and $F, G: \mathcal{C} \rightarrow \mathbf{Chains}_+$ be as before, satisfying 2.. Then any natural transformation $\psi_0: F_0 \rightarrow G_0$ can be extended to a natural transformation $\psi: F \rightarrow G$.

Proof of theorem 2.16. Now to actually prove the Eilenberg-Zilber-Theorem 2.16 (at least in a special case.) Let A, B be simplicial abelian groups. We assume $A = \mathbb{Z}[X]$, $B = \mathbb{Z}[Y]$ for some simplicial sets X, Y . We write $C_*(X), C_*(Y)$. For sets S, T ,

$$\begin{array}{ccc} & \curvearrowright & \\ \mathbb{Z}[S] \otimes \mathbb{Z}[T] & & \mathbb{Z}[S \times T] \\ & \curvearrowleft & \\ s \otimes t & \longrightarrow & (s, t) \end{array}$$

is naturally isomorphic. Dimensionwise this gives $\mathbb{Z}[X] \otimes \mathbb{Z}[Y] \cong \mathbb{Z}[X \times Y]$.

We want to move this further to $C_*(X) \otimes C_*(Y) \cong C_*(X \times Y)$.

Proposition 2.22.

1. For all $p \geq 0$, the simplicial set Δ^p is simplicially contractible.
2. For all $p \geq 0$, the complex $C_*(\Delta^p)$ is chain homotopy equivalent to the complex $\mathbb{Z}[0]$, the complex consisting of \mathbb{Z} in dimension 0.
3. For $p, q \geq 0$, the chain complex $C_*(\Delta^p) \otimes C_*(\Delta^q)$ is chain homotopy equivalent to $\mathbb{Z}[0]$. In particular,

$$H_n(C_*(\Delta^p) \otimes C_*(\Delta^q)) = 0$$

for $n > 0$.

Proof.

1. We define a morphism of simplicial sets $H: \Delta^p \times \Delta^1 \rightarrow \Delta^p$ that contracts Δ^p to the last vertex.³ In dimension n ,

$$H_n: \Delta([n], [p]) \times \Delta([n], [1]) \rightarrow \Delta([n], [p])$$

is given by

$$H_n(\alpha, \beta)(i) = \begin{cases} \alpha(i) & \text{if } \beta(i) = 0 \\ p & \text{if } \beta(i) = 1 \end{cases}$$

for $0 \leq i \leq n$. Let $\gamma: [m] \rightarrow [n]$ be any morphism in Δ . Then

$$H_m(\gamma^*(\alpha, \beta))(j) = H_m(\alpha \circ \gamma, \beta \circ \gamma)(j) = \begin{cases} \alpha(\gamma(j)) & \text{if } \beta(\gamma(j)) = 0 \\ p & \text{if } \beta(\gamma(j)) = 1 \end{cases} = H_n(\alpha, \beta)(\gamma(j)) = \gamma^*(H_n(\alpha, \beta)(j))$$

³remember, that Homotopy is not symmetric in Simplicial sets. This is such an example.

This means H is a homotopy from Id_{Δ^p} to the composite

$$\Delta^p \rightarrow \Delta^0 \xrightarrow{p\text{-th vertex}} \Delta^p$$

2. $C_*: \mathbf{ssets} \rightarrow \mathbf{chains}$ takes simplicial homotopies to chain homotopies. So we know $C_*(\Delta^p)$ is chain homotopy equivalent to

$$C_*(\Delta^0) = (\dots \mathbb{Z} \xrightarrow{\text{Id}} \mathbb{Z} \xrightarrow{0} \mathbb{Z})$$

which is chain homotopy equivalent to

$$(\dots 0 \rightarrow 0 \rightarrow 0 \rightarrow 0 \rightarrow \mathbb{Z}) = \mathbb{Z}[0]$$

3. The tensor product of chain complexes preserves chain homotopy equivalences in each variable separately. So

$$C_*(\Delta^p) \otimes C_*(\Delta^q) \sim \mathbb{Z}[0] \otimes C_*(\Delta^1) \sim \mathbb{Z}[0] \otimes \mathbb{Z}[0] \cong \mathbb{Z}[0].$$

□

We now must produce natural chain homotopies from

$$\mathbf{AW} \circ \mathbf{EZ}: C_*(X) \otimes C_*(Y) \rightarrow C_*(X) \otimes C_*(Y)$$

and

$$\mathbf{EZ} \circ \mathbf{AW}: C_*(X \times Y) \rightarrow C_*(X \times Y)$$

to the respective identities.

Claim. $\mathbf{AW} \circ \mathbf{EZ} - \text{Id}_{C_*(X) \otimes C_*(Y)}: C_*(X) \otimes C_*(Y) \rightarrow C_*(X) \otimes C_*(Y)$ satisfies the hypothesis of acyclic models.

Proof.

$$\begin{array}{ccc} C_0(X) \otimes C_0(Y) & \cong & \mathbb{Z}[X_0] \otimes \mathbb{Z}[Y_0] \\ \parallel & & \uparrow \cong \\ (C_*(X) \otimes C_*(Y))_0 & & \mathbb{Z}[X_0 \times Y_0] \\ & & \downarrow \cong \\ & & C_0(X \times Y) \end{array}$$

Which means $(\mathbf{AW} \circ \mathbf{EZ})_0 = \text{Id}$ and $(\mathbf{EZ} \circ \mathbf{AW})_0 = \text{Id}$. which means $\psi_0 = \text{zero natural transformation}$.

$$(C_*(X) \otimes C_*(Y))_n = \bigoplus_{p+q=n} C_p(X) \otimes C_q(Y) = \bigoplus_{p+q=n} \mathbb{Z}[X_p] \otimes \mathbb{Z}[Y_q] \cong \bigoplus_{p+q=n} \mathbb{Z}[X_p \times Y_q]$$

which is represented by (Δ^p, Δ^q) . Then $H_n(C_*(\Delta^p \otimes \Delta^q)) = 0$ (I think, he erased before I could copy.)

We consider $\phi: \mathbf{EZ} \circ \mathbf{AW} - \text{Id}_{C_*(X \times Y)}: C_*(X \times Y) \rightarrow C_*(X \times Y)$. We know, $\phi_0 = 0$. We need to show, that ϕ satisfies the hypothesis of acyclic models.

$$C_n(X \times Y) = \mathbb{Z}[X_n \times Y_n]$$

is representable by (Δ^n, Δ^n) .

$$H_n(C_*(\Delta^n \times \Delta^n)) \cong H_n(\Delta^0 \times \Delta^0) = H_n(\Delta^0) = 0$$

for $n > 0$, where we used $\Delta^n \sim \Delta^0$ and so $\Delta^n \times \Delta^n \sim \Delta^0 \times \Delta^0$. So acyclic models produces a natural chain nullhomotopy of ϕ . \square

This concludes the proof of the Künneth theorem. \square

2.3 Revisiting Commutativity of the cup-product

The symmetry isomorphism of chain complexes C, D is the morphism.

$$\tau_{C,D}: C \otimes D \xrightarrow{\cong} D \otimes C$$

is given by

$$\begin{aligned} \tau_{C,D_n} &: (C \otimes D)_n && (D \otimes C)_n \\ &\oplus_{p+q=n} C_p \otimes D_q && \oplus_{q+p=n} D_q \otimes C_p \\ &c \otimes d && (-1)^{pq} \cdot d \otimes c \end{aligned}$$

me not being able to keep up.

Fact.

$$\begin{array}{ccc} C_*(X) \otimes C_*(Y) & \xrightarrow{\mathbf{EZ}} & C_*(X, Y) \\ \downarrow \tau & & \downarrow C_*(flip) \\ C_*(Y) \otimes C_*(X) & \xrightarrow{\mathbf{EZ}} & C_*(Y \otimes X) \end{array}$$

commutes. where $flip: X \times Y \rightarrow Y \times X$, $(x, y) \mapsto (y, x)$. Hence, „The Eilenberg-Zilber map is symmetric“.

But however for AW the same diagram does NOT commute. Another diagram is missing.

However it does so up to natural chain homotopy by applying the acyclic models to the difference of the two composites. He explains, why we can apply acyclic models.

Let X be a simplicial set. The diagonal $\Delta: X \rightarrow X \times X$ is flip-invariant, i.e.

$$\begin{array}{ccc} X & \xrightarrow{\Delta} & X \times X \\ & \searrow \Delta & \downarrow flip \\ & & X \times X \end{array}$$

We draw a diagram:

$$\begin{array}{ccccc} C_*(X) & \xrightarrow{C_*(\Delta)} & C_*(X \times X) & \xrightarrow{\mathbf{AW}} & C_*(X) \otimes C_*(X) \\ & \searrow C_*(\Delta) & \downarrow C_*(flip) & & \downarrow \tau \\ & & C_*(X \times X) & \xrightarrow{\mathbf{AW}} & C_*(X) \otimes C_*(X) \end{array}$$

that commutes up to homotopy. We apply the functor $\text{Hom}(_, R)$ to get a new diagram and my speed at copying was not capable of keeping up. You may want to have a look at the videos for this.

[23.04.2025, Lecture 5]

The Plan for today is to show the Künneth theorem for homology. The rough approximation is, that product of spaces goes to Tensorproducts of abelian groups.

2.4 Algebraic Künneth theorem.

If X, Y are simplicial sets, then by EZ we have $H_*(X \times Y; R) = H_*(C_*(X \times Y; R)) \cong H_*((C_*(X, R)) \otimes_R C_*(Y, R))$ and we want to see how that relates to $H_*(X, R) \otimes_R H_*(Y, R)$. This will be the algebraic Künneth theorem.

In the following R is a commutative ring (have integers and fields in mind).

Definition 2.23: Tensor Product of R -chains

Let C, D be chain complexes of R -modules. We define a new complex of R -modules $C \otimes_R D$:

$$(C \otimes_R D)_n = \bigoplus_{p+q=n} C_p \otimes_R D_q$$

with differential

$$d(x \otimes y) = dx \otimes y + (-1)^{pq} x \otimes dy.$$

Note that $R \otimes \mathbb{Z}[S] \cong R[S]$ for S a simplicial set. And $R[S] \otimes_R R[T] \cong R[S \times T]$ for S, T simplicial sets.

For X, Y simplicial sets, we have

$$R \otimes C_*(X, \mathbb{Z}) \otimes C_*(Y, \mathbb{Z}) \xrightarrow{R \otimes \mathbf{EZ}} R \otimes C_*(X \times Y; \mathbb{Z}) \cong C_*(X \otimes Y; R)$$

and for $R \otimes C_*(X; \mathbb{Z}) \otimes C_*(Y; \mathbb{Z}) \cong (R \otimes C_*(X; \mathbb{Z})) \otimes_R (R \otimes C_*(Y; \mathbb{Z})) = C_*(X, R) \otimes_R C_*(Y, R)$, so we get a Eilenberg-Zilber map

$$C_*(X, R) \otimes_R C_*(Y, R) \xrightarrow{\mathbf{EZ}} C_*(X \times Y; R)$$

Aim. relate $H_*(C \otimes_R D)$ to $H_*(C), H_*(D)$. Our hope is to have a map

$$\bigoplus_{p+q=n} H_p(C) \otimes_R H_q(D) \xrightarrow{???} H_n(C \otimes_R D)$$

For example taking $R = \mathbb{Z}$ and $C = D = (\dots, \rightarrow 0 \rightarrow \mathbb{Z} \xrightarrow{\cdot 2} \mathbb{Z} \rightarrow 0)$. Then

$$H_n(C) = H_n(D) = \begin{cases} \mathbb{Z}/2 & n = 0 \\ 0 & n \neq 0 \end{cases}$$

but $C \otimes D = (0 \rightarrow 0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Z}^2 \rightarrow \mathbb{Z} \rightarrow 0)$. And

$$H_1(C \otimes D) = \{(x, -x) \in \mathbb{Z}\} / \{(2y, -2y) \mid y \in \mathbb{Z}\} \cong \mathbb{Z}/2 \neq 0$$

Definition 2.24: Projective R -modules

An R -module P is *projective* if for every epimorphism $\varepsilon: M \rightarrow N$ of R -modules, the map

$$\text{Hom}(P, \varepsilon): \text{Hom}(P, M) \rightarrow \text{Hom}(P, N)$$

is surjective.

$$\begin{array}{ccc} & & M \\ & \nearrow & \downarrow \varepsilon \\ P & \xrightarrow{f} & N \end{array}$$

Fact. P is projective iff P is a direct summand of a free module iff there exists a R -module Q and a set S , such that

$$P \oplus Q \cong R[S].$$

Proof. Free modules are projective:

$$\begin{array}{ccc} & & M \\ & \nearrow g & \downarrow \varepsilon \\ R[S] & \xrightarrow{f} & N \end{array}$$

for every $s \in S$ choose $m_s \in M$ $\varepsilon(m_s) = f(s)$. Then there is a unique homomorphism $g: R[S] \rightarrow M$ such that $g(s) = m_s$.

Let P be projective and Q a summand of P . For reasons I couldn't copy, then Q is also projective.

Let P be a projective R -module. Consider the epimorphism

$$\begin{array}{c} R[P] \rightarrow P \\ p \mapsto p \end{array}$$

Then we have

$$\begin{array}{ccc} & & R[P] \\ & \nearrow g & \downarrow \\ p & \xrightarrow{\text{id}} & P \end{array}$$

So P is a direct summand of $R[P]$.

□

- If R is a field, then all modules are free, hence projective.
- $R = \mathbb{Z}/6$, $P = \mathbb{Z}/2$, $Q = \mathbb{Z}/3$. Then $\mathbb{Z}/6 \cong \mathbb{Z}/2 \oplus \mathbb{Z}/3$, so, as $\mathbb{Z}/6$ is free, $\mathbb{Z}/2$ and $\mathbb{Z}/3$ are projective, but not free.

Proposition 2.25. Let R be a commutative ring, and

$$0 \rightarrow I \xrightarrow{\alpha} M \xrightarrow{\beta} N \rightarrow 0$$

be a short exact sequence of R -modules.

Then for every R -module P , the sequence

$$P \otimes_R I \xrightarrow{P \otimes_R \alpha} P \otimes_R M \xrightarrow{P \otimes_R \beta} P \otimes_R N \rightarrow 0$$

is exact. („ $P \otimes_R _$ is right exact“). If moreover P is projective, then it is also exact with a 0 on the left, i.e. $P \otimes_R \alpha$ is injective. („projective modules are flat“).

Proof.

$$p \otimes_R \beta) \circ (p \otimes_R \alpha) = P \otimes_r (\beta \circ \alpha) = P \otimes_R 0 = 0$$

so $\text{Im}(P \otimes_R \alpha) \subseteq \ker(P \otimes_R \beta)$ so we get an induced homomorphism

$$\gamma \frac{P \otimes_R M}{\text{Im}(P \otimes_R \alpha)} \rightarrow P \otimes_R N$$

exactness is equivalent to δ being an isomorphism. We define a homomorphism $\delta: P \otimes_R N \rightarrow \frac{P \otimes_R M}{\text{Im}(P \otimes_R \alpha)}$ given by $(p, n) \in P \otimes N$ choose $\tilde{n} \in M$, such that $\beta(\tilde{n}) = n$.

Claim. $\delta(p \otimes n) = p \otimes \tilde{n} + \text{Im}(P \otimes_R \alpha)$ is independent of choice of \tilde{n}

Proof. Let $\tilde{\tilde{n}} \in M$ also satisfy $\beta(\tilde{\tilde{n}}) = n$. Then $\beta(\tilde{\tilde{n}} - \tilde{n}) = 0$, so there is $i \in I$ s.t. $\alpha(i) = \tilde{\tilde{n}} - \tilde{n}$. $p \otimes \tilde{\tilde{n}} - p \otimes \tilde{n} = p \otimes (\tilde{\tilde{n}} - \tilde{n}) = p \otimes \alpha(i) \in \text{Im}(P \otimes_R \alpha)$. \square

Claim. The assignment of δ is biadditive and sends (rp, n) and (p, rn) to the same element.

Then this extends to a well defined R -linear map

$$P \otimes_R N \rightarrow \frac{P \otimes_R M}{\text{Im}(P \otimes_R \alpha)}$$

which is isomorphic.

Now let P be projective. We show that then $P \otimes_R \alpha$ is injective.

Case 1 $P = R[S]$ free, S some set. Then

$$P \otimes_R M = R[S] \otimes_R M \cong \bigoplus_{s \in S} s \in SM$$

we have a natural isomorphism of R -modules in M .

From this we get a commutative square of R -modules:

$$\begin{array}{ccc} P \otimes_R I & \xrightarrow{P \otimes_R \alpha} & P \otimes_R M \\ \parallel & & \parallel \\ \bigoplus_{s \in S} I & \xrightarrow{\bigoplus_{s \in S} \alpha} & \bigoplus_{s \in S} M \end{array}$$

where the bottom map is injective.

General case P projective is a summand of a free module F , i.e. there are homomorphisms

$$P \xrightarrow{\lambda} F \xrightarrow{\mu} P$$

s.t. $\mu \circ \lambda = \text{Id}_P$. We consider the commutative square

$$\begin{array}{ccc} P \otimes_R I & \xrightarrow{P \otimes_R \alpha} & P \otimes_R N \\ \downarrow \lambda \otimes_R I & & \downarrow \lambda \otimes_R N \\ F \otimes_R I & \xrightarrow{F \otimes_R \alpha} & F \otimes_R N \end{array}$$

where the bottom map is injective by Case 1 and $\lambda \otimes_R I$ is injective, as it admits a retraction.

□

Definition 2.26: Global dimension of rings

A commutative ring R has global dimension ≤ 1 if every submodule of a projective module is projective.

Example 2.27. Some rings with global dimension ≤ 1 are

- fields
- the ring of integers \mathbb{Z} (subgroups of free abelian groups are free).
- every PID⁴ is of this form. See for example $k[x]$ for k a field or $\mathbb{Z}[i]$ the gaussian integers
- \mathbb{Z}_p the p -adic integers.

Definition 2.28: Tor of nice rings

Let R be a commutative ring of global dimension ≤ 1 . Let M, N be R -modules. Choose an epimorphism $p: P \rightarrow N$ of R -modules with P projective. Define

$$\text{Tor}^R(M, N) = \text{Ker}(M \otimes_R N \xrightarrow{M \otimes_R \text{incl}} M \otimes_R P)$$

Facts. This is independent up to preferred isomorphism of the choice of $p: P \rightarrow N$.

It is symmetric, i.e. we can resolve M instead of N .

If P is projective, then $\text{Tor}^R(P, N) = 0 = \text{Tor}^R(M, P)$.

Construction 2.29. For R a commutative ring, C, D complexes of R -modules. We define a natural homomorphism

$$\Phi: H_p(C) \otimes_R H_q(D) \rightarrow H_{p+q}(C \otimes_R D)$$

via $[x] \otimes [y] \mapsto [x \otimes y]$

We can check this is well defined.

⁴no zero divisors and every ideal is generated by a single element.

Theorem 2.30: Algebraic Künneth theorem

Let R be a commutative ring of global dimension ≤ 1 . Let C, D be complexes of projective R -modules. Then the following map is R -linearly split injective

$$\bigoplus \Phi: \bigoplus_{p+q=n} H_p(C) \otimes_R H_q(D) \rightarrow H_n(C \otimes_R D)$$

Moreover the cokernel is naturally isomorphic to

$$\bigoplus_{p+q=n-1} \mathrm{Tor}^R(H_p(C), H_q(D)).$$

Equivalently, there is a natural and split short exact sequence

$$0 \rightarrow \bigoplus_{p+q=n} H_p(C) \otimes_R H_q(D) \xrightarrow{\Phi} H_n(C \otimes_R D) \rightarrow \bigoplus_{p+q=n-1} \mathrm{Tor}^R(H_p(C), H_q(D)) \rightarrow 0$$

Proof. We let $Z = \{Z_q\}_{q \in \mathbb{Z}}$ be the complex of R modules with $d = 0$ where $Z_q = \mathrm{Ker}(d: D_q \rightarrow D_{q-1})$, let $B = \{B_q\}$ be the complex with $d = 0$ where $B_q = \mathrm{Im}(D: D_{q+1} \rightarrow D_q)$. We have a short exact sequence of complexes of R -modules

$$0 \rightarrow Z \xrightarrow{\mathrm{incl}} D \xrightarrow{d} B[1] \rightarrow 0$$

where $B[1]$ is the complex B shifted up by 1.

We have $B_q \subseteq Z_q \subseteq D_q$ projective by hypothesis. Since R has global dimension ≤ 1 , B_q and Z_q are also projective.

$$0 \rightarrow Z_q \rightarrow D_q \xrightarrow{d} B_{q-1} \rightarrow 0$$

is short exact, B_{q-1} is projective, so the sequence splits.

For every R -module N , the sequence

$$0 \rightarrow N \otimes_R Z_p \rightarrow N \otimes_R D_p \rightarrow N \otimes_R B_{p-1} \rightarrow 0$$

is exact.

This means we get a short exact sequence of complexes

$$0 \rightarrow C \otimes_R Z \rightarrow C \otimes_R D \rightarrow C \otimes_R B[1] \rightarrow 0$$

This means we get a long exact homology sequence

$$\rightarrow H_n(C \otimes_R Z) \xrightarrow{H_n(C \otimes_R \mathrm{incl})} H_n(C \otimes_R D) \xrightarrow{H_n(C \otimes_R d)} H_{n-1}(C \otimes_R B) \xrightarrow{\partial} H_{n-1}(C \otimes_R Z) \rightarrow \dots$$

Since Z has trivial differential:

$$H_n(C \otimes_R Z) = H_n\left(\bigoplus_{q \in \mathbb{Z}} C[q] \otimes Z_q\right) \cong \bigoplus_{q \in \mathbb{Z}} H_n(C[q] \otimes Z_q) \cong \bigoplus_{q \in \mathbb{Z}} H_n(C[q]) \otimes_R Z_q = \bigoplus_{p \in \mathbb{Z}} H_{n-q}(C) \otimes_R Z_q$$

where we use that Z_q is projective.

Similarly $H_n(C \otimes_R B) \cong \bigoplus_{q \in \mathbb{Z}} H_{n-q}(C) \otimes B_q$.

This gives us a long exact sequence

$$\dots \rightarrow \bigoplus_{p+q=n} H_p(C) \otimes_R Z_q \rightarrow H_n(C \otimes_R D) \rightarrow \bigoplus_{p+q=n-1} H_p(C) \otimes B_q \rightarrow \bigoplus_{p+q=n-1} H_p(C) \otimes Z_q$$

This splits up into short exact sequences

$$0 \rightarrow \bigoplus_{p+q=n} \text{Coker}(H_p(C) \otimes B_q \rightarrow H_p(C) \otimes_R Z_p) \rightarrow H_n(C \otimes_R D) \rightarrow \bigoplus_{p+q=n-1} \text{Ker}(H_p(C) \otimes_R B_q \rightarrow H_p(C) \otimes_R Z_q) \rightarrow 0$$

We know $0 \rightarrow B_q \rightarrow Z_q \rightarrow H_q(D)$ is a projective resolution of $H_q(D)$.

This means for all R -modules N ,

$$\text{Tor}^R(N; H_q(D)) = \text{Ker}(N \otimes_R B_q \rightarrow N \otimes_R Z_q)$$

$$N \otimes_R H_q(D) \cong \text{Coker}(N \otimes_R B_q \rightarrow N \otimes_R Z_q)$$

So we get:

$$0 \rightarrow \bigoplus_{p+q=n} H_p(C) \otimes H_q(D) \xrightarrow{\Phi} H_n(C \otimes_R D) \rightarrow \bigoplus_{p+q=n-1} \text{Tor}^R(H_p(C), H_q(D)) \rightarrow 0$$

for next lecture remains, that Φ has a R -linear retraction!

[23.04.2025, Lecture 5]
[28.04.2025, Lecture 6]

For the R -linear spitting.

Because B_q is projective, the following s.e.s. splits:

$$0 \rightarrow Z_q \xrightarrow{\text{incl}} D_q \xrightarrow{d} B_q \rightarrow 0$$

and the map Z_q to D_q admits a retraction. We choose a retraction $r_q: D_q \rightarrow Z_q$ to the inclusion.

Then

$$\begin{array}{ccccc} D_q + 1 & & & & \\ \downarrow d & \searrow 0 & & & \\ B_q & & & & \\ \downarrow \cap & \searrow 0 & & & \\ D_q & \xrightarrow{r_q} & Z_q & \longrightarrow & H_q(D) \end{array}$$

the retraction $\{r_q\}_{q \in \mathbb{Z}}$ for a morphism of chain complexes

$$r: D \rightarrow \{H_q(D), d = 0\}_q$$

that induces the identity on homology.

$H_q(r) \cong H_q(D) \rightarrow H_q(H_*(D), d = 0) = H_q(D)$. Similalry, there is a chain map $\rho: C \rightarrow \{H_p(C), d = 0\}$ that is the identity on homology. This gives a chain mpa $\rho \otimes_R r: C \otimes_R D \rightarrow (H_*(C) \otimes_R H_*(D), d = 0)$ which on homology

$$H_n(\rho \otimes_R r): H_n(C \otimes_R D) \rightarrow H_n(H_*(C) \otimes_R H_*(D), d = 0) = \bigoplus_{p+q=n} H_n(C) \otimes_R H_n(D)$$

which is a retraction to

$$\Psi: \bigoplus_{p+q=n} H_p(C) \otimes_R H_q(D) \rightarrow H_n(C \otimes_R D)$$

□

Example 2.31. Let R be a field. Then every module is free, hence projective, and

$$\mathrm{Tor}^R(M, N) = 0$$

for all R -modules M, N . For all complexes of R -modules C, D ;

$$\psi: \bigoplus_{p+q=n} H_p(C) \otimes_R H_q(D) \xrightarrow{\cong} H_n(C \otimes_R D).$$

is an isomorphism.

If $R = \mathbb{Z}$. Let C, D be a complex of free abelian groups. Then there is a split s.e.s.

$$0 \rightarrow \bigoplus_{p+q=n} H_p(C) \otimes H_q(D) \rightarrow H_n(C \otimes D) \rightarrow \bigoplus_{p+q=n-1} \mathrm{Tor}(H_p(C), H_q(D)) \rightarrow 0$$

2.5 Homological Künneth theorem

Construction 2.32 (Homology exterior pairing). Let X, Y be simplicial sets. Let R be a commutative ring. We define

$$\times: H_p(X, R) \otimes_R H_q(Y, R) \rightarrow H_{p+q}(X \times Y, R)$$

as the composite

$$H_p(C_*(X, R)) \otimes_R H_q(C_*(Y, R)) \xrightarrow{\Phi} H_{p+q}(C_*(X, R) \otimes C_*(Y, R)) \xrightarrow{H_{p+q}(\mathrm{EZ})} H_{p+q}(C_*(X \times Y, R))$$

For topological spaces A, B we Define

$$\times: H_p(A; R) \otimes_R H_q(B, R) \rightarrow H_{p+q}(A \times B, R)$$

as the composite

$$H_p(\mathcal{S}(A), R) \otimes_R H_q(\mathcal{S}(B), R) \xrightarrow{\times} H_{p+q}(\mathcal{S}(A) \otimes \mathcal{S}(B), R) \cong H_{p+q}(\mathcal{S}(A \times B); R)$$

where the isomorphism is given by the fact, that simplicial complex commutes with products. The isomorphism is the canonical map

$$\mathcal{S}(A) \times \mathcal{S}(B) \xleftarrow{(\mathcal{S}(p_A), \mathcal{S}(p_B))} \mathcal{S}(A \times B)$$

Theorem 2.33: Künneth theorem for homology with field coefficients

Let R be a field. Let X, Y be simplicial sets or spaces. Then the homology external product

$$\times: \bigoplus_{p+q=n} H_p(X, R) \otimes_R H_q(X, R) \rightarrow H_n(X \times Y; R)$$

is an isomorphism.

Proof. Follows directly from algebraic Künneth + Eilenberg-Zilber □

Theorem 2.34: Künneth theorem for homology

Let X, Y be spaces or simplicial sets. Then there is a natural and split s.e.s.

$$0 \rightarrow \bigoplus_{p+q=n} H_p(X, \mathbb{Z}) \otimes H_q(Y, \mathbb{Z}) \rightarrow H_n(X \times Y; \mathbb{Z}) \rightarrow \bigoplus_{p+q=n-1} \text{Tor}(H_p(X, \mathbb{Z}), H_q(Y, \mathbb{Z})) \rightarrow 0$$

Special Case. Let X, Y be spaces or simplicial sets. Suppose that $H_n(X, \mathbb{Z})$ is free for all $n \geq 0$. Then

$$\bigoplus_{p+q=n} H_p(X, \mathbb{Z}) \otimes H_q(Y, \mathbb{Z}) \xrightarrow{\Phi} H_n(X \times Y; \mathbb{Z})$$

is an isomorphism.

2.6 Künneth theorem for cohomology

Next we want to show the Künneth theorem for cohomology. The strategy:

- EZ provides a chain homotopy equivalence $C_*(X, \mathbb{Z}) \otimes C_*(Y, \mathbb{Z})$ to $C_*(X \times Y, \mathbb{Z})$.
- $\text{Hom}(_, R): \mathbf{Chains} \rightarrow \mathbf{coChains}_R$ preserves chain homotopies, so

$$\text{Hom}(C_*(X, \mathbb{Z}), R) \otimes \text{Hom}(C_*(Y, \mathbb{Z}), R) \cong \text{Hom}((C_*(X \times Y), \mathbb{Z}), R)$$

- in favorable cases we can relate

$$H^*(\text{Hom}(C, R) \otimes_R \text{Hom}(D, R)) \text{ to } H^*(\text{Hom}(C, R)) \otimes_R H^*(\text{Hom}(D, R))$$

- apply the algebraic Künneth theorem.

Step 3 is the hard step. For that we study relations between Hom and tensors.

Let A be an abelian group and R a commutative ring. We make the set $\text{Hom}(A, R)$ of group homomorphisms into an R module by pointwise addition and skalar multiplication. So $f, g \in \text{Hom}(A, R)$, $r \in R$. then

$$(f + g)(a) = f(a) + g(a), \quad ((r \cdot f)(a) = r \cdot f(a))$$

Let B be another abelian group. Then

$$\bullet: \text{Hom}(A, R) \times \text{Hom}(B, R) \rightarrow \text{Hom}(A \otimes B, R)$$

by $(f \bullet g)(a \otimes b) = f(a) \cdot g(b)$. This is additive in f and g .

$$(f + f') \bullet g = (f \bullet g) + (f' \bullet g)$$

and

$$(rf) \bullet g = r \cdot (f \bullet g) = f \bullet (r \cdot g)$$

for all $r \in R$. This means this extends to a well-defined R -linear map

$$\text{Hom}(A, R) \otimes_R \text{Hom}(B, R) \rightarrow \text{Hom}(A \otimes B, R)$$

Proposition 2.35. Let A, B be abelian groups and R a commutative ring. If A is finitely generated and free, then

$$\text{Hom}(A, R) \otimes_R \text{Hom}(B, R) \rightarrow \text{Hom}(A \otimes B, R)$$

is an isomorphism of R -modules.

Proof. For $A = \mathbb{Z}$:

$$\begin{array}{ccc} \mathrm{Hom}(\mathbb{Z}, R) \otimes_R \mathrm{Hom}(B, R) & \xrightarrow{\bullet} & \mathrm{Hom}(\mathbb{Z} \otimes B, R) \\ \downarrow \mathrm{ev} \otimes_R \mathrm{Hom}(B, R) & & \downarrow \cong \mathrm{Hom}(k, R) \\ R \otimes_R \mathrm{Hom}(B, R) & \xrightarrow[r \otimes g \mapsto r \cdot g]{\cong} & \mathrm{Hom}(B, R) \end{array}$$

where we have $k: B \rightarrow \mathbb{Z} \otimes B$ with $b \mapsto 1 \otimes b$.

Suppose the claim holds for A and A' . Then it holds for $A \oplus A'$.

$$\begin{array}{ccc} \mathrm{Hom}(A \oplus A', R) \otimes_R \mathrm{Hom}(B, R) & \xrightarrow{\bullet} & \mathrm{Hom}((A \oplus A') \otimes B, R) \\ \parallel & & \parallel \\ (\mathrm{Hom}(A, R) \oplus \mathrm{Hom}(A', R)) \otimes_R \mathrm{Hom}(B, R) & & \mathrm{Hom}((A \otimes B) \oplus (A' \otimes B), R) \\ \parallel & & \parallel \\ (\mathrm{Hom}(A, R) \otimes_R \mathrm{Hom}(B, R)) \oplus (\mathrm{Hom}(A', R) \otimes_R \mathrm{Hom}(B, R)) & \xrightarrow[\text{by assumption}]{\cong} & \mathrm{Hom}(A \otimes B, R) \oplus \mathrm{Hom}(A' \otimes B, R) \end{array}$$

The claim holds for $A = \mathbb{Z}^k$, $k \in \mathbb{N}$. any finitely generated free abelian group is isomorphic to \mathbb{Z}^k . \square

Example 2.36. $R = \mathbb{F}_2$ $A = B = \mathbb{Z}[\mathbb{N}]$. Then $\mathrm{Hom}(\mathbb{Z}[\mathbb{N}], R) \cong \mathrm{maps}(\mathbb{N}, R)$ by evaluation of generators. This is R -linear by the R -module structure on $\mathrm{maps}(\mathbb{N}, R)$.

$$\begin{array}{ccc} \mathrm{Hom}(A, R) \otimes \mathrm{Hom}(B, R) & \xrightarrow{\bullet} & \mathrm{Hom}(A \otimes B, R) \\ \mathrm{maps}(\mathbb{N}, R) \otimes_R \mathrm{maps}(\mathbb{N}, R) & & \mathrm{Hom}(\mathbb{Z}[\mathbb{N} \times \mathbb{N}], R) \\ & & \mathrm{maps}(\mathbb{N} \times \mathbb{N}, R) \end{array}$$

This is however not an isomorphism.

$A = B = \mathbb{Z}/2$ and $R = \mathbb{Z}/4$. Then $\mathrm{Hom}(A, R) = \mathrm{Hom}(B, R) = \mathrm{Hom}(\mathbb{Z}/2, \mathbb{Z}/4)$ is cyclic of order two generate by $i: \mathbb{Z}/2 \rightarrow \mathbb{Z}/4$, $n + 2\mathbb{Z} \mapsto 2n + 4\mathbb{Z}$.

$$\begin{array}{ccc} \mathrm{Hom}(A, R) \otimes_R \mathrm{Hom}(B, R) & \xrightarrow{\bullet} & \mathrm{Hom}(A \otimes B, R) \\ \parallel & & \\ \mathrm{Hom}(\mathbb{Z}/2, \mathbb{Z}/4) \otimes_{\mathbb{Z}/4} \mathrm{Hom}(\mathbb{Z}/2, \mathbb{Z}/4) & & \mathrm{Hom}(\mathbb{Z}/2 \otimes \mathbb{Z}/2, \mathbb{Z}/4) \\ \parallel & & \parallel \\ \mathbb{Z}/2 & \xrightarrow{0} & \mathbb{Z}/2 \end{array}$$

This shows, that both assumptions are strictly necessary.

Now let C, D be complexes of abelian groups. Then $\mathrm{Hom}(C, R), \mathrm{Hom}(D, R)$ are cochain complexes of R -modules.

$$\mathrm{Hom}(C, R)^n = \mathrm{Hom}(C_n, R)$$

and

$$d^n: \mathrm{Hom}(C, R)^n \rightarrow \mathrm{Hom}(C, R^{n+1}) = \mathrm{Hom}(D_{n+1}, R)$$

The sum of the \oplus homomorphism gives a cochain map

$$\bigoplus: \operatorname{Hom}(C, R) \otimes_R \operatorname{Hom}(D, R) \rightarrow \operatorname{Hom}(C \otimes D, R)$$

which is in dimension n :

$$\bigoplus_{p+q=n} \operatorname{Hom}(C_p, R) \otimes_R \operatorname{Hom}(D_q, R) \xrightarrow{\text{sum of } \bigoplus} \operatorname{Hom}\left(\bigoplus_{p+q=n} C_p \otimes D_q, R\right)$$

Proposition 2.37. Let C and D be chain complexes of abelian groups, such that $C_n = 0 = D_n$ for $n < 0$ and that C_n is finitely generated and free for all $n \geq 0$. Then \bigoplus is an isomorphism.

$$\bigoplus: \operatorname{Hom}(C, R) \otimes \operatorname{Hom}(D, R) \rightarrow \operatorname{Hom}(C \otimes D, R)$$

is an isomorphism of cochain complexes.

Proof. The vanishing hypothesis makes the potentially infinite sums

$$\bigoplus_{p+q=n} \operatorname{Hom}(C_p, R) \otimes_R \operatorname{Hom}(D_q, R)$$

finite.

Then $\operatorname{Hom}(_, R)$ preserves sums. And

$$\operatorname{Hom}(C_p, R) \otimes \operatorname{Hom}(D_q, R) \xrightarrow{\bigoplus} \operatorname{Hom}(C_p \otimes D_q, R)$$

is an isomorphism by the previous proposition. \square

This is not yet good enough to apply to topological spaces, as they are very not finitely generated.

Proposition 2.38. Let C be a chain complex of free abelian groups, such that $C_n = 0$ for $n < 0$. Suppose that $H_n(C)$ is finitely generated for all $n > 0$.

Then there is a subcomplex B of C , such that

- B_n is finitely generated and free for all $n \geq 0$.
- The inclusion $B \rightarrow C$ is a chain homotopy equivalence.

Proof. We construct subgroups B_n of C_n by induction on $n \geq 0$, such that

- $d(B_n) \subseteq B_{n-1}$
- the inclusions of $0 \rightarrow B_n \xrightarrow{d} B_{n-1} \xrightarrow{d} \cdots \rightarrow B_0 \rightarrow 0$
- into C induces an isomorphism on H_i for all $0 \leq i \leq n-1$ and an epimorphism on H_n .

Induction start: Let x_1, \dots, x_m be elements of C_0 , that generate $H_0(C)$. Select B_0 to be the subgroups of C_0 generated by x_1, \dots, x_m .

Induction step: Suppose B_0, \dots, B_{n-1} have been constructed fulfilling the conditions. Let x_1, \dots, x_m be cycles in C_n whose homology classes generate $H_n(C)$, which is possible because $H_n(C)$ is finitely generated. Set

$$Z = \operatorname{Ker}(d: B_{n-1} \rightarrow B_{n-2}) \cap \operatorname{Im}(d: C_n \rightarrow C_{n-1})$$

which is finitely generated because B_{n-1} is. Let z_1, \dots, z_k generate this intersection. Choose $y_1, \dots, y_k \in C_n$, such that $d(y_i) = z_i$ for $1 \leq i \leq k$.

Let B_n be the subgroup generated by $x_1, \dots, x_m, y_1, \dots, y_k$. Then $d(B_n) \subseteq B_{n-1}$.

Let $B_{\leq n}$ and $B_{< n}$ be the subcomplexes of C generated by B_0, \dots, B_n and B_0, \dots, B_{n-1}

Then $B_{< n} \subseteq B_{\leq n} \subseteq C$ where $B_{< n}$ induces isomorphism on H_i for $0 \leq i \leq n-2$ and epi on H_{n-1} . Similarly $B_{< n} \rightarrow B_{\leq n}$ is iso in dimension $\leq n-1$.

Then $B_{\leq n}$ is an Isomorphism on H_i for $0 \leq i \leq n-2$ and surjective on H_n because we include x_1, \dots, x_m that generate $H_n(C)$.

Let $x \in B_{n-1}$ be any cycle whose class is in the kernel of $H_{n-1}(B_{< n}) \rightarrow H_{n-1}(C)$. Then $x \in Z$ so x is a linear combination of the classes z_1, \dots, z_k and hence a boundary of a linear combination of y_1, \dots, y_k . So $x = d(w)$ for some $w \in B_n$. Then

$$\begin{array}{ccc} & H_{n-1}(B_n) & \\ \nearrow & & \searrow \\ H_{n-1}(B_{< n}) & \xrightarrow{\quad\quad\quad} & H_{n-1}(C) \end{array}$$

the class of x maps to 0 and the map becomes injective and hence an isomorphism.

We let B be the subcomplex of C generated by all B_i for all $i \geq 0$. Then the inclusion $B \rightarrow C$ induces an isomorphism on H_i for all $i \geq 0$, so it is a quasi-isomorphism.

By the end of last term we proved, it is already a chain homotopy equivalence! \square

[23.04.2025, Lecture 6]
[30.04.2025, Lecture 7]

Theorem 2.39: Algebraic Künneth theorem, cohomology

Let R be a commutative ring of global dimension ≤ 1 . Let C, D be chain complexes of abelian groups such that $C_n = 0 = D_n$ for $n < 0$ and all C_n are free and $H_n(C)$ is finitely generated free.

Then for all $n \geq 0$:

$$\bigoplus_{p+q=n} H^p(\text{Hom}(C, R)) \otimes_R H^q(\text{Hom}(D, R)) \xrightarrow{\Phi} H^n(\text{Hom}(C \otimes D, R))$$

is injective and its cokernel is isomorphic to

$$\bigoplus_{p+q=n+1} \text{Tor}^R(H^p(\text{hom}(C, R)), H^q(\text{Hom}(D, R)))$$

Warning. We do not assume, that there is a splitting.

Proof. „Basically just putting all the hard stuff we’ve already done together in the right way.“

Case 1 Suppose that also C_n is finitely generated for all $n \geq 0$. Then $\bullet: \text{Hom}(C, R) \otimes_R \text{Hom}(D, R) \rightarrow \text{Hom}(C \otimes D, R)$ is an isomorphism of cochain complexes. Applying the homological algebraic Künneth theorem to

$$H^n(\text{Hom}(C \otimes D, R)) \cong H^n(\text{Hom}(C, R) \otimes_R \text{Hom}(D, R))$$

since C_n is finitely generated and free, it is isomorphic to \mathbb{Z}^k for some $k \geq 0$, so $\text{Hom}(C, R)^n = \text{Hom}(C_n, R) \cong \text{Hom}(\mathbb{Z}^k, R) = R^k$ which is free hence projective as an R -module for all $n \geq 0$.

Caveat 1. we make cochain complexes into chain complexes, then apply Künneth, then come back. This turns $n-1$ in the \bigoplus for $\text{Tor} R$ into $n+1$.

Caveat 2. The proof of the homological Künneth theorem (without the splitting) used only that one complex is dimensionwise projective. Hence it is no problem, that D is not projective.

General case We choose a subcomplex B of C such that B_n is finitely generated for all $n \geq 0$ and $B \hookrightarrow C$ is a chain homotopy equivalence. Then

$$\mathrm{Hom}(i, R): \mathrm{Hom}(B, R) \rightarrow \mathrm{Hom}(C, R)$$

is a chain homotopy equivalence of R -module complexes.⁵

Note Additive functors preserve chain homotopy equivalences, however not quasi-isomorphisms. Because of that, quasi-Isomorphisms and chain homotopy equivalences are quite different.

Similarly we see

$$\mathrm{Hom}(i \otimes D, R): \mathrm{Hom}(C \otimes D, R) \rightarrow \mathrm{Hom}(B \otimes D, R)$$

is a chain homotopy equivalence.

This gives a commutative square in $\mathbf{coChains}_R$:

$$\begin{array}{ccc} \bigoplus_{p+q=n} H^p(\mathrm{Hom}(C, R)) \otimes H^q(\mathrm{Hom}(D, R)) & \xrightarrow{\Phi} & H^n(\mathrm{Hom}(C \otimes D, R)) \\ \downarrow \cong & & \downarrow \cong \\ \bigoplus_{p+q=n} H^p(\mathrm{Hom}(B, R)) \otimes_R H^q(\mathrm{Hom}(D, R)) & \xrightarrow[\text{by special case}]{\Phi} & H^n(\mathrm{Hom}(B \otimes D, R)) \end{array}$$

□

Construction 2.40. Let X, Y be spaces or simplicial sets. R an commutative ring. The *exterior cup product*

$$\times: H^p(X, R) \times H^q(Y, R) \rightarrow H^{p+q}(X \otimes Y, R)$$

is defined by $(x, y) \mapsto p_1^*(x) \cup p_2^*(y)$, where $p_1: X \times Y \rightarrow X$ and $p_2: X \times Y \rightarrow Y$.

Recall. The AW-map is

$$\mathrm{AW}: C_*(X \times Y) \rightarrow C_*(X) \otimes C_*(Y)$$

Proposition 2.41. Let X, Y be simplicial sets, R commutative ring. Then the composite

$$H^p(X, R) \otimes_R H^q(Y, R) \xrightarrow[\lfloor f]{\otimes} [g] \mapsto [f \otimes g] \Phi H^{p+q}(\mathrm{Hom}(C_*(X), R) \otimes_R \mathrm{Hom}(C_*(Y), R)) \xrightarrow{H^{p+q}(\bullet)} H^{p+q}(\mathrm{Hom}(C_*(X \otimes Y), R))$$

equals the external cup product.

Proof. In the notes. □

Theorem 2.42: Künneth theorem in cohomology

Let R be a commutative ring of global dimension ≤ 1 . Let X, Y be spaces such that $H_n(X, \mathbb{Z})$ is finitely generated for all $n \geq 0$. Then the total exterior cup product map

$$\bigoplus_{p+q=n} H^p(X, R) \otimes_R H^q(Y, R) \rightarrow H^n(X \times Y, R)$$

is injective, and its cokernel is naturally isomorphic to

$$\bigoplus_{p+q=n+1} \mathrm{Tor}^R(H^p(X, R), H^q(Y, R))$$

⁵This is due to the Hom-functor being additive. Unfortunately I don't know what that means.

Proof. Similar to the homological one. Use the cohomological algebraic Künneth theorem and the Eilenberg-Zilber theorem. You can read it up somewhere. \square

Remark 2.43. Let X be a CW-complex of finite type i.e. such that it has only finitely many cells in every dimension. (ex. $\mathbb{R}P^\infty$). Then

$$C_*^{Cell}(X, \mathbb{Z})$$

is finitely generated free in every dimension, hence $H_n^{cell}(X, \mathbb{Z}) \cong H_n(X, \mathbb{Z})$ if finitely generated, so Künneth theorem applies.

Construction 2.44. Let A, B be graded-commutative⁶ rings. Then $A \otimes B$ is another graded-commutative ring by

$$(A \otimes B)_n = \bigoplus_{p+q=n} A_p \otimes B_q$$

with multiplication for $a \in A_p, b \in B_q, a' \in A_{p'}, b' \in B_{q'}$.

$$(a \otimes b) \cdot (a' \otimes b') = (-1)^{p' \cdot q} (aa') \otimes (bb')$$

Check for well-definedness yourself.

Corollary 2.45. Let R be a field, X, Y spaces and suppose, that $H_n(X, \mathbb{Z})$ is finitely generated for all $n \geq 0$. Then

$$\times : H^*(X, R) \otimes_R H^*(Y, R) \rightarrow H^*(X \times Y, R)$$

is an isomorphism of graded-commutative R -algebras.

Note. We already knew that this is a isomorphism of abelian groups. The new information is, that this is compatible with ring structure.

Proof. We take $x \in H^p(X, R), x' \in H^{p'}(X, R), y \in H^q(Y, R), y' \in H^{q'}(Y, R)$ and then

$$\begin{aligned} (x \cup x') \times (y \cup y') &= p_1^*(x \cup x') \cup p_2^*(y \cup y') \\ &= (p_1^*(x) \cup p_1^*(x')) \cup (p_2^*(y) \cup p_2^*(y')) \\ &= (-1)^{p' \cdot q} (p_1^*(x) \cup p_2^*(y)) \cup (p_1^*(x') \cup p_2^*(y')) \\ &= (-1)^{p \cdot q'} (x \times y) \cup (x' \times y') \end{aligned}$$

\square

Corollary 2.46. Let X, Y be spaces such that $H_n(X, \mathbb{Z})$ is finitely generated and free for all $n \geq 0$. Then

$$H^*(X, \mathbb{Z}) \otimes H^*(Y, \mathbb{Z}) \rightarrow H^*(X \times Y, \mathbb{Z})$$

is an isomorphism of graded-commutative rings.

2.7 Example calculations for Cohomology-rings

Now we are actually calculating some cohomology rings. Namely for $S^k \times S^l, S^1 \times \cdots \times S^1$ and $\mathbb{C}P^2$.

Remember

$$H^n(S^k) = \begin{cases} \mathbb{Z} & n = 0, k \\ 0 & n \neq 0, k \end{cases}$$

⁶ $a \cdot b = (-1)^{\deg(A) \cdot \deg(B)} b \cdot a$

and assume $k \geq 1$. For dimensional reasons, the cup product on $H^*(S^k, \mathbb{Z})$ is trivial. $H^*(S^k, \mathbb{Z})$ is dimensionwise finitely generated free, and hence for every space Y the exterior cup product

$$H^*(S^k, \mathbb{Z}) \otimes H^*(Y, \mathbb{Z}) \rightarrow H^*(S^k \times Y, \mathbb{Z})$$

is an isomorphism of graded-commutative rings. Take $Y = S^l$ for $l \geq 1$.

Let $e_k \in H^n(S^k; \mathbb{Z})$ be one of the two generators. Then $H^+(S^k, \mathbb{Z}) = \Lambda(e_k)$ where Λ denotes an exterior product. This includes $e_k^2 = 0$. We define $a := p_1^*(e_k) \in H^k(S^k \times S^l; \mathbb{Z})$ and $b := p_2^*(e_l) \in H^l(S^k \times S^l; \mathbb{Z})$. Then

$$H^*(S^k, \mathbb{Z}) \otimes H^*(S^l, \mathbb{Z}) \xrightarrow[x]{\cong} H^*(S^k \times S^l; \mathbb{Z}) = \mathbb{Z}\{1 \times 1, 1 \times e_l, e_k \times 1, e_k \cdot e_l\}$$

where we have $1 \times 1 = 1, 1 \times e_l = b, e_k \times 1 = a, e_k \cdot e_l = a \cup b$.

We look at multiplicative relations:

$$a^2 = 0, b^2 = 0$$

and so

$$a^2 = (p_1^*(e_k))^2 = p_1^*(e_k^2) = p_1^*(0) = 0$$

If k or l is even, then $a \cup b = b \cup a$ and if both are odd, then $a \cup b = -b \cup a$.

We summarize, if k and l are even, then

$$H^*(S^k \times S^l; \mathbb{Z}) = \mathbb{Z}[a, b]/(a^2 = 0, b^2 = 0)$$

and if one is odd

$$H^*(S^k \times S^l; \mathbb{Z}) = \Lambda(a, b)$$

where Λ again denotes exterior products.

We give an inductive description of $H^*(S^1 \times \cdots \times S^1; \mathbb{Z})$ n -times. We use, that

$$\times: H^*(S^1; \mathbb{Z}) \otimes H^*(\underbrace{S^1 \times \cdots \times S^1}_{n-1 \text{ times}}) \cong H^*(S^1 \times \cdots \times S^1, \mathbb{Z})$$

we define $a_i = p_i^*(e_1) \in H^1(\underbrace{S^1 \times \cdots \times S^1}_n; \mathbb{Z})$, where $p_i: (S^1)^n \rightarrow S^1$ is projection to the i -th factor for $1 \leq i \leq n$. We get $a_i^2 = 0$ and $a_i \cup a_j = -a_j \cup a_i$ for $i \neq j$. This gives us, that an additive basis of $H^*(S^1)^n; \mathbb{Z}$ is given by

$$a_{i_1} \cup \cdots \cup a_{i_k} \text{ for all tuples } 1 \leq a_i < a_2 < \cdots < a_k \leq n$$

This gives us $\text{rank}(H^*((S^1)^n; \mathbb{Z})) = 2^n$. The multiplicative structure is given by $H^*((S^1)^n, \mathbb{Z}) = \Lambda(a_1, \dots, a_n)$.

Later we will compute $H^*(\mathbb{C}P^n; \mathbb{Z})$ via Poincaré-duality to get $\cong \mathbb{Z}[X]/(X^{n+1})$ for $x \in H^2(\mathbb{C}P^n, 2)$.

We will now use a trick to at least calculate $H^*(\mathbb{C}P^2; \mathbb{Z})$. We know, that

$$H^n(\mathbb{C}P^2, \mathbb{Z}) = \begin{cases} \mathbb{Z} & n = 0, 2, 4 \\ 0 & \text{else} \end{cases}$$

we take $x \in H^2(\mathbb{C}P^2; \mathbb{Z})$ a generator. The multiplicative structure is completely defined by which multiple of the generator of $H^4(\mathbb{C}P^2, \mathbb{Z})$ x^2 is.

We use homogenous coordinate notation for $\mathbb{C}P^2$. For $0 \neq (x, y, z) \in \mathbb{C}^3$ we write $[x, y, z] := \mathbb{C} \cdot (x, y, z) \in \mathbb{C}P^2$. We define a continuous map

$$\mu: \mathbb{C}P^1 \times \mathbb{C}P^1 \rightarrow \mathbb{C}P^2$$

given by $([v, w], [x, y]) \mapsto [vx, vy + wx, wy]$. We let $e = [1, 0]$ a basepoint in $\mathbb{C}P^1$. Then $\mu(e, _), \mu(_, e): \mathbb{C}P^1 \rightarrow \mathbb{C}P^2$. are both the „standard inclusions“ $[x, y] \mapsto [x, y, 0]$.

Proposition 2.47. The map $\mu^*: H^4(\mathbb{C}P^2, \mathbb{Z}) \rightarrow H^4(\mathbb{C}P^1 \times \mathbb{C}P^1, \mathbb{Z})$ is injective and its image has index 2.

proof next time.

[30.04.2025, Lecture 7]

[5.05.2025, Lecture 8] Rather sleepy today, quality may be accordingly.

Note. Remember $\mathbb{C}P^2 \cong S^2$.

Proof. We will drop coefficients from the notation. $H^*(X) := H^*(X; \mathbb{Z})$. The continuous map $\mathbb{C}^2 \rightarrow \mathbb{C}P^2$, $\pi(a, b) = (a^2 - b, 2a, 1)$ is an open embedding and a homoeomorphism onto the open 4-cell $\mathbb{C}P^2 \setminus \mathbb{C}P^1$. That is just the set $[x, y, 1]$ for $(x, y) \in \mathbb{C}^2$. Then

$$(x, y) = (a^2 - b, 2a) \implies a = y/2, b = (a - x = y^2/4 - x)$$

This gives an isomorphism of relative cohomology groups

$$\pi^*: H^4(\mathbb{C}P^2 \setminus \mathbb{C}P^1, \mathbb{C}P^2 \setminus (\mathbb{C}P^1 \cup [0, 0, 1])) \rightarrow H^4(\mathbb{C}^2, \mathbb{C}^2 \setminus (0, 0))$$

Then we have an excision isomorphism:

$$H^4(\mathbb{C}P^2, \mathbb{C}P^2 \setminus [0, 0, 1]) \cong H^4(\mathbb{C}P^2 \setminus \mathbb{C}P^1, \mathbb{C}P^2 \setminus \mathbb{C}P^1 \cup [0, 0, 1])$$

The long exact sequence of the pair gives an isomorphism

$$H^4(\mathbb{C}P^2, \mathbb{C}P^2 \setminus [0, 0, 1]) \rightarrow H^4(\mathbb{C}P^2)$$

We also Define

$$\pi': \mathbb{C}^2 \rightarrow \mathbb{C}P^1 \times \mathbb{C}P^1, (a, b) \mapsto ([a + b, 1], [a - b, 1])$$

A similar calculation gives

$$H^4(\mathbb{C}P^1 \times \mathbb{C}P^1, \mathbb{C}P^1 \times \mathbb{C}P^1 \setminus ([0, 1], [0, 1]))$$

as isomorphic to $H^4(\mathbb{C}P^1 \times \mathbb{C}P^1)$.

We now also define ν .

$$\nu: \mathbb{C}^2 \rightarrow \mathbb{C}^2; \quad (a, b) \mapsto (a, b^2)$$

Now a diagram I didn't copy commutes.

The problem now reduces to show that

$$\nu^*: H^4(\mathbb{C}^2, \mathbb{C}^2 \setminus (0, 0)) \rightarrow H^4(\mathbb{C}^2, \mathbb{C}^2 \times 0), 0$$

is multiplication by 2.

A diagram I didn't copy. He applied Künneth and found out some map is multiplication by 2. \square

Proposition 2.48. Let $x \in H^2(\mathbb{C}P^2, \mathbb{Z})$ be an additive generator. Then x^2 is an additive generator of $H^4(\mathbb{C}P^2, \mathbb{Z})$. So $H^*(\mathbb{C}P^2, \mathbb{Z})$ is a truncated polynomial algebra i.e.

$$H^*(\mathbb{C}P^2, \mathbb{Z}) = \mathbb{Z}[X]/(x^3)$$

Outlook. $H^*(\mathbb{C}P^m; \mathbb{Z}) = \mathbb{Z}[X]/(x^{m+1})$ This will be proven later using Poincaré-Duality.

Proof. We write $i: \mathbb{C}P^1 \rightarrow \mathbb{C}P^2$ for „the inclusion“, $i[x, y] = [x, y, 0]$. Then

$$H^*(\mathbb{C}P^1, \mathbb{Z}) = \mathbb{Z}\{1, i^2(x)\}$$

$$\times: H^*(\mathbb{C}P^2) \otimes H^*(\mathbb{C}P^1) \cong H^*(\mathbb{C}P^1 \times \mathbb{C}P^1)$$

we write $a := p_1^*(i^*(x)), b := p_2^*(i^*(x))$. Then

$$H^*(\mathbb{C}P^1 \times \mathbb{C}P^1) = \mathbb{Z}\{1, a, b, a \cdot b\}$$

with $a^2 = b^2 = 0, ab = ba$.

Claim. We have

$$\begin{array}{ccc} \mu^*(x) = a + b & \in & H^2(\mathbb{C}P^1 \times \mathbb{C}P^1) \\ & & \downarrow \cong \\ & & H^2(\mathbb{C}P^1 \vee \mathbb{C}P^1) \\ & & \downarrow \cong \\ & & H^2(\mathbb{C}P^1) \times H^2(\mathbb{C}P^1) \end{array}$$

where we use that the wedge is an isomorphism on the 2-skeleton of $\mathbb{C}P^1 \times \mathbb{C}P^1$. The composite map is given by

$$z \mapsto ((e, _)^*(z)), (_ , e)^*(z))$$

We note

$$\begin{aligned} (e, _)^*(a + b) &= (e, _)^*(p_1^*(i^*(x))) + (e, _)^*(p_2^*(i^*(x))) \\ &= \underbrace{(i \circ p_1 \circ (e, _))^*(x)}_{\text{constant}} + \underbrace{(i \circ p_2 \circ (e, _))^*(x)}_{\text{identity}} \\ &= i^*(x) \end{aligned}$$

and also

$$(e, _)^*(\mu^*(x)) = \underbrace{(\mu \circ (e, _))^*}_{=1} = i^*(x)$$

This gives $\mu^*(x) = a + b$. Now let $y \in H^4(\mathbb{C}P^2)$ be a generator and let $n \in \mathbb{Z}$ be such, that $x^2 = n \cdot y$. Now

$$2ab = (a + b)^2 = (\mu^*(x))^2 = \mu^*(x^2) = \mu^*(ny) = n \cdot \mu^*(y) = n \cdot 2 \cdot ab$$

where the last equality uses degree 2 of μ . This holds in the free abelian group $H^4(\mathbb{C}P^1 \times \mathbb{C}P^1) = \mathbb{Z}\{a, b\}$. This means $2 = 2n$ and hence $n = 1$ and so $x^2 = y$. \square

Application to the Hopf map.

The Hopf map $\eta: S^3 \rightarrow S^2$ is defined as

$$S^3 = S(\mathbb{C}^2) \rightarrow \mathbb{C}P^1 \cong S^2$$

given by $(x, y) \mapsto [x, y]$.

Then $0 \neq [y] \in \pi_3(S^2, *) \cong \mathbb{Z}\{y\}$.

Proposition 2.49. Attaching a 4-cell to $\mathbb{C}P^1$ yields a space homeomorphic to $\mathbb{C}P^2$. Informally: „ η is the attaching map of the 4-cell in $\mathbb{C}P^2$.“

Proof. Consider the map $\alpha: D(\mathbb{C}^2) \rightarrow \mathbb{C}P^2$, $(x, y) \mapsto [x, y, 1 - |x|^2 - |y|^2]$.

This restricts to a homeomorphism from $D(\mathbb{C}^4) \setminus S(\mathbb{C}^2)$ onto $\mathbb{C}P^2 \setminus \mathbb{C}P^1$ and the following commutes:

$$\begin{array}{ccc} S(\mathbb{C}^2) & \xrightarrow{\eta} & \mathbb{C}P^1 \\ \downarrow & & \downarrow i \\ D(\mathbb{C}^2) & \xrightarrow{\alpha} & \mathbb{C}P^2 \end{array} \quad \begin{array}{c} [x, y] \\ \downarrow \\ [x, y, 0] \end{array}$$

this gives a well-defined continuous map $D(\mathbb{C}^2) \cup_{S(\mathbb{C}^2), \eta} \mathbb{C}P^1 \rightarrow \mathbb{C}P^2$, This is a continuous bijection between compact Hausdorff spaces, hence a homeomorphism. \square

Theorem 2.50: Hopf map is not constant

The Hopf map η is not homotopic to a constant map.

Proof. By contradiction. If η was homotopic to the constant map $c: S^3 \rightarrow S^2$, then $D^4 \cup_{S^3, \eta} \mathbb{C}P^1$ would be homotopy-equivalent to $D^4 \cup_{S^3, \text{const}} \mathbb{C}P^1 = \mathbb{C}P^1 \vee (D^4/S^3) \cong S^2 \vee S^4$.

These spaces have the same additive cohomology. However, their cup-product differs. Namely in $H^*(\mathbb{C}P^1 \vee S^4, \mathbb{Z})$ the square of every 2-dimensional class is 0.

As such, $\mathbb{C}P^1 \vee S^4 \not\sim \mathbb{C}P^2$. \square

Outlook. The Hopf map is sometimes presented as the map

$$S(\mathbb{C}^2) \rightarrow \mathbb{C} \cup \{\infty\} = \text{one point compactification of } \mathbb{C} \cong S^2$$

given by $(x, y) \mapsto x/y$. For $\mathbb{H} =$ the quaternions $= \mathbb{R}^4$ with the skew-field multiplication $= \mathbb{R}\{1, i, j, k\}$ and $i^2 = j^2 = k^2 = ijk = -1$. And then we get

$$\nu: S^7 = S(\mathbb{H}^2) \mapsto \mathbb{H} \cup \{\infty\} = S^4$$

given by $(x, y) \mapsto x/y = xy^{-1} \vee y^{-1}x$. This map is also called the second Hopf-map. Using that most of linear algebra still applies to skew-fields, we can define $\mathbb{H}P^n$ and see by a similar argument, that ν is not nullhomotopic. Then $[\nu] \in \pi_7(S^4, *) \cong \mathbb{Z}\{\nu\} \oplus \mathbb{Z}/?$ Schwede doesn't remember what exactly π_7 is.

Then we also have $\mathbb{O} =$ Cayley octonians $= \mathbb{R}^8$ with a nonassociative, noncommutative division algebra structure $\cdot: \mathbb{O} \times \mathbb{O} \rightarrow \mathbb{O}$. Then there is still an $\mathbb{O}P^2$ but no general $\mathbb{O}P^n$.

However this is enough to still calculate that $H^*(\mathbb{O}P^2, \mathbb{Z}) = \mathbb{Z}[w]/w^3$ where $w \in H^8(\mathbb{O}P^2, \mathbb{Z})$. And you can show

$$\sigma: S(\mathbb{O}^2) \rightarrow \mathbb{O}P^1 = \mathbb{O} \cup \{\infty\}$$

given by $(x, y) \mapsto x/y$ is non zero-homotopic. And $[\sigma] \in \pi_{15}(S^8) = \mathbb{Z} \oplus \mathbb{Z}/120$.

He also talks about a theorem, that these are all the Hopf-Maps that exist. No more in higher dimensions.

[5.05.2025, Lecture 8]
[07.05.2025, Lecture 9]

Part II

Manifolds and Poincaré Duality

Chapter 3

Topological manifolds

The long-time goal is to prove Poincaré duality. For that we first need to study manifolds.

Definition 3.1: Manifold

An m -manifold is a Hausdorff space M such that every point of M has an open neighborhood homeomorphic to \mathbb{R}^m .¹

¹This is sometimes called a topological manifold to differentiate from smooth ones.

Remark 3.2. • The empty space is an m -manifold for all $m \geq 0$.

- Let M be a non empty manifold. Then the dimension m is an intrinsic invariant. Let $x \in M$ be a point, let U be an open neighborhood of x homeomorphic to \mathbb{R}^m . Let $\varphi: \mathbb{R}^m \rightarrow U$ be a homeomorphism such that $\varphi(0) = x$. Then

$$H_i(M, M \setminus \{x\}, \mathbb{Z}) \xleftarrow{\cong} H_i(U, U \setminus \{x\}, \mathbb{Z}) \xleftarrow{\varphi^*} H_i(\mathbb{R}^m, \mathbb{R}^m \setminus \{0\}, \mathbb{Z})$$

where we use excision for the first homeomorphism. Furthermore we see

$$H_i(\mathbb{R}^m, \mathbb{R}^m \setminus \{0\}, \mathbb{Z}) \sim H_i(D^m, S^{m-1}, \mathbb{Z}) = \begin{cases} \mathbb{Z} & i = m \\ 0 & i \neq m \end{cases}$$

We call this the local homology of x . From this we can reproduce the dimension of M .

- The Hausdorff condition is important to rule out pathological examples such as the „line with double origin“:

$$\mathbb{R} \amalg \mathbb{R} / (x, 0) \sim (x, 1) \text{ for all } x \in \mathbb{R} \setminus \{0\}$$

Can't draw the picture of the space.

This is not Hausdorff, but locally \mathbb{R}^1 . we don't want this to be a manifold.

Example 3.3. • open subsets of \mathbb{R}^m are m -manifolds.

- Let M be a Hausdorff space, such that every point has an open neighborhood that is an m -manifold. Then M is an m -manifold.
- Let M be an m -manifold and N an n -manifold. Then $M \times N$ is an $m + n$ -manifold.
- The m -sphere $S^m = \{(x_1, \dots, x_{m+1}) \in \mathbb{R}^m \mid x_1^2 + \dots + x_{m+1}^2 = 1\}$ is an m -manifold.

Let $x = (x_1, \dots, x_{m+1}) \in S^m$ be a point. Let $V = \{y \in \mathbb{R}^{m+1} \mid \langle y, x \rangle = 0\}$ be the orthogonal complement of x . The stereographic projection is a homeomorphism

$$x \in S^m \setminus \{-x\} \rightarrow V$$

given by some formula I couldn't copy before it was erased and he also had a nice picture.

- The real projective space $\mathbb{R}P^m \cong S^m / x \sim -x$ is an m -manifold. Let $\{x, -x\}$ be a point in $\mathbb{R}P^m$ for $x \in S^m$. Let x be one of the representatives. Let $\mathbb{R}^m \cong U = \{z \in S^m \mid \langle z, x \rangle \geq 0\}$

„The northern hemisphere with north-pole x “. As $U \subseteq S^m$ we get via projection a map to $\mathbb{R}P^m$. This is an open embedding onto a neighborhood.

- Let $\mathbb{C}P^m = \{l \in \mathbb{C}^{n+1} : L \text{ complex line through } 0\}$. is a $2m$ manifold. Consider first the point $[0, 0, \dots, 0, 1]$.

Then $\mathbb{R}^{2n} \cong \mathbb{C}^n \rightarrow \mathbb{C}P^n$ given by $(z_1, \dots, z_m) \mapsto [z_1, \dots, z_m, 1]$ is an homeomorphism onto a open neighborhood U of $[0, 0, \dots, 0, 1]$.

Let $l \in \mathbb{C}P^n$ be any point, let $v \in l$ be a nonzero vector in l . Let $A \in GL_n(\mathbb{C})$ such that $A \cdot (0, \dots, 0, 1) = v$. Then $A: \mathbb{C}P^n \rightarrow \mathbb{C}P^n$, $L_0 = [0, \dots, 0, 1]$ given by $L \mapsto A \cdot L$ sends $A \cdot L_0 = L$. So we can take $A(U)$ as an open neighborhood of L homeomorphic to \mathbb{R}^{2n} .

Now we do some examples that are a little more involved.

Construction 3.4 (Stiefel manifold). Let $0 \leq k \leq n$. The Stiefel manifold $V_{k,n} = \{(v_1, \dots, v_k) \in \mathbb{R}^n\} \mid \text{orthonormal set}\}$. We call this the „ k -frame“ this means

$$\langle v_i, v_j \rangle = \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases}$$

note, that each v_i is a vector in \mathbb{R}^n . We give $V_{k,n}$ the subspace topology of $(\mathbb{R}^n)^k$. This is even a closed subspace of $(S^{n-1})^k$, so $V_{k,n}$ is compact.

Examples are,

- $V_{0,n} = \{\emptyset\}$ is a point hence a 0-manifold.
- $V_{1,n} = S^{n-1}$.
- $V_{n,n} = O(n)$ the n -th orthogonal group.
- $V_{n-1,n} \xrightarrow{\cong} SO(n)$ given by $(Ae_1, \dots, A \cdot e_{n-1}) \mapsto A$ where $e_i = {}^t(0, \dots, 1, \dots, 0)$. This is bijective because it sends orthogonal matrices to the orthogonal vectors that span it. That is not what was written on the board. That was erased before I could copy.

Proposition 3.5. $V_{k,n}$ is a manifold of dimension $(n-1) + (n-2) + \dots + (n-k) = nk - \frac{k(k+1)}{2}$

Proof. By induction on k . We have already seen $V_{0,n} = \{\emptyset\}$ as a 0-manifold and $V_{1,n} = S^{n-1}$ a $(n-1)$ -manifold.

Now let $k \geq 2$. Let $S_+^{n-1} = \{(x_1, \dots, x_n) \in S^{n-1} : x_1 \geq 0\}$ be the „northern hemisphere“. We define a continuous map $\psi: S_+^{n-1} \rightarrow O(n)$ as the following composite

$$S_*^{n-1} \rightarrow GL_n(\mathbb{R}) \xrightarrow{\text{Gram-Schmidt}} O(n) \quad w \mapsto ({}_t w, e_2, \dots, e_n) \mapsto \dots$$

where Gram-Schmidt is a continuous way to orthonormalize a matrix.

We remember the properties:

- ψ is continuous
- $\psi(e_1) = \psi(1, 0, 0, \dots, 0) = E_n$
- $\psi(w) \cdot e_1 = w$.

Warning. There is no continuous map $\tilde{\psi}: S^{n-1} \rightarrow O(n)$ such that $\tilde{\psi}(w) \cdot e_1 = w$.

We show the manifold condition around $(e_1, \dots, e_k) \in V_{k,n}$. We set $U = \{(v_1, \dots, v_k) \in V_{k,n} : v_1 \in S_+^{n-1}\}$ is open in $V_{k,n}$ around (e_1, \dots, e_k) . The map

$$U \rightarrow S^{n-1} \times V_{k-1,n-1}, \quad (v_1, \dots, v_k) \mapsto (v_1, (\psi(v_1))^{-1}(v_2), \dots, (\psi(v_1))^{-1}(v_k))$$

where $(\psi(v_1))^{-1}(v_i)$ are in $0 \times \mathbb{R}^{n-1}$. The well-definedness follows from $\psi(v_1)^{-1}$ is an orthogonal matrix such that $\psi(v_1)^{-1}(v_1) = e_1$. This means, that $\psi(v_1)^{-1}(v_2, \dots, v_k)$ will be an orthonormal $k-1$ -set that is also orthogonal to e_1 , i.e. they sit in $0 \times \mathbb{R}^{n-1}$. He also rambles, as to why this is continuous.

It is a homeomorphism. This shows that around e_1, e_2, \dots, e_k $V_{k,n}$ is locally a manifold of dimension $(n-1) + \dim(V_{k-1,n-1}) = (n-1) + (n-2 + \dots + n-k)$.

We have a continuous inverse:

$$S_+^{n-1} \times V_{k-1,n-1} \rightarrow U \quad (v, w_1, \dots, w_{k-1}) \mapsto (v, \psi(v)(0, w_1), \dots, \psi(v)(0, w_{k-1}))$$

Now let $(v_1, \dots, v_k) \in V_{k,n}$ be any point. We choose an extension to an orthonormal basis $(v_1, \dots, v_k, v_{k+1}, \dots, v_n)$. Set $A = (v_1, \dots, v_n) \in O(n)$. then

$$A \cdot _ : V_{k,n} \rightarrow V_{k,n}$$

is a self homeomorphism that sends $(w_1, \dots, w_k) \mapsto (A \cdot w_1, \dots, A \cdot w_k)$ and specifically e_1, \dots, e_k to v_1, \dots, v_k . So the homeomorphism takes the previous neighborhood U homeomorphically onto the neighborhood $A \cdot U$ of (v_1, \dots, v_k) \square

Remark 3.6. What we really showed is, that $V_{k,n} \rightarrow S^{n-1}$, $(v_1, \dots, v_k) \mapsto v_1$ is a smooth locally trivial fiberbundle with fiber $V_{k-1,n-1}$.

Note. Complex Stiefel Manifold. We can also define

$$V_{k,n}^{\mathbb{C}} = \{(v_1, \dots, v_k) \in \mathbb{C}^n : \langle v_i, v_j \rangle = \delta_{i,j}\}$$

where δ denotes the Kronecker-symbol and we use the hermitian complex bilinear product.

This is a manifold of dimension $(2n-1) + (2n-3) + (2n-5) + \dots + (2n-2k+1) = 2nk - k^2$. We will see.

$$V_{0,n}^{\mathbb{C}} = \{\cdot\} \quad V_{1,n}^{\mathbb{C}} = S^{2n-1}, \quad V_{n-1,n} \cong SU(n), \quad V_{n,n} \cong U(n)$$

For the quaternions $\mathbb{H} = \mathbb{R}\{1, i, j, k\}$ with $i^2 = j^2 = k^2 = ijk = -1$, we have quaternionic conjugation $\lambda = a + bi + cj + dk \rightarrow \bar{\lambda} = a - bi - cj - dk$ that is an anti-isomorphism: $\lambda \bar{\cdot} \mu = \bar{\mu} \cdot \bar{\lambda}$. This gives a „Quaternionic skalar product“ on \mathbb{H}^n is defined by $[x, y] := \bar{x}_1 y_1 + \dots + \bar{x}_n y_n$ for $x, y \in \mathbb{H}^n$. This is an \mathbb{H} -sesquilinear, non degenerate positive definite \mathbb{R} -bilinear form.

With the right definitions and being careful, all of this works.

This gives Quaternionic Stiefel manifolds:

$$V_{k,n}^{\mathbb{H}} = \{(v_1, \dots, v_k) \in (\mathbb{H}^n)^k : [v_i, v_j] = \delta_{i,j}\}$$

is a manifold of dimension $(4n-1) + (4n-5) + \dots + (4n-4k+3) = 4nk - k(2k-1)$. And we see again

$$V_{1,n}^{\mathbb{H}} = S^{4n-1}, \quad V_{n,n}^{\mathbb{H}} = Sp(n) = \{A \in M(n \times n, \mathbb{H}) : A \cdot \bar{A}^t = \bar{A}^t \cdot A = E_n\}$$

Where Sp is the symplectic group. There is no such thing as a special symplectic group, because you would need determinant for that, which then really needs commutativity.

Construction 3.7 (Grassmann manifolds). Let $0 \leq k \leq n$ The Grassmann manifold of k -pairs in \mathbb{R}^n is

$$Gr(k, n) = Gr_k(n) = Gr_k(\mathbb{R}^n) = \{L \subseteq \mathbb{R}^n : L \text{ is } k\text{-dimensional } \mathbb{R}\text{-subspace.}\}$$

There is a surjective map

$$\text{span}: V_{k,n} \rightarrow Gr(k, n) \quad (v_1, \dots, v_k) \mapsto \text{span}(v_1, \dots, v_k).$$

we give $Gr(k, n)$ the quotient topology. Next time we will see $Gr(k, n)$ is a compact manifold of dimension $k \cdot (n - k)$.

The map $Gr(k, n) \mapsto Gr(n - k, n)$ given by $L \mapsto L^\perp$ is a homeomorphism.

[07.05.2025, Lecture 9]

[12.05.2025, Lecture 10]

Example. We have $Gr(1, n) = \mathbb{R}P^{n-1}$.

Theorem 3.8: Grassmann Manifolds

$Gr(k, n)$ is a compact manifold of dimension $k \cdot (n - k)$.

Proof. We first show compactness. Quasicompactness is clear, as it is a quotient space of a compact space.

We will show Hausdorff by constructing an injection into a Hausdorff space. For $V \in Gr(k, n)$ we consider the orthogonal projection $p_V: \mathbb{R}^n \rightarrow \mathbb{R}^n$. Let (v_1, \dots, v_k) be an orthonormal basis. then

$$p_V(x) = \langle x, v_1 \rangle \cdot v_1 + \dots + \langle x, v_k \rangle \cdot v_k$$

We will sometimes also write $p_V: \mathbb{R}^n \rightarrow \mathbb{R}^k$.

The map $Gr(k, n) \rightarrow \text{Hom}_{\mathbb{R}}(\mathbb{R}^n \rightarrow \mathbb{R}^n)$ given by $V \mapsto p_V$ is injective. Claim: this map is continuous.

By the quotient topology, we need to show, that the composite $V_{k,n} \rightarrow Gr(k, n) \rightarrow \text{Hom}_{\mathbb{R}}(\mathbb{R}^n, \mathbb{R}^n)$ is continuous. This map is

$$(v_1, \dots, v_k) \mapsto \sum_{i=1, \dots, k} \langle _, v_i \rangle \cdot v_i$$

and as a sum of continuous maps it is continuous. Because $Gr(kn)$ admits an injective continuous map to a Hausdorff space, it is Hausdorff.

Manifold property. Let $V \in Gr(k, n)$ be any k -plane. Set $U := \{L \in Gr(k, n) : L \cap V^\perp = \{0\}\}$. Claim: U is an open subset of $Gr(k, n)$. We choose an orthonormal basis (v_1, \dots, v_k) of V .

Claim. $\text{span}^{-1}(U) = \{(l_1, \dots, l_k) : \det(\langle l_i, v_j \rangle)_{1 \leq i, j \leq k} \neq 0\} \subseteq V_{k,n}$.

If we show this, we are done, as $\det \neq 0$ is an open condition.

Note. V^\perp is the kernel of $p_V: \mathbb{R}^n \rightarrow \mathbb{R}^n$. So $L \cap V^\perp = \{0\} \Leftrightarrow pr|_L: L \rightarrow V$ is injective.

As $\dim(L) = \dim(V) = k$, this is equal to $pr|_L: L \rightarrow V$ is bijective. Since $(\langle l_i, v_j \rangle)_{1 \leq i, j \leq k}$ is the matrix that expresses $(pr)|_L$ in terms of the basis $(l_i)_{1 \leq i \leq k}$ and $(v_j)_{1 \leq j \leq k}$, this is equivalent to $\det(\langle l_i, v_j \rangle) \neq 0$.

The map $V_{k,n} \rightarrow \mathbb{R}, (l_1, \dots, l_k) \mapsto \det(\langle l_i, v_j \rangle)$ is continuous, so $\text{span}^{-1}(U)$ is open in $V_{k,n}$, hence U is open in $Gr(k, n)$.

Next, we exhibit a homeomorphism

$$\begin{array}{ccc} & \Psi & \\ U & \xrightarrow{\quad} & \text{Hom}_{\mathbb{R}}(V, V^\perp) \\ & \Gamma & \end{array}$$

We then use $\dim(V) = k, \dim(V^\perp) = n - k$, so $\text{Hom}_{\mathbb{R}}(V, V^\perp) \cong \mathbb{R}^{k(n-k)}$.

Note that $\Gamma(f) \cap V^\perp = \{v \oplus f(V) : v = 0\} = \{0, 0\}$.

We define $\Gamma: \text{Hom}(V, V^\perp) \rightarrow U$ using that $\mathbb{R}^n = V \oplus V^\perp$. Then

$$\Gamma(f: V \rightarrow V^\perp) = \text{Graph of } f = \{v \oplus f(v) : v \in V\}$$

The graph map factors as the composite after choice of orthonormal basis v_1, \dots, v_k of V as

$$\text{Hom}_{\mathbb{R}}(V, V^\perp) \xrightarrow{\text{Gram-Schmidt}} V_{k,n} \xrightarrow{\text{span}} Gr(k, n)$$

so Γ is a continuous map.

We define $\Psi: U \rightarrow \text{Hom}_{\mathbb{R}}(V, V^\perp)$ as follows: If $L \in U$, then $p_V|_L: L \rightarrow V$ is a linear isomorphism.

We define $\Psi(L)$ as the composite $V \xrightarrow{(p_V|_L)^{-1}} L \xrightarrow{(p_{V^\perp}|_L)} V^\perp$.

This is inverse to Γ by go check yourself.

For Continuity of $\Psi: U \rightarrow \text{Hom}_{\mathbb{R}}(V, V^\perp)$. Since $\text{span}: V_{k,n} \rightarrow Gr(k, n)$ is a quotient map, so is its restriction

$$\text{span}: \text{span}^{-1}(U) \rightarrow U$$

So it suffices to show, that the composite

$$\text{span}^{-1}(U) \rightarrow U \xrightarrow{\Psi} \text{Hom}_{\mathbb{R}}(V, V^\perp)$$

is continuous.

To prove that, we choose orthonormal bases (v_1, \dots, v_k) of V and w_1, \dots, w_{n-k} of V^\perp . Expressing a linear map in the basis is a linear isomorphism

$$\text{Hom}_{\mathbb{R}}(V, V^\perp) \cong M(k \times (n - k), \mathbb{R}).$$

So we only need to show that

$$\text{span}^{-1}(U) \xrightarrow{\text{span}} U \xrightarrow{\Gamma} \text{Hom}_{\mathbb{R}}(V, V^\perp) \cong M(k \times (n - k), \mathbb{R})$$

is continuous. Did not copy the argument. Something about how we just compose matrices. \square

Corollary 3.9. The map $Gr(k, n) \rightarrow P_{k,n} := \{q \in \text{Hom}_{\mathbb{R}}(\mathbb{R}^n, \mathbb{R}^n) : q^2 = q = q^*, \text{trace}(q) = k\}$ is a homeomorphism.

Corollary 3.10. For all $0 \leq k \leq n$, the map $Gr(k, n) \rightarrow Gr(n - k, n)$ given by $V \mapsto V^\perp$ is an homeomorphism.

Proof. We need only show continuity.

$$\begin{array}{ccc} Gr(k, n) & \xrightarrow{V \mapsto V^\perp} & Gr(n - k, n) \\ \parallel & & \parallel \\ P_{k,n} & \xrightarrow{f \mapsto \text{Id} - f} & P_{n-k,n} \end{array}$$

\square

We can define the complex analogue: $Gr^{\mathbb{C}}(k, n) = \{L \subseteq \mathbb{C}^n : L \text{ is a complex linear subspace}\}$ with quotient topology by $V_{k,n}^{\mathbb{C}} \xrightarrow{\text{span}} Gr^{\mathbb{C}}(k, n)$ is a compact manifold of dimension $2k \cdot (n - k)$.

We can even define this for Quaternions:

$$Gr^{\mathbb{H}}(k, n) = \{L \subseteq \mathbb{H}^n : \mathbb{H}\text{-right submodule of dimension } k\}$$

is a compact manifold of dimension $4 \cdot k \cdot (n - k)$.

Bigger picture. The orthogonal group $O(n)$ acts transitively on $V_{k,n}$. This gives an isomorphism $O(n)/(1 \times O(n - k)) \rightarrow V_{k,n}$, that is even a homeomorphism.

Similarly, we have a transitive action $O(n) \rightarrow Gr(k, n)$. Looking at the stabilizer of \mathbb{R}^k . We get $O(n)/O(k) \times O(n - k) \xrightarrow{\cong} Gr(k, n)$ an homeomorphism.

This works similarly for complex and quaternionic Stiefel/Graßmann manifolds. This can be summarized as: „Stiefel manifolds and Grassmannians are homogenous spaces“.

Fact. Let G be a liegroup. H a closed subgroup. Then G/H is a (smooth) manifold of dimension $\dim G - \dim H$.

3.1 Orientations

We will (again) try $H_n(X)$ for $H_n(X, \mathbb{Z})$ and see how long it takes for him to forget he wanted to drop the \mathbb{Z} from the notation.

Notation 3.11. For $Y \subseteq X$, write $H_n(X | Y) := H_n(X, X \setminus Y; \mathbb{Z})$, we call the „local homology of X at Y “.

This is because for $Y \subseteq U \subseteq X$, U a neighborhood of Y , then excision gives

$$H_n(U | Y) = H_n(U; U \setminus Y; \mathbb{Z}) \xrightarrow{\cong} H_n(X, X \setminus Y; \mathbb{Z}) = H_n(X | Y)$$

If M is an m -manifold, and $x \in M$, then $H_n(X | x) = H_n(X | \{x\})$. This is \mathbb{Z} iff $m = n$ and else 0.

Definition 3.12: Local orientation

Let M be an m -manifold. A local orientation of M at $x \in M$ is a generator of $H_m(X | x)$.

There are exactly two local orientations at every point.

Construction 3.13 (Orientation covering). Let M be an m -manifold. We define the set $\tilde{M} = \{(x, \mu) : x \in M, \mu \text{ is a local orientation at } x\}$. This comes with a map $p: \tilde{M} \rightarrow M$, $p(x, \mu) = x$. This map is surjective and every point in M has exactly two preimages.

A subset B of M is a *Local ball* if B is a local subset of M , such that there exists a homeomorphism $\phi: \mathbb{R}^n \rightarrow M$ onto some open subset, such that $\phi(\langle D \rangle^n) = B$.

Note. If B is a local ball in M , then $M \setminus B \rightarrow M \setminus \{x\}$ is a homotopy-equivalence (here we need the special definition of open ball). This induces an isomorphism $r_x^B: H_m(M | B) \rightarrow H_m(X | x) \cong \mathbb{Z}$ for all $x \in B$. If μ is a local orientation at x , i.e. a generator of $H_m(X | B)$, we set $U(B, \mu) = \{(x, r_x^B(\mu)) : x \in B\} \subseteq \tilde{M}$.

Theorem 3.14: Orientation covering

Let M be an m -manifold.

1. As (B, μ) varies over all pairs of local balls B and generators μ of $H_m(M \mid B)$, the subset $U(B, \mu)$ of \tilde{M} are the basis of a topology on \tilde{M} .
2. In this topology on \tilde{M} , the map $p: \tilde{M} \rightarrow M$, $p(x, \mu) = x$ is a twofold covering, the orientation covering of M .
3. \tilde{M} is an m -manifold.

Proof. 1. We need to show, that for all local balls B, B' and all generators $\mu \in H_m(X \mid B)$, $\mu' \in H_m(X \mid B')$, the set $U(B, \mu) \cap U(B', \mu')$ is a union of basiss sets. Let $(x, \nu) \in U(B, \mu) \cap U(B', \mu')$. so $x \in B \cap B'$. and $r_x^B(\mu) = r_x^{B'}(\mu') := \nu$.

Choose a smaller local ball, s.t. $x \in B'' \subseteq B \cap B'$. We consider the following diagram of local homology groups:

$$\begin{array}{ccccc}
 H_m(X \mid B) & & & & \\
 \downarrow & \searrow \cong & & \searrow \cong & \\
 H_m(X \mid B \cap B') & \longrightarrow & H_m(X \mid B'') & \xrightarrow[\cong]{r_x^{B''}} & H_m(X \mid x) \\
 \uparrow & \nearrow \cong & & \nearrow \cong & \\
 H_m(X \mid B') & & & &
 \end{array}$$

so μ and μ' map to the same generator of $H_m(X \mid B'')$. Set $\mu'' = \text{incl}_*(\mu) = \text{incl}'_*(\mu')$. Then $(x, \nu) \in U(B'', \mu'') \subseteq U(B, \mu) \cap U(B', \mu')$

2. Because M is a manifold, the local balls form a basis of a topology of M . So it suffices to establish for all local balls B in M a homeomorphism

$$\begin{array}{ccc}
 p^{-1}(B) & \cong & B \amalg B \\
 \downarrow p & \swarrow \text{fold} & \\
 B & &
 \end{array}$$

I did not manage to copy the rest of this argument.

3. is a special case of

Proposition 3.15. Let $p: N \rightarrow M$ be a covering map and M an m -manifold. Then N is an m -manifold.

Proof. Hausdorff is clear.

For $y \in N$ choose an open neighborhood U of $p(y) = x$ in M , such that $U \cong \mathbb{R}^m$ and p is locally trivial over U . Choose a homeomorphism of $p^{-1}(U) \cong U \times F$ for F some discrete space. Then $U \times f$ is again homeomorphic to \mathbb{R}^n and its preimage is an open neighborhood of $y \in N$. □

□

[12.05.2025, Lecture 10]
[19.05.2025, Lecture 11]¹

¹I recently got a new laptop (including a new keyboard), which may negatively affect my writing speed and subsequently quality of the script for the next few lectures

Definition 3.16: Orientation

An orientation of an m -manifold is a continuous section $s: M \rightarrow \tilde{M}$ of p the orientation covering.

Definition 3.17: Orientability

An manifold is orientable, if it has an orientation.

Remark 3.18. If M is connected and orientable, $s: M \rightarrow \tilde{M}$ an orientation. Then $\tau \circ s: M \rightarrow \tilde{M}$ is another orientation where $\tau: \tilde{M} \rightarrow \tilde{M}, (x, \mu) \mapsto (x, -\mu)$.

Then $M \amalg M \cong \tilde{M}$ is a homeomorphism and $p: \tilde{M} \rightarrow M$ is the trivial covering.

So for connected M , the following are equivalent: M is orientable, $p: \tilde{M} \rightarrow M$ has a continuous section, p is trivial, i.e. $\tilde{M} \cong M \amalg M$.

An orientable connected manifold has exactly two orientations

If M is orientable and has n path-components, then M has exactly 2^n orientations.

Corollary 3.19. Let M be a connected m -manifold such that for some (hence any) $x \in M$, the group $\pi_1(M, x)$ does not have a subgroup of index 2. Then M is orientable.

Proof. We argue by contradiction. If M was not orientable, then $p: \tilde{M} \rightarrow M$ is not a product covering. So \tilde{M} is path connected, so for every $\tilde{x} \in \tilde{M}$ the homomorphism on fundamental groups $p_*: \pi_1(\tilde{M}, \tilde{x}) \rightarrow \pi_1(M, p(\tilde{x}))$ is a monomorphism with image of index 2. So $\text{Im}(p_*)$ is an index 2 subgroup of $\pi_1(M, x)$. \square

This gives that in particular every simply connected manifold is orientable.

Example 3.20. The spaces S^n (for $n \geq 2$), $\mathbb{C}P^n, \mathbb{H}P^n$ are orientable manifolds.

He continues to draw, that S^1 is also orientable.

Let M be an m -manifold, that is also a topological group, i.e. there is a continuous map $m: M \times M \rightarrow M$ that is also a group structure on M and such that $m \mapsto m^{-1}$ is continuous. Then M is orientable.

Proof. choose a local orientation $\mu \in H_m(M | 1)$, where $1 \in M$ is the multiplicative unit. For every $m \in M$,

$$m \cdot _ : M \rightarrow M$$

is a homeomorphism that takes 1 to m , so $(m \cdot _)*: H_m(M | 1) \rightarrow H_m(M | m)$ is an isomorphism. Set $\mu_m := (m \cdot _)*(\mu)$. Then $\{\mu_m\}_{m \in M}$ is an orientation of M . \square

Examples for this are $S^1, O(n), U(n), \text{Sp}(n), SO(n), SU(n), \dots$

Proposition 3.21. Let M be any n -manifold.

1. The manifold \tilde{M} is orientable, and the map $\tau: \tilde{M} \rightarrow \tilde{M}$ given by $\tau(x, \mu) = (x, -\mu)$ is orientation reversing.
2. Let $q: N \rightarrow M$ be a twofold covering and N be orientable manifold, $\tau: N \rightarrow N$ the free deck-transformation. If $\tau: N \rightarrow N$ is orientation reversing, then $q: N \rightarrow M$ is isomorphic as a covering to $p: \tilde{M} \rightarrow M$.

Proof.

1. Let $\tilde{x} = (x, \mu) \in \tilde{M}$ be any point in \tilde{M} . Since $p: \tilde{M} \rightarrow M$ is a local homeomorphism. so $p: H_n(\tilde{M}, \tilde{x}) \xrightarrow{\cong} H_n(M, x) \ni \mu$. Set $\mu_{\tilde{x}} := p_*^{-1}(\mu)$. then $\{\mu_{\tilde{x}}\}_{\tilde{x} \in \tilde{M}}$ is an orientation of \tilde{M} .

The map $\tau: \tilde{M} \rightarrow \tilde{M}$ reverses this orientation.

$$\tau_*: H_n(\tilde{M}, \tilde{x}) \rightarrow H_n(\tilde{M}, \tau_*(\tilde{x})) \quad p_*^{-1}(\mu) \mapsto \tau_*(p_*^{-1}(\mu)) = p_*^{-1}(\mu) = -p_*^{-1}(-\mu) = \mu_{\tau(\tilde{x})}$$

2. We have

$$\begin{array}{ccc} N & & \tilde{M} \\ & \searrow q & \swarrow p \\ & M & \end{array}$$

and we look for a map $N \rightarrow \tilde{M}$. Define $f: N \rightarrow \tilde{M}$ by $f(y) = (q(y), q_*(\mu_y))$. We use $q_*: H_n(N | y) \xrightarrow{\cong} H_n(M | q(y))$. this f is continuous. We will not check this, f commutes with the free involution:

$$f(\tau y) = (q(y), q_*(\mu_{\tau y})) = (q(y), q_*(-\tau_*(\mu_y))) = (q(y), -q_*(\mu_y)) = \tau(q(y), q_*(\mu_y)) = \tau(f(y))$$

So f is a continuous bijection over M , hence a homeomorphism.

□

3.1.1 Orientability of $\mathbb{R}P^n$

We already know $\mathbb{R}P^1$ and $\mathbb{R}P^3$ are orientable, as $\mathbb{R}P^1 \cong S^1$ and $\mathbb{R}P^3 \cong SO(3)$.

Recall. The antipodal map $A: S^n \rightarrow S^n$, given by $x \mapsto -x$ has degree $(-1)^{n+1}$.

Let $\mu \in H_n(S^n, \mathbb{Z})$ be any generator, define an orientation on S^n by $x \in S^n : \mu_x := r_x^{S^n}(\mu) \in H_n(S^n(x))$.

We look at

$$\begin{array}{ccc} H_n(S^n, \mathbb{Z}) & \xrightarrow[r_x^{S^n}]{\cong} & H_n(S^n | x) \\ \downarrow A_* & & \downarrow A_* \\ H_n(S^n, \mathbb{Z}) & \xrightarrow{\cong} & H_n(S^n | -x) \end{array}$$

Some of this diagram is missing.

this gives if n is even, then $A: S^n \rightarrow S^n$ is orientation reversing. if n is odd, then $A: S^n \rightarrow S^n$ is orientation preserving:

So for even n , then $q: S^n \rightarrow \mathbb{R}P^n$ is twofold covering and flip reverses orientation, so this „is“ the orientation covering. As $S^n \not\cong \mathbb{R}P^n \amalg \mathbb{R}P^n$ we have no continuous section to $S^n \xrightarrow{q} \mathbb{R}P^n$ and $\mathbb{R}P^n$ is not orientable.

For n odd we have

Proposition 3.22. Let $f: N \rightarrow N$ be continuous free involution of a connected oriented m -manifold. Then

1. $M := N/x \sim f(x)$ is an m -manifold.
2. If N is orientable and f is orientation preserving, then M is orientable.

Proof. 1. We have implicitly already done he's trying to convince me.

2. Choose an orientation $\{\mu_y\}_{y \in N}$ of N . We define an orientation of M as follows: For $x \in M$ choose $y \in N$, such that $p(y) = x$. Then we have

$$p_*: H_n(N | y) \xrightarrow{\cong} H_n(M | x)$$

and set $\mu_x := q_*(\mu_y)$. This is independent of the choice of y : the other choice is $f(y)$.

Some diagram I couldn't copy. As f is orientation preserving, the choice does not matter.

□

Then $\mathbb{R}P^n$ is orientable for n odd.

Next. We want to show, that for a m -manifold and all $n > m$: $H_n(M, A) = 0$. As a slogan „Homology vanishes above the geometric dimension.“

In many examples, m -manifolds M have m -dimensional CW-structure, e.g.

$$S^n, \mathbb{R}P^n, \mathbb{H}P^n, \mathbb{C}P^n$$

We could also produce CW-structure on the Grassmannians and Stiefel-Manifolds.

Warning. an m -manifold need not admit a CW-structure! Smooth manifolds admit triangulations, hence CW-structures. But there are non-smoothable manifolds, that do not admit CW-structures.

Theorem 3.23: Vanishing homology in high dimensions

Let M be an m -manifold; let K be a compact subset of M . Then

1. $H_i(M, M \setminus K; A) = 0$ for all $i > n$ and A any abelian group.
In particular, if M is compact, then $H_i(M; A) = 0$ for all $i > n$ and all A .
2. A class in $H_n(M, M \setminus K, A)$ is zero if and only if for all $x \in K$, its image under $r_x^K: H_n(M, M \setminus K, A) \rightarrow H_n(M, M \setminus \{x\}; A)$ is zero.
In particular, if M is compact, the maps $r_x: H_n(M, A) \rightarrow H_n(M, M \setminus \{x\}; A)$ are jointly injective.

Proof. In 6 bootstrapping-steps.

Step 1 If $M = \mathbb{R}^n$, and K is a convex compact subset. Choose $R > 0$ such that K is contained in the open ball of radius R around $y \in K$.

$$\{z \in \mathbb{R}^n : |z - y| = R\} =: S_R^{n-1} \subseteq M \setminus K \subseteq M \setminus \{y\}$$

and these are homotopy equivalences.

So $H_i(M | K) \cong H_i(M | x) = 0$ for $i > n$.

Step 2 K_1, K_2 two compact subsets of M . Suppose the claim holds for K_1, K_2 and $K_1 \cap K_2$. Then it also holds for $K_1 \cup K_2$.

We do this by a Mayer-Vietoris argument for local homology.

$$M \setminus (K_1 \cap K_2) = (M \setminus K_1) \cup (M \setminus K_2)$$

$$\text{And } M \setminus K_1 \cap M \setminus K_2 = M \setminus (K_1 \cup K_2)$$

Remembering the theorem of small simplices we get long fractions of chain complexes, which I did not copy. We get a long exact sequence of homology groups

$$H_{n+1}(M | K_1 \cap K_2) \xrightarrow{\partial} H_n(M | K_1 \cup K_2) \rightarrow H_n(M | K_1) \oplus H_n(M | K_2) \rightarrow H_n(M | K_1 \cap K_2)$$

For $i > n$ we have $H_i(M \mid K_1 \cup K_2)$ lies between $H_{i+1}(M \mid K_1 \cap K_2) = 0$ and $H_i(M \mid K_1) \oplus H_i(M \mid K_2) = 0$.

For $i = n$

$$0 = H_{n+1}(M \mid K_1 \cap K_2) \rightarrow H_n(M \mid K_1 \cup K_2) \hookrightarrow H_n(M \mid K_1) \oplus H_n(M \mid K_2)$$

Let $Z \in H_n(M \mid K_1 \cup K_2)$ such that $r_x^{K_1 \cup K_2}(z) = 0$ for all $x \in K_1 \cup K_2$.

Claim. $r_{K_1}^{K_1 \cup K_2}(z) = 0$. To see this pick $x \in K_1$. Then

$$\begin{array}{ccc} H_n(M \mid K_1 \cup K_2) & \xrightarrow{r_{K_1}^{K_1 \cup K_2}} & H_n(M \mid K_1) \\ & \searrow & \downarrow r_x^{K_1} \\ & & H_n(M \mid x) \end{array}$$

For all $x \in K_1$, $r_x^{K_1}(r_{K_1}^{K_1 \cup K_2}(z)) = 0$ and so $r_{K_1}^{K_2 \cap K_1}(z) = 0$ because the claim ii) holds for K_1 . Similarly $r_{K_2}^{K_1 \cap K_2}(z) = 0$ and then also $z = 0$ by the injectivity of

$$(r_{K_1}^{K_1 \cap K_2}, r_{K_2}^{K_1 \cup K_2})$$

[19.05.2025, Lecture 11]
[21.05.2025, Lecture 12]

Interlude.

Construction 3.24. Let $\mu \in H_m(M \mid x)$ be a local orientation of a manifold M at x . Let $\nu \in H_n(N \mid y)$ similarly. We construct the relative Künneth isomorphism

$$\begin{array}{ccc} H_m(M \mid x) \otimes H_n(N \mid y) & & H_{m+n}(M \times N \mid (x, y)) \\ \parallel & & \parallel \\ H_m(M, M \setminus \{x\}; \mathbb{Z}) \otimes H_n(N, N \setminus \{y\}; \mathbb{Z}) & \xrightarrow{\times} & H_{m+n}(M \times N, M \times (N \setminus \{y\}) \cup (M \setminus \{x\}) \times N; \mathbb{Z}) \end{array}$$

„We can combine local orientations to get some in the product.“ $\mu \times \nu$ is a local orientation of $M \times N$ at (x, y) . Because we didn't make any choices, this is probably continuous. Let $\{\mu_x\}_{x \in X}$ and $\{\nu_y\}_{y \in Y}$ be orientation of M and N . Then $\{\mu_x \times \nu_y\}_{(x,y) \in X \times Y}$ is an orientation of $M \times N$.

End of interlude.

Step 3 Let $M = \mathbb{R}^n$, $K = K_1 \cup \dots \cup K_m$ for K_1, \dots, K_m being compact convex subsets of $M = \mathbb{R}^n$.

Proof. By induction on m . $m = 1$ is clear by step 1.

Now let $K = (K_1 \cup \dots \cup K_{m-1}) \cup K_m$ where we have for both terms the conditions by induction. We need to look at

$$(K_1 \cup \dots \cup K_{m-1}) \cap K_m = (K_1 \cap K_m) \cup \dots \cup (K_{m-1} \cap K_m)$$

where we use that the intersection of convex subsets is again convex. Then the union is again fulfilling the conditions by induction. \square

Step 4 $M = \mathbb{R}^n$ and K any compact subset of \mathbb{R}^n . We need the following claim.

Claim. Let $\alpha \in H_i(\mathbb{R}^n | K)$. Then there is a compact neighborhood N of K and a class $\alpha' \in H_i(\mathbb{R}^n | N)$ such that $\alpha \in r_K^N(\alpha')$.

Proof. Let $\alpha = [x + C_i(\mathbb{R}^n \setminus K)]$ for some $x \in C_i(\mathbb{R}^n)$ such that $d_i(x) \in C_{i-1}(\mathbb{R}^n \setminus K)$. Then

$$d_i(x) = \sum_{\text{finite}} \alpha_j \cdot (f_j: \nabla^{i-1} \rightarrow \mathbb{R} \setminus K)$$

with $\alpha_j \in A$ and f_j continuous. Set $L = \text{supp}(d_i(x)) := \bigcup f_j(\nabla^{i-1})$ a compact subset of $\mathbb{R}^n \setminus K$. Now K, L are disjoint subsets of \mathbb{R}^n . so there is an N , compact neighborhood of K with $N \cap L = \emptyset$. You can for example look at $\text{dist}(L, K) > 0$ and take N accordingly.

Then $d_i(x) \in C_{i-1}(L) \subseteq C_{i-1}(\mathbb{R}^n \setminus N)$ so $\alpha' := [x + C_i(\mathbb{R}^n \setminus N)] \in H_i(\mathbb{R}^n | N)$ satisfies $r_K^N(\alpha') = \alpha$. \square

Now for the proof of step 4: either $i > n$ or $i = n$ and $\text{res}_x^K(\alpha) = 0$ for all $x \in K$.

Let N be a compact neighborhood of K and $\alpha' \in H_i(\mathbb{R}^n | N)$ with $r_K^N(\alpha') = \alpha$. Since K is compact, it can be covered with finitely many metric open balls in N . Let b_1, \dots, b_m be the closed metric balls, which still lie in N . Now $K \subseteq B_1 \cup \dots \cup B_m \subseteq N$ and B_i is a convex compact subset, so laim i) and ii) hold for $B_1 \cup \dots \cup B_m$. So

$$r_{B_1 \cup \dots \cup B_m}^n(\alpha') = 0$$

if $i > n$. and

$$\text{res}_X^{B_1 \cup \dots \cup B_m}(\text{res}_{B_1 \cup \dots \cup B_m}^N(\alpha')) = 0$$

for $i = n, x \in B_1 \cup \dots \cup B_m$. And then

$$r_{B_1 \cup \dots \cup B_m}^N(\alpha') = 0$$

by step 3. and then $\alpha = r_K^N(\alpha') = r_K^{B_1 \cup \dots \cup B_m}(\underbrace{r_{B_1 \cup \dots \cup B_m}^N(\alpha')}_{=0}) = 0$

Step 5 M arbitrary, there is an open neighborhood U of K homeomorphic to \mathbb{R}^n .

Choose a homeomorphism $\phi: U \xrightarrow{\cong} \mathbb{R}^n$. contemplate the commutative diagram

$$\begin{array}{ccc} H_i(M | K) & \xrightarrow[\cong]{\text{excision}} H_i(U | K) & \xrightarrow[\phi_*]{\cong} H_i(\mathbb{R}^n | \phi(K)) \\ \downarrow r_*^K & & \downarrow r_{\phi(x)}^{\phi(K)} \\ H_i(U | x) & \xrightarrow[\phi_*]{\cong} & H_i(\mathbb{R}^n | \phi(x)) \end{array}$$

Step 6 General case.

Claim. There are compact subsets K_1, \dots, K_m of M such that

- K_i is contained in an open subset homeomorphic to \mathbb{R}^m .
- $K = K_1 \cup \dots \cup K_m$.

When we prove this claim, we are done by step 2.

Proof of claim. Each $x \in K$ has an open neighborhood homeomorphic to \mathbb{R}^n . So there is a compact neighborhood N_x of x such that $x \in N_x \subseteq U \cong \mathbb{R}^n$. $K \subseteq \bigcup_{x \in K} N_x$ so by compactness $K = N_{x_1} \cup \dots \cup N_{x_m}$. \square

\square

3.2 The fundamental class

We know $H_n(S^n; \mathbb{Z}) \cong \mathbb{Z}$, $H_{k+l}(S^k \times S^l; \mathbb{Z}) \cong \mathbb{Z}$, $H_n(S^n \amalg S^n; \mathbb{Z}) \cong \mathbb{Z} \oplus \mathbb{Z}$ and $H_{2n}(\mathbb{C}P^n; \mathbb{Z}) \cong \mathbb{Z}$, $H_{4n}(\mathbb{H}P^n; \mathbb{Z}) \cong \mathbb{Z}$

$$H_n(\mathbb{R}P^n, \mathbb{Z}) \cong \begin{cases} \mathbb{Z} & n \text{ odd} \\ 0 & n \text{ even.} \end{cases}$$

$$H_m(\mathbb{R}^n; \mathbb{Z}) = 0 \text{ for } n \geq 1.$$

The pattern should be for connected manifolds, their top homology is \mathbb{Z} , if M is compact and orientable.

We want to show:

Proposition 3.25. Let M be a compact connected n -manifold. Then

$$H_n(M; \mathbb{Z}) \cong \begin{cases} \mathbb{Z} & \text{if } M \text{ is orientable} \\ 0 & M \text{ is not orientable.} \end{cases}$$

For that we first study fundamental classes.

Theorem 3.26: Fundamental Classes

Let M be an oriented n -manifold and K a compact subset of M . then there is a unique class $\mu_K \in H_n(M, M \setminus K; \mathbb{Z}) \cong H_n(M | K)$ such that

$$r_x^K(\mu_K) = \mu_x \in H_n(M | x)$$

for all $x \in K$.

The class μ_K is the relative fundamental class .

Special Case. Let M be a compact oriented m -manifold. Then there is a unique class $[M] \in H_n(M; \mathbb{Z})$ such that $r_x^K([M]) = \mu_x$ for all $x \in M$. Then $[M]$ is the fundamental class.

If in addition M is connected, then $[M]$ generates $H_n(M, \mathbb{Z}) \cong \mathbb{Z}$.

Proof. Uniqueness is already done. We make a construction in three steps.

Step 1 Suppose K is contained in an open subset U homeomorphic to \mathbb{R}^n . Then K is contained in a local ball B in the sense of the definition of the orientation covering. We consider the commutative diagram: For $x, y \in K$:

$$\begin{array}{ccccc} & & H_n(M | B) & & \\ & \swarrow r_x^B \cong & \downarrow r_K^B & \searrow r_y^B \cong & \\ & & H_n(M | K) & & \\ & \swarrow r_x^K & & \searrow r_y^K & \\ H_n(M | x) & & & & H_n(M | y) \end{array}$$

Then $r_K^B(\mu_B)$ has the desired property.

Step 2 Suppose that $K = K_1 \cup K_2$, K_1, K_2 compact, claim true for K_1 and K_2 . Let $\mu_{K_1} \in H_n(M | K_1)$ and $\mu_{K_2} \in H_n(M | K_2)$ be the relative fundamental classes. We showed in the proof of Step 2 of the previous result that the following is exact:

$$0 \rightarrow H_n(M | K) \xrightarrow{(r_{K_1}^K, r_{K_2}^K)} H_n(M | K_1) \oplus H_n(M | K_2) \rightarrow H_n(M | K_1 \cap K_2) \rightarrow \dots$$

so we see $(\mu_{K_1}, \mu_{K_2}) \mapsto r_{K_1 \cap K_2}^{K_1}(\mu_{K_1}) - r_{K_1 \cap K_2}^{K_2}(\mu_{K_2})$, where both terms are equal to $\mu_{K_1 \cap K_2}$, so their difference is 0. Any $x \in K$ is contained in $K_1 \cap K_2$. If $x \in K_1$, then

$$r_x^K(\mu_K) = r_x^{K_1}(r_{K_1}^K(\mu_K)) = r_x^{K_1}(\mu_{K_1}) = \mu_x$$

Step 3 General case. As in step 6 of the previous proof, we can write $K = K_1 \cup \dots \cup K_m$ where all K_i are compact and all K_i are contained in an open subset homeomorphic to \mathbb{R}^n . Then μ_{K_i} exists for all $i = 1, \dots, m$ by step 1. By induction on m and Step 2 it holds for $K = K_1 \cup \dots \cup K_m$. □

We draw some corollaries:

Theorem 3.27: Top Homology of connected Manifolds

Let M be a connected, compact, oriented n -manifold. Then $H_n(M; \mathbb{Z}) \cong \mathbb{Z}$, generated by $[M]$.

Moreover, for all $x \in M$, the restriction $r_x^M: H_n(M, \mathbb{Z}) \rightarrow H_n(M | x)$ is an isomorphism.

Proof. Claim. Let $\alpha \in H_n(M, \mathbb{Z})$. Then the set $x \in \{M \mid r_x^M(\alpha) = 0 \text{ in } H_n(M | x)\}$ is open and closed in M .

Let $x \in M$ be such that $r_x^M(\alpha) = 0$ (or $r_x^M(\alpha) \neq 0$). Let B be a local ball in M containing x . We get a commutative diagram: For $x, y \in B$

$$\begin{array}{ccc} & H_n(M) & \\ & \downarrow r_B^M & \\ & H_n(M | B) & \\ \swarrow \cong \quad r_x^B & & \searrow \cong \quad r_y^B \\ H_n(M | x) & & H_n(M | y) \end{array}$$

For some reason, this shows the claim.

Then $r_x^M: H_n(M; \mathbb{Z}) \rightarrow H_n(M | x) \cong \mathbb{Z}$ is surjective, because $r_x^M([M]) = \mu_x$ generates $H_n(M | x)$. If $\alpha \in H_n(M, \mathbb{Z})$ is such that $r_x^M(\alpha) = 0$, then $\{y \in M : r_y^M(\alpha) = 0\}$ is open, closed and nonempty. Since M is connected, this set is all of M , so $\alpha = 0$ by the detection property (ii) in the previous theorem. □

Corollary 3.28. Let M be a compact connected and non-orientable n -manifold. Then $H_n(M; \mathbb{Z}) = 0$.

Proof. Let $\alpha \in H_n(M, \mathbb{Z})$. As in the previous proof, $\{x \in M : r_x^M(\alpha) = 0\}$ is an open and closed subset. Also $r_x^M: H_n(M, \mathbb{Z}) \rightarrow H_n(M | x)$ is injective for all $x \in M$. Hence $H_n(M; \mathbb{Z}) \hookrightarrow H_n(M | x) \cong \mathbb{Z}$, so $H_n(M; \mathbb{Z})$ is torsion free.

Let $p: \tilde{M} \rightarrow M$ be the orientation covering. Since M is connected and not orientable, \tilde{M} is connected, compact and orientable. Let $\{\mu_x\}_{x \in \tilde{M}}$ be an orientation of \tilde{M} . Then

$$H_n(\tilde{M}; \mathbb{Z}) \cong \mathbb{Z}$$

generated by $[\tilde{M}]$. Let $\tau: \tilde{M} \rightarrow \tilde{M}$ be the nonidentity deck transformation. Then τ is orientation reversing, $\tau_*[\tilde{M}] = -[\tilde{M}]$ (A diagram I didn't copy as proof).

So $p_*[\tilde{M}] = p_*(\tau_*([\tilde{M}])) = p_*(-[\tilde{M}]) = -p_*[\tilde{M}]$ so $2 \cdot p_*[\tilde{M}] = 0$. so by torsion-freeness $p_*([\tilde{M}]) = 0$.

As $[\tilde{M}]$ generates $H_n(\tilde{M}, \mathbb{Z})$, $p_* = 0: H_n(\tilde{M}; \mathbb{Z}) \rightarrow H_n(M; \mathbb{Z})$.

Let $\text{tr}: H_n(M, \mathbb{Z}) \rightarrow H_n(\tilde{M}; \mathbb{Z})$ be the transfer map. Then $0 = p_* \circ \text{tr} =$ multiplication by 2 on $H_n(M; \mathbb{Z})$ so $H_n(M, \mathbb{Z}) = 0$. \square

[19.05.2025, Lecture 12]

[26.05.2025, Lecture 13]

———— Was unfortunately unable to attend the lecture. ————

[26.05.2025, Lecture 13]

[28.05.2025, Lecture 14]

Chapter 4

Cap Product

Due to missing the last lecture I intended to copy the repetition Schwede does at the beginning of his lecture. However, public transport prevented me from copying the start of the Repetition. The most important definition of last lecture was the Cap-Product:

It will later be used for Poïncare-Duality.

Definition 4.1: Cap-Product

Let X be a simplicial set, $Y \subseteq X$ simplicial subset, R any commutative ring. For $0 \leq i \leq n$ we define

$$\cap: C_n(X, Y; R) \times C^i(X, Y, R) \rightarrow C_{n-i}(X; R)$$

given by

$$x \in C_n(X, Y; R) = \frac{R[X_n]}{R[Y_n]}, f \in C^i(X, Y; R), \text{ i.e. } f: X_i \rightarrow R, f(Y_i) = \{0\}$$

and

$$x \cap f = \underbrace{f(x[0, i])}_{\in R} \cdot x[i, n]$$

and then i missed something due to broken board

Proposition 4.2. Let $Y \subseteq X$ be simplicial sets, R some commutative ring. Then the following hold:

1. $d(x \cap f) = (-1)^i \cdot (dx \cap f - x \cap df)$ for $x \in C_n(X, Y; R), f \in C^i(X, Y; R)$
2. The cap product descends to a well defined and R -bilinear map

$$\cap: H_n(X, Y; R) \times H^i(X, Y; R) \rightarrow H_{n-1}(X, R), \quad [x] \cap [f] := [x \cap f]$$

3. If $Y = \emptyset$ $\xi \in H_n(X, R), \alpha \in H^i(X, R), \beta \in H^j(X, R)$ we have

$$(\xi \cap \alpha) \cap \beta = \xi \cap (\alpha \cup \beta)$$

4. if $Y = \emptyset$, then $\xi \cap 1 = \xi$.

5. Let $\Psi: X \rightarrow X'$ be a morphism of simplicial sets, s.t. $\Psi(Y) \subseteq Y', \xi \in H_n(X, Y; R), \alpha \in H^i(X', Y', R)$. Then

$$\Psi_*(\xi) \cap \alpha = \Psi_*(\xi \cap \Psi^*(\alpha))$$

Proof. 1. Was done last time.

2. Suppose $dx = 0, df = 0$, then

$$d(x \cap f) = (-1)^i \underbrace{((dx) \cap f)}_{=0} - x \cap \underbrace{df}_{=0} = 0$$

so $x \cap f$ is a cycle. For $y \in C_{n+1}(X, Y; R)$

$$(x + dy) \cap f = x \cap f \pm d(y \cap f)$$

which gives $[(x + dy) \cap f] = [x \cap f]$ and similarly $[x \cap (f + dg)] = [x \cap f]$ for $g \in C^{i-1}(X, Y; R)$.

3. Let $x \in C_n(X; R)$ represent $\xi \in H_n(X, R)$. Let $a: X_i \rightarrow R, b: X_j \rightarrow R$ represent $\alpha \in H^i(X, R)$ and $\beta \in H^j(X, R)$. Then

$$\begin{aligned} (x \cap a) \cap b &= (a(x[0, i]) \circ x[i, n]) \cap b \\ &= a(x[0, i]) \cdot (x[i, n] \cap b) \\ &= a(x[0, 1]) \cdot b(x[i, n][0, j]) \cdot x[i, n][j, n-1] \\ &= a(x[0, i]) \cdot b(x[i, i+j]) \cdot x[i+j, n] \\ &= a(x[0, i+j][0, i]) \cdot b(x[0, i+j][i, i+j]) \cdot x[i+j, n] \\ &= (a \cup b)(x[0, i+j]) \cdot x[i+j, n] \\ &= x \cap (a \cup b) \end{aligned}$$

4. For some reason clear.

5. We let $x \in X_n$ be an n -simplex, $f \in C^i(X', Y'; R)$ representing a chain in $H^i(X', Y'; R)$.

$$\begin{aligned} \Psi_*(x) \cap f &= \Psi_n(x) \cap f \\ &= f((\Psi_n(x))[0, i]) \cdot (\psi_n(x))[i, n] \\ &= f(\psi_i(x[0, i])) \cdot \Psi[n-i](x[i, n]) \\ &= \Psi_*(f(\Psi_i(x[0, i]))) \cdot x[i, n] \\ &= \Psi_*(x \cap \Psi^*(f)) \end{aligned}$$

□

Chapter 5

Poincaré Duality

5.1 Cohomology with compact support

We are working towards:

Theorem 5.1: Poïncare-duality

Let M be a compact n -manifold, $i \geq 0$.

- If M is oriented, then $[M] \cap _ : H^i(M; \mathbb{Z}) \rightarrow H_{n-i}(M; \mathbb{Z})$ is an isomorphism.
- The map $\nu_M \cap _ : H^i(M; \mathbb{F}_2) \rightarrow H_{n-i}(M; \mathbb{F}_2)$ is an isomorphism.

Our idea is to proof this by

- Proof for \mathbb{R}^n
- Patching/Mayer-Vietoris argument $M = U_1 \cup U_2$.

However, we have a problem: $M = \mathbb{R}$ is an oriented 1-manifold. $H^1(\mathbb{R}; \mathbb{Z}) \cong 0$, but $H_0(\mathbb{R}; \mathbb{Z}) \cong \mathbb{Z}$, which is not the same.

To solve this, we introduce Compactly supported cohomology as a variation of singular cohomology that has Poincaré duality for not necessarily compact manifolds: We will get

$$H_{comp}^i(M; \mathbb{Z}) \xrightarrow{\cong} H_{n-i}(M; \mathbb{Z})$$

for compact M .

Construction 5.2. Let X be a topological space, A an abelian group. A singular cochain $f \in C^n(\mathcal{S}(X), A)$

$$f : \mathcal{S}(X)_n = \text{maps}^{\text{cont}}(\nabla^n; X) \rightarrow A$$

is supported in a subset K of X if $f(\phi) = 0$ for all $\phi : \nabla^n \rightarrow X$ with $\phi(\nabla^n) \subseteq X \setminus K$. Equivalently f belongs to the kernel of the homomorphism

$$C^n(\mathcal{S}(X); A) \rightarrow C^n(\mathcal{S}(X \setminus K); A)$$

$f \in C^n(\mathcal{S}(X); A)$ is *compactly supported* if there is a compact subset K of X such that f is supported on K .

Proposition 5.3. Compactly supported cochains form a subcomplex of $C^*(\mathcal{S}(X); A)$.

Proof. Suppose $f : \mathcal{S}(X)_n \rightarrow A$ is supported on K with K compact. Then

$$(df)(\Psi) = \sum_{i=0, \dots, n+1} (-1)^i f(d_i^*(\Psi)) = \sum_{i=0, \dots, n+1} (-1)^i f(\Psi \circ (d_i)_*)$$

and then he outspeeded me. □

Example 5.4. There are no continuous maps $\phi: \nabla^n \rightarrow X$ with image in $X \setminus X$, so every cochain $f \in C^n(\mathcal{S}(X), A)$ is supported on X .

So if X is compact, then every cochain is compactly supported, hence $C_{comp}^*(X, A) = C^*(X, A)$.

Example 5.5. Note: $\{0\}$ is a compact subset of \mathbb{R}^n . So

$$\underbrace{C^*(\mathcal{S}(X), \mathcal{S}(X \setminus \{0\}); A)}_{\text{Cochains supported on } \{0\}} \subseteq C_{comp}^*(\mathcal{S}(X), A)$$

Proposition 5.6. For all $n \geq 1$, all abelian groups A , the inclusion

$$C^*(\mathcal{S}(\mathbb{R}^n); \mathcal{S}(\mathbb{R}^n \setminus \{0\}); A) \rightarrow C_{comp}^*(\mathcal{S}(\mathbb{R}^n); A)$$

is a quasi-isomorphism. In particular:

$$H_{comp}^i(\mathbb{R}^n; A) = H^i(\mathbb{R}^n; \mathbb{R}^n \setminus \{0\}; A) \cong \begin{cases} A & \text{if } i = n \\ 0 & \text{else} \end{cases}$$

Proof. By the five lemma, it suffices to show that the quotient complex

$$\frac{C_{comp}^*(\mathcal{S}(\mathbb{R}^n))}{C^*(\mathcal{S}(\mathbb{R}^n), \mathcal{S}(\mathbb{R}^n \setminus \{0\}))}$$

is acyclic. Let $f \in C_{comp}^i(\mathcal{S}(X))$ be a compactly supported cochain class in the quotient complex in a cocycle, i.e. $df \in C^{i+1}(\mathcal{S}(X), \mathcal{S}(X \setminus \{0\}))$. Then ?? of f is supported on $\{0\}$.

Since f is compactly supported and every compact subset of \mathbb{R}^n is contained in a sufficiently large ball, there is a $r > 0$ s.t. f is supported on $D_r^n = \{x \in \mathbb{R}^n : |x| \leq r\}$ the disc of radius r around 0. Write $\text{res}(f)$ for the restriction of f to $C^*(\mathcal{S}(X), \mathcal{S}(X \setminus D_r^n))$. The inclusion

$$\mathbb{R}^n \setminus D_r^n \rightarrow \mathbb{R}^n \setminus \{0\}$$

is a homotopy equivalence, so the cohomology of the pair $\mathbb{R}^n \setminus \{0\}; \mathbb{R}^n \setminus D_r^n$ is trivial. Equivalently, the residue cochain complex

$$C^*(\mathcal{S}(\mathbb{R}^n \setminus \{0\}), \mathcal{S}(\mathbb{R}^n \setminus D_r^n))$$

is trivial. So the cycle $\text{res}(f)$ in the acyclic complex is a coboundary. So there exists $g \in C^{i-1}(\mathcal{S}(\mathbb{R}^n \setminus \{0\}), \mathcal{S}(\mathbb{R}^n \setminus D_r^n))$ s.t. $dg = \text{res}(f)$. We extend g to a cochain

$$\tilde{g}: \mathcal{S}(X)_{i-1} \rightarrow A$$

by zero, i.e.

$$\tilde{g}(\phi) = \begin{cases} g(\phi) & \text{if } \phi(\nabla^{i-1}) \subseteq \mathbb{R}^n \setminus 0 \\ 0 & \text{else} \end{cases}$$

Then $\text{res}(\tilde{g}) = g$. In particular, \tilde{g} is supported on D_r^n because g is.

Then

$$\text{res}(d\tilde{g}) = d(\text{res}(\tilde{g})) = dg = \text{res}(f)$$

so $\text{res}(f - d\tilde{g}) = 0$ in $C^i(\mathcal{S}(\mathbb{R}^n \setminus \{0\}), \mathcal{S}(\mathbb{R}^n \setminus D_r^n))$. so $f - d\tilde{g}$ is supported on 0 , so ?? to $C^*(\mathcal{S}(\mathbb{R}^n); \mathcal{S}(\mathbb{R}^n \setminus \{0\}))$. Since $[f] = [f - d\tilde{g}] = 0$. \square

He talks about how this could also be done using some category theory.

Note. We can take $\mathbb{R}^n \rightarrow \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$ taking inclusion and projection. So we get $\mathbb{Z} \cong H_{comp}^n(\mathbb{R}^n; \mathbb{Z})$ is a retract of $H_{comp}^n(\mathbb{R}^{n+1}; \mathbb{Z}) \cong 0$. That doesn't make sense and gives the following

Warning! H_{comp}^* is not functorial for arbitrary continuous maps!

Compactly supported cohomology is

- contravariantly functorial in *proper* continuous maps
- covariantly functorial in open embeddings.

Definition 5.7: Proper maps

A continuous map $f: X \rightarrow Y$ is *proper* if for every compact subset K of Y , the set $f^{-1}(K)$ is compact with the subspace topology of X .

Example 5.8. Let X be a space. Then $X \rightarrow \{x\}$ is proper iff X is compact.

If K is compact and X is any space, then the projection $X \times K \xrightarrow{p} X$ is proper: For $L \subseteq X$ compact, $p^{-1}(L) = L \times K$ is compact.

Proposition 5.9. Let $\psi: X \rightarrow Y$ be a proper continuous map. Then the cochain map

$$\psi^*: C^*(\mathcal{S}(Y), A) \rightarrow C^*(\mathcal{S}(X), A)$$

takes compactly supported cochains to compactly supported cochains, so it restricts to a chain map

$$\psi^*: C_{comp}^*(\mathcal{S}(Y), A) \rightarrow C_{comp}^*(\mathcal{S}(X), A)$$

This restriction induces group homomorphisms $\psi^*: H_{comp}^i(Y; A) \rightarrow H_{comp}^i(X, A)$

Proof. We let $f: \mathcal{S}(Y)_n \rightarrow A$ be a simplicial cochain that is supported on the compact subset K of Y . then $\psi^{-1}(K)$ is compact because f is proper.

Claim. $\psi^*(f)$ is supported on $\psi^{-1}(K)$.

Let $\phi: \nabla^n \rightarrow X$ be continuous with image in $X \setminus \psi^{-1}(K)$. Then

$$(\psi^*(f))(\phi) = f(\psi \circ \phi) = 0$$

because $\psi \circ \phi$ has image in $Y \setminus K$. □

[28.05.2025, Lecture 14]

[02.06.2025, Lecture 15]

So f from the last proposition induces a homomorphism

$$f^*: H_{comp}^i(Y, A) \rightarrow H_{comp}^i(X, A)$$

and we get a contravariant functor in proper maps.

Today we will talk about

5.1.1 Covariant functoriality for open embeddings

This comes formally from understanding that compactly supported Cohomology is a colimit of some relative Cohomologies.

Let K be a compact subset of some space X . Then

$$C^*(\mathcal{S}(X), \mathcal{S}(X \setminus K)) \subseteq C_{comp}^*(X, A)$$

We write

$$\lambda_K: H^i(X, X \setminus K; A) \rightarrow H_{comp}^i(X, A)$$

for the induced map in cohomology.

For $K \subseteq L \subseteq X$, L also compact, then $X \setminus K \supseteq X \setminus L$ so

$$C^*(\mathcal{S}(X), \mathcal{S}(X \setminus K), A) \hookrightarrow C^*(\mathcal{S}(X), \mathcal{S}(X \setminus L); A) \subseteq C_{comp}^*(X, A)$$

giving a commutative triangle of cohomology groups:

$$\begin{array}{ccc} H^i(X, X \setminus K; A) & & \\ \downarrow \text{incl}^* & \searrow \lambda_K & \\ & & H_{comp}^i(X; A) \\ & \nearrow \lambda_L & \\ H^i(X, X \setminus L; A) & & \end{array}$$

Proposition 5.10. Let X be a Hausdorff space, A and B two abelian groups. Let $\alpha_K: H^i(X, X \setminus K; A) \rightarrow B$ be a homomorphism, for all compact subsets K of X , such that for all $K \subseteq L \subseteq X$, with L compact, the following commutes:

$$\begin{array}{ccc} H^i(X, X \setminus K; A) & & \\ \downarrow j_L^K & \searrow \alpha_K & \\ & & B \\ & \nearrow \alpha_L & \\ H^i(X, X \setminus L; A) & & \end{array}$$

Then there is a unique homomorphism $\alpha: H_{comp}^i(X, A) \rightarrow B$ such that $\alpha \circ \lambda_K = \alpha_K$ for all $K \subseteq X$ compact.¹

Proof. We drop the coefficients A from the notation (but only when he remembers he did this). Furthermore we set for this proof $n = i$.

Uniqueness We let $\alpha: H_{comp}^i(X) \rightarrow B$ be a homomorphism such that $\alpha \circ \lambda_K = \alpha_K$ for all K compact in X .

Let $f \in C_{comp}^n(X, A)$ be a compactly supported cochain. Let f be supported on the compact subset K . Then $f \in C^n(X, X \setminus K)$, and $\lambda_K[f] = [f]$. Hence

$$\alpha[f] = \alpha(\lambda_K[f]) = \alpha_K[f]$$

Existence/Construction Given $f \in C_{comp}^n(X, A)$ let K be a compact subset on which f is supported.

Claim. $\alpha[f] := \alpha_K[f]$ is independent of the choice of K .

Let K' be another compact subset of X on which f is supported. Let $L = K \cup K'$, another compact subset of X . We consider

$$\alpha_K[f] = \alpha_L(j_K^L[f]) = \alpha_L[f] = \alpha_L(j_{K'}^L[f]) = \alpha_{K'}[f]$$

Claim. $\alpha_K[f]$ only depends on the cohomology class of f .

¹The morphisms $\{\lambda_K\}_K$ express $H_{comp}^i(X, A)$ as a colimit of the groups $H^i(X, X \setminus K; A)$

Let $g \in C_{comp}^{n-1}(X, A)$ be any cochain. Then g is supported on some compact subset K' of X . Then f and g are supported on $L = K \cup K'$. So $\alpha[f] = \alpha_K[f] = \alpha_L[f + dg] = \alpha[f + dg]$, so $\alpha: H_{comp}^i(X, A) \rightarrow B$ is a well defined map and $\alpha \circ \lambda_K = \alpha_K$ for all K compact.

We still need to show, that α is a group homomorphism: Let $f, f' \in C_{comp}^n(X, A)$. Let f be supported on K , f' supported on K' , for K, K' compact in X . Then f and f' are both supported on $L = K \cup K'$, and

$$\alpha[f + f'] = \alpha_L[f + f'] = \alpha_L[f] + \alpha_L[f'] = \alpha[f] + \alpha[f']$$

□

Excursion 5.11 (filtered colimits.).

Definition. A PoSet (partially ordered set) is a pair (P, \leq) consisting of a set P and a relation \leq , that is

- reflexve
- transitive
- antisymmetric, i.e. if $x \leq y$ and $y \leq x$, then $x = y$.

A poset (P, \leq) is *filtered* if for all $x, y \in P$, there is $z \in P$ s.t. $x \leq z, y \leq z$.

Some examples are

- $(\mathbb{N}, \leq), (\mathbb{R}, \leq)$
- the set of finite subsets of a given set, under \subseteq .
- The set of finite dimension vector subspaces of a vector space under \subseteq
- the set of compact subsets $\mathcal{C}(X)$ in a Hausdorff space under \subseteq .

Every poset P, \subseteq gives rise to a category \underline{P} with

- $\text{Ob}(\underline{P}) = P$
- $\underline{P}(x, y) = \begin{cases} \{(x, y)\} & \text{if } x \leq y \\ \emptyset & \text{else} \end{cases}$

Composition is then uniquely defined.

A filtered colimit is a colimit over a filtered poset.

Example 5.12. Let X be a Hausdorff space, $\mathcal{C}(X)$ the poset of its compact subsets. A functor $\mathcal{C}(X) \rightarrow \mathbf{Ab}$ is given by $K \mapsto H^i(X, X \setminus K; A)$ and for $K \subseteq L$, we get j_L^K .

We can now say

$$C_{comp}^i(X, A) = \text{colim}_{\mathcal{C}(X)} C^*(X, X \setminus K; A)$$

Proposition (filtered colimits are exact). Let (P, \leq) be a filtered poset, $F, G, H: \underline{P} \rightarrow \mathbf{Ab}$ functors, $\phi: F \rightarrow G$ and $\psi: G \rightarrow H$ natural transformations of functors. Suppose that for all $x \in P$,

$$0 \rightarrow F(x) \xrightarrow{\phi(x)} G(x) \xrightarrow{\psi(x)} H(x) \rightarrow 0$$

is exact. Then the sequence

$$0 \rightarrow \text{colim}_{\underline{P}}(F) \xrightarrow{\text{colim}_{\underline{P}} \phi} \text{colim}_{\underline{P}} G \xrightarrow{\text{colim}_{\underline{P}} \psi} \text{colim}_{\underline{P}} H \rightarrow 0$$

is exact.

This might become an exercise.

Corollary 5.13. Let (P, \leq) be a filtered coposet, $F: \underline{P} \rightarrow \mathbf{Chains}$ a functor. Then the canonical morphism

$$\operatorname{colim}_P (H_n \circ F) \rightarrow H_n(\operatorname{colim}_P (F))$$

is an isomorphism.

Corollary 5.14.

$$H_{comp}^i(X, A) = H^i(\operatorname{colim}_{\mathcal{C}(X)} C^*(X, X \setminus K; A)) \xleftarrow{\cong} \operatorname{colim}_{\mathcal{C}(X)} H^i(C^*(X, X \setminus K; A)) = \operatorname{colim}_{\mathcal{C}(X)} H^i(X, X \setminus K; A)$$

Definition. Let (P, \leq) be a poset. Let $Q \subseteq P$ a subset. Q is *cofinal* in P , if for all $x \in P$, there is $y \in Q$ such that $x \leq y$.

If P is filtered, Q cofinal in P , then Q is also filtered.

Proposition. Let Q be a cofinal subset of a filtered Poset P . Let $F: \underline{P} \rightarrow \mathcal{C}$ be a functor to any category that has a colimit. Then $\operatorname{colim}_Q F = \operatorname{colim}_P F$.

Proof might be another exercise.

Example 5.15. $\mathcal{C}(\mathbb{R}^n) \subseteq \{D_r^n : r \geq 0\} \subseteq D_r^n : r \in \mathbb{N}$ are cofinal, where D_r^n again denotes the ball of radius r around 0. This gives

$$\begin{aligned} H_{comp}^i(\mathbb{R}^n; A) &= \operatorname{colim}_{K \in \mathcal{C}(X)} H^i(X, X \setminus K; A) \cong \operatorname{colim}_{\text{cofinal } r \in (\mathbb{N}, \leq)} H^i(\mathbb{R}^n, \mathbb{R}^n \setminus D_r^n; A) \\ &= \operatorname{colim}(H^i(\mathbb{R}^n, \mathbb{R}^n \setminus \{0\}) \xrightarrow{\cong} H^i(\mathbb{R}^n, \mathbb{R}^n \setminus D_1^n) \xrightarrow{\cong} \dots \xrightarrow{\cong} H^i(\mathbb{R}^n, \mathbb{R}^n \setminus D_k^n) \xrightarrow{\cong} \dots) \end{aligned}$$

which is by category theory already isomorphic to $H^i(\mathbb{R}, \mathbb{R}^n \setminus 0)$. This roughly represents the proof we did last week in more detail.

Construction 5.16. Let Y be a open subset of a Hausdorff space X . We'll define a homomorphism

$$\iota_Y^X: H_{comp}^i(Y; A) \rightarrow H_{comp}^i(X; A)$$

Let K be a compact subset of Y . We define

$$\alpha_K: H^i(Y, Y \setminus K; A) \rightarrow H_{comp}^i(X, A)$$

by looking at $H^i(Y, Y \setminus K; A) \xleftarrow[\text{excision}]{\cong} H^i(X, X \setminus K; A)$ is an isomorphism and we can invert it.

We use (X, Y, K) is excisive. We then compose with $\lambda_K: H^i(X, X \setminus K; A) \xrightarrow{\lambda_K} H_{comp}^i(X, A)$.

The maps α_K are compatible for $K \subseteq L \subseteq Y$ all compact.

$$\begin{array}{ccccc} & & & \alpha_K & \\ & & & \curvearrowright & \\ H^i(Y, Y \setminus K; A) & \xleftarrow{\cong} & H^i(X, X \setminus K; A) & & \\ \downarrow j_L^K & & \downarrow j_L^K & \searrow \lambda_K & \\ & & & & H_{comp}^i(X; A) \\ & & & \nearrow \lambda_L & \\ H^i(Y, Y \setminus L; A) & \xleftarrow{\cong} & H^i(X, X \setminus L; A) & & \\ & & & \curvearrowleft & \\ & & & \alpha_L & \end{array}$$

The universal property of $H_{comp}^i(Y, A)$ as a colimit yields a unique homomorphism ι_Y^X that admits a commutative square I couldn't copy for all $K \subseteq Y$ and K compact.

Proposition 5.17. Let X be a Hausdorff space.

1. The homomorphism $\iota_X^X: H_{comp}^i(X, A) \rightarrow H_{comp}^i(X, A)$ is the identity.
2. If $Z \subseteq Y \subseteq X$ with Z and Y open, then

$$\iota_Y^X \circ \iota_Z^Y = \iota_Z^X: H_{comp}^i(Z, A) \rightarrow H_{comp}^i(X, A)$$

Proof. We let K be any compact subset of Z .

$$\begin{array}{ccccc}
 & & \cong & & \\
 & & \text{excision} & & \\
 & \nearrow & & \searrow & \\
 H^i(X, X \setminus K; A) & \xrightarrow[\text{excision}]{\cong} & H^i(Y, Y \setminus K; A) & \xrightarrow[\text{excision}]{\cong} & H^i(Z, Z \setminus K; A) \\
 \downarrow \lambda_K & & \downarrow \lambda_K & & \downarrow \lambda_K \\
 H_{comp}^i(X, A) & \xleftarrow[\iota_Y^X]{} & H_{comp}^i(Y; A) & \xleftarrow[\iota_Z^Y]{} & H_{comp}^i(Z; A) \\
 & \nwarrow & & \nearrow & \\
 & & \iota_Z^X & &
 \end{array}$$

Something about precomposing with any $H^i(Z, Z \setminus K; A)$ is enough. □

The maps ι_Y^X form a covariant functor from the poset of open subsets of X .

[02.06.2025, Lecture 15]
[04.06.2025, Lecture 16]

5.2 The duality map

For compact oriented manifolds M , Poincaré duality says that

$$[M] \cap _ : H^i(M; \mathbb{Z}) \rightarrow H_{n-i}(M; \mathbb{Z})$$

is an isomorphism.

If M is oriented, but not necessarily compact, this is generalized by the duality map, for which we will write

$$D_M: H_{comp}^i(M; \mathbb{Z}) \rightarrow H_{n-i}(M; \mathbb{Z})$$

This will be an isomorphism.

Construction 5.18. Let (M, μ) be an oriented n -manifold. $\mu = \{\mu_x\}_{x \in M}$ an orientation; earlier we saw: For every compact subset K of M there is a unique class („relative fundamental class“) $\mu_K \in H_n(M, M \setminus K; \mathbb{Z})$ such that

$$r_x^K(\mu_K) = \mu_x$$

for all $x \in K$. So the cap product gives a map

$$\mu_K \cap _ : H^i(M, M \setminus K; \mathbb{Z}) \rightarrow H_{n-i}(M; \mathbb{Z})$$

Claim. for $K \subseteq L$ both compact, the map

$$\text{incl}_*: H_n(M, M \setminus L; \mathbb{Z}) \rightarrow H_n(M, M \setminus K; \mathbb{Z})$$

satisfies $\text{incl}_*(\mu_L) = \mu_K$.

Indeed for all $x \in K$,

$$r_x^K(\text{incl}_*(\mu_L)) = r_x^L(\mu_L) = \mu_x$$

for all $x \in K$, so $\text{incl}_*(\mu_L)$ has the properties that characterize μ_K .

Claim. For $\alpha \in H^i(M; M \setminus K; \mathbb{Z})$ we have

$$\mu_K \cap \alpha = \text{incl}_*(\mu_L) \cap \alpha = \mu_L \cap \text{incl}^*(\alpha)$$

in $H_{n-i}(M; \mathbb{Z})$ using the mixed functoriality of cap.

So the following commutes

$$\begin{array}{ccc} H^i(M, M \setminus K; \mathbb{Z}) & & \\ \downarrow j_L^K = \text{incl}^* & \searrow \mu_K \cap - & \\ H^i(M, M \setminus L; \mathbb{Z}) & \xrightarrow{\mu_L \cap -} & H_{n-i}(M, \mathbb{Z}) \end{array}$$

so there is a unique homomorphism $D_M: H_{comp}^i(M; \mathbb{Z}) \rightarrow H_{n-i}(M; \mathbb{Z})$ such that for all $K \subseteq M$ compact, the following commutes:

$$\begin{array}{ccc} H_{comp}^i(M; \mathbb{Z}) & \longrightarrow & H_{n-i}(M; \mathbb{Z}) \\ \lambda_K \uparrow & \nearrow \mu_K \cap - & \\ H^i(M, M \setminus K; \mathbb{Z}) & & \end{array}$$

This can be thought of to some extent as „capping with a non-existent fundamental class.“

Proposition 5.19. For any orientation of \mathbb{R}^n , all $i \geq 0$, the duality map

$$D_{\mathbb{R}^n}: H_{comp}^i(\mathbb{R}^n; \mathbb{Z}) \rightarrow H_{n-i}(\mathbb{R}^n; \mathbb{Z})$$

is an isomorphism.

Proof. Earlier: $H_{comp}^i(\mathbb{R}^n; \mathbb{Z}) = \begin{cases} \mathbb{Z} & \text{if } i = n \\ 0 & \text{else} \end{cases}$ and $H_{n-i}(\mathbb{R}^n; \mathbb{Z}) = 0$, if $i \neq n$, so D_M is an isomorphism between trivial groups for $i \neq n$.

For $i = n$, let

$$\Psi: H^n(\mathbb{R}^n; \mathbb{R}^n \setminus \{0\}; \mathbb{Z}) \rightarrow \text{Hom}(H_n(\mathbb{R} \mid 0), \mathbb{Z})$$

be the evaluation homomorphism from the universal coefficient theorem. This map is surjective by the universal coefficient theorem. Let $\mu_0 \in H_n(\mathbb{R}^n \mid 0)$ be the chosen local orientation. So there is a class $\alpha \in H^n(\mathbb{R}^n, \mathbb{R}^n \setminus \{0\}; \mathbb{Z})$ such that

$$\Phi(\alpha)(\mu_0) = 1.$$

Since \mathbb{R}^n is path connected, every point $x \in \mathbb{R}^n$ represents the same element $e = [x] \in H_0(\mathbb{R}^n; \mathbb{Z})$ a generator. Let $f \in C^n(\mathcal{S}(\mathbb{R}^n), \mathcal{S}(\mathbb{R}^n \setminus 0); \mathbb{Z})$ be a cochain that represents α , and let

$$\sum a_i \cdot \psi_i \in C_n(\mathcal{S}(\mathbb{R}^n), \mathcal{S}(\mathbb{R}^n \setminus 0); \mathbb{Z})$$

represent μ_0 . with some $\psi_i: \nabla^n \rightarrow \mathbb{R}^n, a_i \in \mathbb{Z}$. Then

$$\begin{aligned}
 \mu_0 \cap \alpha &= [\sum a_i \cdot \psi_i] \cap [f] \\
 &= \sum a_i \cdot [\psi_i \cap f] \\
 &= \sum a_i \cdot f(\psi_i) \cdot [\psi(0, 0, \dots, 0, 1)] \\
 &= (\sum a_i \cdot f(\psi_i)) \cdot e \\
 &= \underbrace{\Psi(\alpha)(\mu_0)}_{=1} \cdot e = e
 \end{aligned}$$

where $e \in H_0(\mathbb{R}, \mathbb{Z})$ is the geometric generator. So $\mu_0 \cap \alpha$ generates $H_0(\mathbb{R}^n; \mathbb{Z})$.

We can conclude:

$$\begin{array}{ccc}
 H^n(\mathbb{R}^n, \mathbb{R}^n \setminus 0; \mathbb{Z}) & \xrightarrow[\cong]{\lambda_{\{0\}}} & H^n_{comp}(\mathbb{R}^n; \mathbb{Z}) \\
 & \searrow \mu_0 \cap - & \downarrow D_{\mathbb{R}^n} \\
 & & H_0(\mathbb{R}^n; \mathbb{Z})
 \end{array}$$

So D_M is a surjection between free abelian groups of rank 1, hence an isomorphism! □

We now proved Poincaré duality for \mathbb{R}^n . This is in itself of course not very exciting, but the first step for the bootstrapping we will do later. For that we will develop a Mayer-Vietoris sequence for compactly supported cohomology.

We now first need to show naturality of the duality map.

We let M be a n -manifold, $\mu = \{\mu_x\}_{x \in X}$ an orientation and U an open subset of M . Then $\{r_U^M(\mu_x)\}_{x \in U}$ is an orientation of U .

Proposition 5.20. Let (M, μ) be an oriented n -manifold and U an open subset of M with induced orientation. Then the following diagram commutes:

$$\begin{array}{ccc}
 H^i_{comp}(M; \mathbb{Z}) & \xrightarrow{D_M} & H_{n-i}(M; \mathbb{Z}) \\
 \iota_U^M \uparrow & & \uparrow \text{incl}_* \\
 H^i_{comp}(U; \mathbb{Z}) & \xrightarrow{D_U} & H_{n-i}(U; \mathbb{Z})
 \end{array}$$

Proof. Let K be a compact subset of U .

Claim. The map

$$\text{incl}_*: H_n(U, U \setminus K; \mathbb{Z}) \rightarrow H_n(M, M \setminus K; \mathbb{Z})$$

takes μ_K^U to μ_K^M .

Indeed for all $x \in K$,

$$r_x^M(\text{incl}_*(\mu_K^U)) = r_x^U(\mu_K^U) = \mu_x^U = \mu_x$$

Since this property characterise μ_K^M , the claim holds.

Consider the diagram

$$\begin{array}{ccccc}
 & & \mu_K^U \cap - & & \\
 & \nearrow & & \searrow & \\
 H^i(U, U \setminus K; \mathbb{Z}) & \xrightarrow{\lambda_K} & H_{comp}^i(U; \mathbb{Z}) & \xrightarrow{D_U} & H_{n-i}(U; \mathbb{Z}) \\
 \uparrow \text{incl}^* \cong & & \downarrow \iota_U^M & & \downarrow \text{incl}_* \\
 H^i(M, M \setminus K; \mathbb{Z}) & \xrightarrow{\lambda_K} & H_{comp}^i(M; \mathbb{Z}) & \xrightarrow{D_M} & H_{n-i}(M; \mathbb{Z}) \\
 & \searrow & & \nearrow & \\
 & & \mu_K^M \cap - & &
 \end{array}$$

where we now the left square commutes but don't know about the right one yet.

Every class in $H_{comp}^i(U; \mathbb{Z})$ is of the form $\lambda_K(\alpha)$ for some compact subset K of U , some $\alpha \in H^i(U, U \setminus K; \mathbb{Z})$. By excision there is a class $\beta \in H^i(M, M \setminus K; \mathbb{Z})$, such that $\text{incl}^*(\beta) = \alpha$. So using the mixed functoriality of \cap and the definition of D_M , we get

$$\begin{aligned}
 \text{incl}_*(D_U(\lambda_K(\alpha))) &\stackrel{\text{def. } D_U}{=} \text{incl}_*(\mu_K^U \cap \alpha) \\
 &= \text{incl}_*(\mu_K^U \cap \text{incl}^*(\beta)) \\
 &= \mu_K^M \cap \beta \\
 &= D_M(\lambda_K(\beta)) \\
 &= D_M(\iota_U^M(\lambda_K(\text{incl}^*(\beta)))) \\
 &= D_M(\iota_U^M(\lambda_K(\alpha)))
 \end{aligned}$$

Since all classes in $H_{comp}^i(U; \mathbb{Z})$ are of the form $\lambda_K(\alpha)$, this proves the proposition. \square

5.2.1 Mayer-Vietoris sequences for compactly supported cohomology

Lemma 5.21: Compact unions

Let X be a locally compact Hausdorff space. Let U, V be open subsets of X , such that $X = U \cup V$. Then every compact subset of X is of the form $K \cup L$ for a compact subset K of U and a compact subset L of V .

Proof. Let C be any compact subset of $X = U \cup V$. Then every point $x \in C$ is contained in U or in V . If $x \in U$, then U is a neighborhood of x in X . Since x is locally compact, there is a compact neighborhood N_x of x in U . If $x \in V$, there is a compact neighborhood N_x of x in V . Since C is compact, it is covered by finitely many $N_{x_1} \cup \dots \cup N_{x_n}$ of these compact neighborhoods, with each N_{x_i} contained in U or V .

Set $\bar{K} = \bigcup N_{x_i}$ such that $N_{x_i} \subseteq U$ and $\bar{L} = \bigcup N_{x_i}$ such that $N_{x_i} \subseteq V$. So \bar{K}, \bar{L} are compact, $\bar{K} \subseteq U, \bar{L} \subseteq V$. So $K = C \cap \bar{K}$ is compact in U and $L = C \cap \bar{L}$ is compact in V . We see $C = K \cup L$. \square

Construction 5.22 (Connecting homomorphism). Let X be a locally compact hausdorff space, $U, V \subseteq X$ open, such that $X = U \cup V$. Let A be any coefficient group, that we drop from the notation. We define

$$\partial: H_{comp}^i(X) \rightarrow H_{comp}^{i+1}(U \cap V)$$

For every compact subset C of X , we define

$$\partial_C: H^i(X, X \setminus C) \rightarrow H_{comp}^{i+1}(U \cap V)$$

as follows. We choose compact subsets K of U and L of V , such that $C = K \cup L$ and define ∂_C as the composite

$$H^i(X, X \setminus C) = H^i(X, X \setminus (C \cup L)) \xrightarrow{\partial} H^{i+1}(X, X \setminus (K \cap L)) \xrightarrow{\text{excision}} H^{i+1}(U \cap V, (U \cap V) \setminus (K \cap L)) \xrightarrow{\lambda_{K \cap L}} H_{comp}^{i+1}(U \cap V)$$

using $X \setminus (K \cap L) = (X \setminus K) \cup (X \setminus L)$ and $(X \setminus K) \cap (X \setminus L) = X \setminus (K \cup L)$

Claim.

1. the definition of ∂_C is independent of the choice of K, L .
2. If $C \subseteq \bar{C} \subseteq X$ with C, \bar{C} compact, then

$$\begin{array}{ccc} H^i(X, X \setminus C) & & \\ \downarrow \iota_{\bar{C}} & \searrow \partial_C & \\ H^i(X, X \setminus \bar{C}) & \xrightarrow{\partial_{\bar{C}}} & H_{comp}^{i+1}(U \cap V) \end{array}$$

Appendix

List of definitions

1.3	Differential graded ring	6
2.2	Simplicial abelian group	12
2.6	Tensor product of simplicial abelian groups	13
2.9	(p,q) -shuffle	14
2.13	Eilenberg-Zilber map	15
2.18	Representable functor	17
2.23	Tensor Product of R -chains	22
2.24	Projective R -modules	23
2.26	Global dimension of rings	25
2.28	Tor of nice rings	25
3.1	Manifold	40
3.12	Local orientation	45
3.16	Orientation	47
3.17	Orientability	47
4.1	Cap-Product	55
5.7	Proper maps	59

List of statements

1.2	fundamental properties of cup product	4
1.6	Properties of the cup-product on homology	6
1.8	Commutativity of the cup-product	7
1.10	\cup_1 -Product	8
2.1	Künneth, simple version	12
2.15	Shuffle maps form a chain map	15
2.16	Eilenberg-Zilber	15
2.17	Yoneda lemma	16
2.21	Acyclic models	18
2.30	Algebraic Künneth theorem	26
2.33	Künneth theorem for homology with field coefficients	28
2.34	Künneth theorem for homology	29
2.39	Algebraic Künneth theorem, cohomology	32
2.42	Künneth theorem in cohomology	33
2.50	Hopf map is not constant	38
3.8	Graßmann Manifolds	43
3.14	Orientation covering	46
3.23	Vanishing homology in high dimensions	49
3.26	Fundamental Classes	52
3.27	Top Homology of connected Manifolds	53
5.1	Poincaré-duality	57
5.21	Compact unions	66

Index

(p,q) -shuffle, 14

Alexander Whitney map, 4

compactly supported, 57

cup-product, 4, 6

differential graded ring, 6

duality map, 63

Eilenberg-Zilber map, 15

exterior cup product, 33

finite type CW-complex, 34

fundamental class, 52

Global dimension, 25

graded ring, 6

Künneth theorem, 12

Local homology, 45

manifold, 40

Orientation, 47

Orientation covering, 46

Projective Module, 23

representable functor, 17

ring, 4

shuffle map, 15

simplicial abelian group, 12

Tensor product of simplicial abelian groups,

13