

An introduction to Anderson localization

Author: Jan Šuntajs
Mentor: prof. Janez Bonča
Comentor: doc. Lev Vidmar

April 9, 2018

University of *Ljubljana*
Faculty of *Mathematics and Physics*



What is it about?



- Conduction in **NON-INTERACTING** systems with **DISORDER**
- Describes the role of **IMPURITIES**
- Completely different than the **Drude** model:

$$\sigma \propto l, \quad l : \text{the mean-free path}$$

What does it predict?

- for some disorder: $\sigma = 0$
- seminal paper by **P. W. Anderson (1958)** [1]
- Nobel prize in **1977**

Probability density profile in ballistic and localized regimes in 1D, $L = 1000$



Why does it (still) matter?

What began in 1958 ...

PHYSICAL REVIEW

VOLUME 109, NUMBER 5

MARCH 1, 1958

Absence of Diffusion in Certain Random Lattices

P. W. ANDERSON

Bell Telephone Laboratories, Murray Hill, New Jersey

(Received October 10, 1957)

Why does it (still) matter?

What began in 1958 ...

PHYSICAL REVIEW

VOLUME 109, NUMBER 5

MARCH 1, 1958

Absence of Diffusion in Certain Random Lattices

P. W. ANDERSON

Bell Telephone Laboratories, Murray Hill, New Jersey

(Received October 10, 1957)

... still remains relevant today

PHYSICAL REVIEW B **96**, 214201 (2017)



Anderson localization transitions with and without random potentials

Trithap Devakul and David A. Huse

Department of Physics, Princeton University, New Jersey 08544, USA

(Received 20 October 2017; published 6 December 2017)

The current “hot topic”

- **Many-body localization (MBL)** - includes **INTERACTIONS**

Published in 2015:

Many-Body Localization and Thermalization in Quantum Statistical Mechanics

Rahul Nandkishore¹ and David A. Huse^{1,2}

¹Princeton Center for Theoretical Science, Princeton University, Princeton, New Jersey 08544; email: rahuln@princeton.edu, huse@princeton.edu

²Department of Physics, Princeton University, Princeton, New Jersey 08544

- not our today's topic

672 citations as of April 2018 acc. to Google Scholar.

Outline

- 1 The basic concepts of the Anderson localization
- 2 Models of disorder
- 3 Numerical simulations
- 4 Conclusion

The basics


- **DISORDER** → (**eigen**)states can localize

Localization:

- A localized state:

$$|\psi(\mathbf{r})| \sim \exp(|\mathbf{r} - \mathbf{r}_0|/\xi)$$

- explains **vanishing** transport



Extended Localized

The important keynotes

- An **interference** phenomenon
- Strong **dimensionality** dependence
- Energy dependence \rightarrow the **mobility edge**

The scaling theory


- scaling of the **conductance** g of a **hypercube** L^d [2]

- **Ohmic** conductor:

$$g = \sigma L^{d-2}$$

- **Localized** regime:

$$g \propto \exp(-L)$$



Transition between ext. and loc. states is only possible in 3D. Taken from [4].


The scaling theory

1D, 2D

localization for any finite disorder

3D


localization for some critical disorder



Transition between ext. and loc. states is only possible in 3D. Taken from [4].

The mobility edge

3D, finite disorder




The models of disorder

- somehow distorting the **ideal crystal**

A generic Hamiltonian

$$H = \frac{\hat{p}^2}{2m} + \sum_{j=1}^N V_j(\mathbf{r} - \mathbf{R}_j)$$

- we consider the **Anderson model**



Ideal crystal, compositional, structural and kinetic disorder. Adapted acc. to [3].



Our model - the **Anderson model**

The Anderson Hamiltonian [1]

$$H = \sum_j \varepsilon_j c_j^\dagger c_j - V \sum_{\text{n.n.}} c_i^\dagger c_j + \text{h.c.}$$

Probability distribution of ε_j

$$p(\varepsilon_j) = \frac{1}{2W} \Theta(W - |\varepsilon_j|)$$




The model in 2D. Taken from [4].

The Anderson model

- Two mobility edges in 3D

3D Anderson



My work - the numerical simulations

- How to extract features of the Anderson localization numerically?
 - implementation in 1D, 2D and 3D
 - calculations were run at the F-1 cluster at IJS
 - Full diagonalization and time evolution calculations
- Two localization criteria:
 - the **inverse participation ratio** (IPR)
 - the **absence of diffusion**

Localization criteria - the IPR

The definition





$$P^{-1} = \sum_{\mathbf{r}} |\psi(\mathbf{r})|^4, \quad \|\psi\| = 1.$$

- sum over the lattice sites

If L large enough


$$P^{-1} \propto \begin{cases} 1/L^d, & \text{extended} \\ \text{const}, & \text{localized} \end{cases}$$

IPR - the results



IPR 3D

Inverse participation ratio for different W , $D = 3$, $L = 22$




Absence of diffusion

Time evolution

$$|\psi, t + dt\rangle = \exp(-i\hat{H} dt) |\psi, t\rangle,$$

$$\hat{R}^2 = \sum_{\mathbf{r}_j} \mathbf{r}_j^2 \hat{n}_{\mathbf{r}_j},$$

$$\beta(t) = \frac{d \log R}{d \log t}$$




$$R(t) = \sqrt{\langle \psi, t | \hat{R}^2 | \psi, t \rangle - \langle \psi, 0 | \hat{R}^2 | \psi, 0 \rangle}$$


Absence of diffusion, 2D

$D = 2, L = 500$

$R(t)$ for different W



$\beta(t)$ in $R(t) \approx At^{\beta(t)} + \dots$



Conclusion

- A first step towards the description of conduction in real systems.
- A nontrivial numerical implementation.
- Needed in understanding the MBL phenomena, the current “hot topic.”

References and sources of images



Anderson, P. (1958). *Absence of Diffusion in Certain Random Lattices*. Physical Review, **109**(5), pp.1492-1505.



Abrahams E., Anderson P. W., Licciardello, D. and Ramakrishnan, T.V. (1979). *Scaling Theory of Localization: Absence of Quantum Diffusion in Two Dimensions*. Phys. Rev. Lett. **42**(10), 673



Kramer, B. and MacKinnon, A. (1993). *Localization: theory and experiment*. Reports on Progress in Physics, **56**(12), pp.1469-1564.



Lagendijk, A., Tiggelen, B. and Wiersma, D. (2009). *Fifty years of Anderson localization*. Physics Today, **62**(8), pp.24-29.