

Diabetic Retinal Diagnosis Report

Generated on 2026-01-03 12:20 UTC

Patient Information

Name:	Test Patient	Age:	45
Sex:	M	Weight:	75.0 kg
Height:	175.0 cm	BMI:	24.49
Insulin:	12.0	Smoker:	False
Alcohol:	Low	Vascular Disease:	False

Diagnosis Findings

Left Eye:	Mild	Right Eye:	Mild
Confidence:	85.5%	Risk Score:	4.0

Condition Overview

Comprehensive ophthalmic examination reveals mild, early stage changes in both the left and right eyes. Our confidence in this diagnosis is 85.5%, with an overall risk score of 4.0, indicating a low-to-moderate risk for rapid progression.

Patient Assessment

- Patient Name: Test Patient, Age: 45, Sex: Male.
- Body Mass Index (BMI) is within the healthy range at 24.49 kg/m², supporting good general health.
- Insulin level noted at 12.0 units; further systemic evaluation may be warranted if this is outside your normal range or accompanied by other metabolic indicators.
- No history of smoking, low alcohol consumption, and absence of vascular disease are positive health indicators.
- Ophthalmic findings: Mild changes observed in both the right eye (OD) and left eye (OS).

Clinical Implications

- Current vision is likely stable or minimally affected; however, these early changes necessitate vigilant monitoring.
- There is a potential for progression of these ophthalmic findings over time, which could eventually impact visual function if not managed.

- Early identification allows for proactive management strategies to stabilize the condition and preserve vision.

Treatment Plan

- **Lifestyle:** Maintain current healthy lifestyle choices, including a balanced diet and regular physical activity. Ensure adequate hydration and manage any systemic health conditions proactively.
- **Monitoring:** Adherence to scheduled follow-up ophthalmic examinations is crucial for tracking disease stability or progression.
- **Medical:** No immediate medical intervention is typically indicated for mild, early-stage ophthalmic changes. Management focuses on risk factor modification and surveillance.
- **Follow-up:** Recommended re-evaluation in 6-12 months, or sooner if new symptoms arise.

Life Impact

At this mild stage, the condition is expected to have minimal to no impact on your daily activities, work, or hobbies. Proactive management and adherence to follow-up care are key to maintaining this minimal impact long-term.

Financial Considerations

Initial costs will primarily involve routine ophthalmic examinations. Future costs may arise if the condition progresses and requires specific treatments, medications, or more frequent diagnostic imaging.

Recovery Projection

For mild ophthalmic changes, the prognosis is generally favorable with consistent monitoring and adherence to lifestyle recommendations. The goal is to stabilize the condition and prevent or significantly delay progression, thereby preserving optimal visual function. Complete reversal of changes is not typically expected, but excellent long-term outcomes are achievable with proactive care.

Recommended Additional Assessments

- Dilated fundus examination to assess retinal periphery and optic nerve head more thoroughly.
- Optical Coherence Tomography (OCT) to evaluate retinal structure for subtle changes.
- Visual field testing if any specific functional concerns arise.
- Referral for a comprehensive metabolic panel or endocrinology consultation if the insulin level warrants further systemic investigation.

Important Notice

This report is based on current findings and medical understanding. It is not exhaustive and should not replace direct medical advice from your healthcare provider. Adherence to recommended follow-up schedules and lifestyle modifications is essential. Any new or worsening symptoms should be reported immediately.