


SURFACES

Rodrigo Silveira


Curve and Surface Design
Facultat d'Informàtica de Barcelona
Universitat Politècnica de Catalunya

A surface




Source: <https://openhome.cc/eGossip/OpenSCAD/BezierSurface.htm>

PARAMETRIZING SURFACES



PARAMETRIZING SURFACES


1) Explicit equation



PARAMETRIZING SURFACES

1) Explicit equation

$z = f(x, y)$, for $x \in [x_0, x_1]$, $y \in [y_0, y_1]$, for f a continuous function




PARAMETRIZING SURFACES

1) Explicit equation

$z = f(x, y)$, for $x \in [x_0, x_1]$, $y \in [y_0, y_1]$, for f a continuous function

The surface is $S = \{(\mathbf{x}, \mathbf{y}, \mathbf{f}(\mathbf{x}, \mathbf{y}))/x \in [x_0, x_1], y \in [y_0, y_1]\}$




PARAMETRIZING SURFACES


1) Explicit equation

$z = f(x, y)$, for $x \in [x_0, x_1]$, $y \in [y_0, y_1]$, for f a continuous function

The surface is $S = \{(\mathbf{x}, \mathbf{y}, \mathbf{f}(\mathbf{x}, \mathbf{y}))/x \in [x_0, x_1], y \in [y_0, y_1]\}$



[From Wikimedia commons - by Ag2gaeh - Own work]





PARAMETRIZING SURFACES

1) Explicit equation

$z = f(x, y)$, for $x \in [x_0, x_1]$, $y \in [y_0, y_1]$, for f a continuous function

The surface is $S = \{(\mathbf{x}, \mathbf{y}, \mathbf{f}(\mathbf{x}, \mathbf{y}))/x \in [x_0, x_1], y \in [y_0, y_1]\}$

Example: $z = \sin x \sin y$



2) Implicit equation

$F(x, y, z) = 0$, for F a continuous function

2) Implicit equation

$F(x, y, z) = 0$, for F a continuous function

The surface is $S = \{(x, y, z) / F(x, y, z) = 0\}$

2) Implicit equation

$F(x, y, z) = 0$, for F a continuous function

The surface is $S = \{(x, y, z) / F(x, y, z) = 0\}$

Example: $x^2 + y^2 + z^2 = 4$


PARAMETRIZING SURFACES

2) Implicit equation

$F(x, y, z) = 0$, for F a continuous function

The surface is $S = \{(x, y, z) / F(x, y, z) = 0\}$

Example: $x^2 + y^2 + z^2 = 4$



3) Parametric equation

$$\begin{array}{l} x = x(u, v) \\ y = y(u, v) \\ z = z(u, v) \end{array} \left| \begin{array}{l} \longrightarrow u \in [u_0, u_1], v \in [v_0, v_1], x, y, z : \mathbb{R}^2 \rightarrow \mathbb{R} \text{ continuous} \end{array} \right.$$

PARAMETRIZING SURFACES

3) Parametric equation

$$\begin{array}{l} x = x(u, v) \\ y = y(u, v) \\ z = z(u, v) \end{array} \left| \begin{array}{l} \longrightarrow \\ \longrightarrow \\ \longrightarrow \end{array} \right. u \in [u_0, u_1], v \in [v_0, v_1], x, y, z : \mathbb{R}^2 \rightarrow \mathbb{R} \text{ continuous}$$

The surface is $S = \{(\mathbf{x}(\mathbf{u}, \mathbf{v}), \mathbf{y}(\mathbf{u}, \mathbf{v}), \mathbf{z}(\mathbf{u}, \mathbf{v})) / u \in [u_0, u_1], v \in [v_0, v_1]\}$

PARAMETRIZING SURFACES

3) Parametric equation

$$\begin{array}{l} x = x(u, v) \\ y = y(u, v) \\ z = z(u, v) \end{array} \left| \begin{array}{l} \longrightarrow \\ \longrightarrow \\ \longrightarrow \end{array} \right. u \in [u_0, u_1], v \in [v_0, v_1], x, y, z : \mathbb{R}^2 \rightarrow \mathbb{R} \text{ continuous}$$

The surface is $S = \{(\mathbf{x}(\mathbf{u}, \mathbf{v}), \mathbf{y}(\mathbf{u}, \mathbf{v}), \mathbf{z}(\mathbf{u}, \mathbf{v})) / u \in [u_0, u_1], v \in [v_0, v_1]\}$

Example: $(\cos(u), \sin(u), v)$, for $0 \leq u \leq 2\pi$ and $0 \leq v \leq 2$


PARAMETRIZING SURFACES

3) Parametric equation

$$\left. \begin{array}{l} x = x(u, v) \\ y = y(u, v) \\ z = z(u, v) \end{array} \right| \rightarrow u \in [u_0, u_1], v \in [v_0, v_1], x, y, z : \mathbb{R}^2 \rightarrow \mathbb{R} \text{ continuous}$$

The surface is $S = \{(\mathbf{x}(\mathbf{u}, \mathbf{v}), \mathbf{y}(\mathbf{u}, \mathbf{v}), \mathbf{z}(\mathbf{u}, \mathbf{v})) / u \in [u_0, u_1], v \in [v_0, v_1]\}$

Example: $(\cos(u), \sin(u), v)$, for $0 \leq u \leq 2\pi$ and $0 \leq v \leq 2$




PARAMETRIZING SURFACES

3) Parametric equation

$$\left. \begin{array}{l} x = x(u, v) \\ y = y(u, v) \\ z = z(u, v) \end{array} \right\} \rightarrow u \in [u_0, u_1], v \in [v_0, v_1], x, y, z : \mathbb{R}^2 \rightarrow \mathbb{R} \text{ continuous}$$

The surface is $S = \{(\mathbf{x}(\mathbf{u}, \mathbf{v}), \mathbf{y}(\mathbf{u}, \mathbf{v}), \mathbf{z}(\mathbf{u}, \mathbf{v})) / u \in [u_0, u_1], v \in [v_0, v_1]\}$

Example: $(\cos(u), \sin(u), v)$, for $0 \leq u \leq 2\pi$ and $0 \leq v \leq 2$



Example: $(u, u \cos(v), u \sin(v))$, for $u, v \in \mathbb{R}$



PARAMETRIZING SURFACES

3) Parametric equation

$$\left. \begin{array}{l} x = x(u, v) \\ y = y(u, v) \\ z = z(u, v) \end{array} \right\} \rightarrow u \in [u_0, u_1], v \in [v_0, v_1], x, y, z : \mathbb{R}^2 \rightarrow \mathbb{R} \text{ continuous}$$

The surface is $S = \{(\mathbf{x}(\mathbf{u}, \mathbf{v}), \mathbf{y}(\mathbf{u}, \mathbf{v}), \mathbf{z}(\mathbf{u}, \mathbf{v})) / u \in [u_0, u_1], v \in [v_0, v_1]\}$

Example: $(\cos(u), \sin(u), v)$, for $0 \leq u \leq 2\pi$ and $0 \leq v \leq 2$



Example: $(u, u \cos(v), u \sin(v))$, for $u, v \in \mathbb{R}$

PARAMETRIZING SURFACES

Advantages and disadvantages?

PARAMETRIZING SURFACES

Advantages and disadvantages?

- **Explicit equations:** easy to discretize and represent, but cannot express all surfaces

PARAMETRIZING SURFACES

Advantages and disadvantages?

- **Explicit equations:** easy to discretize and represent, but cannot express all surfaces
- **Implicit equations:** allow to represent all surfaces, also proving properties and testing points for containment. However, they are not easy to discretize and to represent graphically

PARAMETRIZING SURFACES

Advantages and disadvantages?

- **Explicit equations:** easy to discretize and represent, but cannot express all surfaces
- **Implicit equations:** allow to represent all surfaces, also proving properties and testing points for containment. However, they are not easy to discretize and to represent graphically
- **Parametric equations:** allow to represent all surfaces, and are easy to discretize. However, algebraic manipulation is not easy

PARAMETRIZING SURFACES

Advantages and disadvantages?

- **Explicit equations:** easy to discretize and represent, but cannot express all surfaces
- **Implicit equations:** allow to represent all surfaces, also proving properties and testing points for containment. However, they are not easy to discretize and to represent graphically
- **Parametric equations:** allow to represent all surfaces, and are easy to discretize. However, algebraic manipulation is not easy



→ the choice for surface design for CAD and graphics

EXAMPLES OF SURFACES

Some surfaces easier to parametrize: curve-based

EXAMPLES OF SURFACES

Some surfaces easier to parametrize: curve-based

1) Ruled surfaces

EXAMPLES OF SURFACES

Some surfaces easier to parametrize: curve-based

1) Ruled surfaces

Surfaces that can be drawn by moving a line in space

→ For every point p on the surface, there exists a line $\ell(x)$ that is contained in the surface

EXAMPLES OF SURFACES

Some surfaces easier to parametrize: curve-based

1) Ruled surfaces

Surfaces that can be drawn by moving a line in space

→ For every point p on the surface, there exists a line $\ell(x)$ that is contained in the surface

Examples:

EXAMPLES OF SURFACES


Some surfaces easier to parametrize: curve-based

1) Ruled surfaces

Surfaces that can be drawn by moving a line in space

→ For every point p on the surface, there exists a line $\ell(x)$ that is contained in the surface

Examples:



Cone

$$(u, u \cos(v), u \sin(v))$$

EXAMPLES OF SURFACES


Some surfaces easier to parametrize: curve-based

1) Ruled surfaces

Surfaces that can be drawn by moving a line in space


→ For every point p on the surface, there exists a line $\ell(x)$ that is contained in the surface

Examples:



Cone

$$(u, u \cos(v), u \sin(v))$$



Hyperboloid

$$(\sqrt{1+u^2} \cos v, \sqrt{1+u^2} \sin v, u)$$

EXAMPLES OF SURFACES

Some surfaces easier to parametrize: curve-based


Source: <http://math.arizona.edu/~models>

1) Ruled surfaces

Surfaces that can be drawn by moving a line in space


→ For every point p on the surface, there exists a line $\ell(x)$ that is contained in the surface

Examples:



Cone

$$(u, u \cos(v), u \sin(v))$$



Hyperboloid

$$(\sqrt{1+u^2} \cos v, \sqrt{1+u^2} \sin v, u)$$




EXAMPLES OF SURFACES

Ruled surface: helicoid

EXAMPLES OF SURFACES

Ruled surface: helicoid




EXAMPLES OF SURFACES

Ruled surface: helicoid

Consider a circular helix with axis $0z$

The *helicoid* associated is the set of all lines perpendicular to $0z$ that go through a point in $0z$ and one in the helix.




EXAMPLES OF SURFACES

Ruled surface: helicoid

Consider a circular helix with axis $0z$

The *helicoid* associated is the set of all lines perpendicular to $0z$ that go through a point in $0z$ and one in the helix.

Parametrization



EXAMPLES OF SURFACES

Ruled surface: helicoid

Consider a circular helix with axis $0z$


The *helicoid* associated is the set of all lines perpendicular to $0z$ that go through a point in $0z$ and one in the helix.

Parametrization

A point on circular helix:

$$P(t) = (a \cos t, a \sin t, bt)$$

A point on the axis $0z$: $Q(s) = (0, 0, s)$



EXAMPLES OF SURFACES

Ruled surface: helicoid

Consider a circular helix with axis $0z$

The *helicoid* associated is the set of all lines perpendicular to $0z$ that go through a point in $0z$ and one in the helix.

Parametrization


A point on circular helix:

$$P(t) = (a \cos t, a \sin t, bt)$$

A point on the axis $0z$: $Q(s) = (0, 0, s)$

Horizontal line through $P(t)$ and $Q(s)$:

To be horizontal, we set $s = bt$



EXAMPLES OF SURFACES

Ruled surface: helicoid

Consider a circular helix with axis $0z$

The *helicoid* associated is the set of all lines perpendicular to $0z$ that go through a point in $0z$ and one in the helix.

Parametrization

A point on circular helix:


$$P(t) = (a \cos t, a \sin t, bt)$$

A point on the axis $0z$: $Q(s) = (0, 0, s)$

Horizontal line through $P(t)$ and $Q(s)$:

To be horizontal, we set $s = bt$

$$\begin{aligned}\text{Thus: } S(t, \lambda) &= (1 - \lambda)Q(t) + \lambda P(t) \\ &= (1 - \lambda)(0, 0, bt) + \lambda(a \cos t, a \sin t, bt) \\ &= (a\lambda \cos t, a\lambda \sin t, bt)\end{aligned}$$



EXAMPLES OF SURFACES

2) Surfaces of revolution


2) Surfaces of revolution

Surface created by rotating a curve (*generatrix* or *profile*) around a line (*axis*)

EXAMPLES OF SURFACES

2) Surfaces of revolution

Surface created by rotating a curve (*generatrix* or *profile*) around a line (*axis*)




$$\gamma(t) = (x(t), y(t), z(t))$$

EXAMPLES OF SURFACES

2) Surfaces of revolution


Surface created by rotating a curve (*generatrix* or *profile*) around a line (*axis*)



EXAMPLES OF SURFACES

2) Surfaces of revolution



Surface created by rotating a curve (*generatrix* or *profile*) around a line (*axis*)



EXAMPLES OF SURFACES

2) Surfaces of revolution

Surface created by rotating a curve (*generatrix* or *profile*) around a line (*axis*)





rotation from 0 to π


EXAMPLES OF SURFACES

2) Surfaces of revolution

Surface created by rotating a curve (*generatrix* or *profile*) around a line (*axis*)



rotation from 0 to π



rotation from 0 to 2π

EXAMPLES OF SURFACES

Surfaces of revolution


Surface created by rotating a curve (*generatrix*)
around a line (*axis*)

EXAMPLES OF SURFACES

Surfaces of revolution

Surface created by rotating a curve (*generatrix*) around a line (*axis*)

Given a parametrization of the generatrix curve, say, in the xy -plane, so $P(t) = (x(t), y(t), 0)$, $t \in [0, 1]$, and an axis, say $0y$, we obtain the parametrization of the surface of revolution around the axis as follows:




EXAMPLES OF SURFACES

Surfaces of revolution

Surface created by rotating a curve (*generatrix*) around a line (*axis*)

Given a parametrization of the generatrix curve, say, in the xy -plane, so $P(t) = (x(t), y(t), 0)$, $t \in [0, 1]$, and an axis, say $0y$, we obtain the parametrization of the surface of revolution around the axis as follows:



EXAMPLES OF SURFACES


Surfaces of revolution

Surface created by rotating a curve (*generatrix*) around a line (*axis*)

Given a parametrization of the generatrix curve, say, in the xy -plane, so $P(t) = (x(t), y(t), 0)$, $t \in [0, 1]$, and an axis, say $0y$, we obtain the parametrization of the surface of revolution around the axis as follows:

$$T_y(w) = \begin{pmatrix} \cos w & 0 & \sin w \\ 0 & 1 & 0 \\ -\sin w & 0 & \cos w \end{pmatrix}$$

rotation by angle w around y -axis



EXAMPLES OF SURFACES

Surfaces of revolution

Surface created by rotating a curve (*generatrix*) around a line (*axis*)

Given a parametrization of the generatrix curve, say, in the xy -plane, so $P(t) = (x(t), y(t), 0)$, $t \in [0, 1]$, and an axis, say $0y$, we obtain the parametrization of the surface of revolution around the axis as follows:


$$T_y(w) = \begin{pmatrix} \cos w & 0 & \sin w \\ 0 & 1 & 0 \\ -\sin w & 0 & \cos w \end{pmatrix}$$

rotation by angle w around y -axis

We apply the rotation as follows

$$P(u)T_y(w)$$

for $u \in [0, 1]$ and $w \in [0, 2\pi]$



EXAMPLES OF SURFACES


Surfaces of revolution

Surface created by rotating a curve (*generatrix*) around a line (*axis*)

Given a parametrization of the generatrix curve, say, in the xy -plane, so $P(t) = (x(t), y(t), 0)$, $t \in [0, 1]$, and an axis, say $0y$, we obtain the parametrization of the surface of revolution around the axis as follows:

$$T_y(w) = \begin{pmatrix} \cos w & 0 & \sin w \\ 0 & 1 & 0 \\ -\sin w & 0 & \cos w \end{pmatrix}$$

rotation by angle w around y -axis



We apply the rotation as follows


$$\begin{aligned} &P(u)T_y(w) \\ &= (x(u) \cos w, y(u), x(u) \sin w) \end{aligned}$$

for $u \in [0, 1]$ and $w \in [0, 2\pi]$

EXAMPLES OF SURFACES

Example of surface of revolution: cone

What is a cone? Let's start from the *implicit equation*




EXAMPLES OF SURFACES

Example of surface of revolution: cone

What is a cone? Let's start from the *implicit equation*

A right circular cone with *apex point* v , *axis* ℓ (a line through v), and *aperture angle* 2α (for $0 < \alpha < \pi/2$) is....





EXAMPLES OF SURFACES

Example of surface of revolution: cone

What is a cone? Let's start from the *implicit equation*

A right circular cone with *apex point* v , *axis* ℓ (a line through v), and *aperture angle* 2α (for $0 < \alpha < \pi/2$) is....

... the set of all lines through v forming angle α with ℓ .



Warning: only half cone shown here

EXAMPLES OF SURFACES



Example of surface of revolution: cone

What is a cone? Let's start from the *implicit equation*

A right circular cone with *apex point* v , *axis* ℓ (a line through v), and *aperture angle* 2α (for $0 < \alpha < \pi/2$) is....

... the set of all lines through v forming angle α with ℓ .

Example: apex is $(0, 0, 0)$, axis is $0z$, an equation is:



Warning: only half cone shown here

EXAMPLES OF SURFACES



Example of surface of revolution: cone

What is a cone? Let's start from the *implicit equation*

A right circular cone with *apex point* v , *axis* ℓ (a line through v), and *aperture angle* 2α (for $0 < \alpha < \pi/2$) is....

... the set of all lines through v forming angle α with ℓ .

Example: apex is $(0, 0, 0)$, axis is $0z$, an equation is:



Warning: only half cone shown here

EXAMPLES OF SURFACES



Example of surface of revolution: cone

What is a cone? Let's start from the *implicit equation*

A right circular cone with *apex point* v , *axis* ℓ (a line through v), and *aperture angle* 2α (for $0 < \alpha < \pi/2$) is....

... the set of all lines through v forming angle α with ℓ .

Example: apex is $(0, 0, 0)$, axis is $0z$, an equation is:



Warning: only half cone shown here

EXAMPLES OF SURFACES

Example of surface of revolution: cone



What is a cone? Let's start from the *implicit equation*

A right circular cone with *apex point* v , *axis* ℓ (a line through v), and *aperture angle* 2α (for $0 < \alpha < \pi/2$) is....

... the set of all lines through v forming angle α with ℓ .

Example: apex is $(0, 0, 0)$, axis is $0z$, an equation is:

$$\tan(\pi/2 - \alpha) = \frac{z}{\sqrt{x^2 + y^2}} \quad (\text{slope of orange segment})$$



Warning: only half cone shown here

EXAMPLES OF SURFACES

Example of surface of revolution: cone

What is a cone? Let's start from the *implicit equation*

A right circular cone with *apex point* v , *axis* ℓ (a line through v), and *aperture angle* 2α (for $0 < \alpha < \pi/2$) is....



... the set of all lines through v forming angle α with ℓ .

Example: apex is $(0, 0, 0)$, axis is $0z$, an equation is:

$$\tan(\pi/2 - \alpha) = \frac{z}{\sqrt{x^2 + y^2}} \quad (\text{slope of orange segment})$$

$$\tan(\pi/2 - \alpha) = \frac{1}{\tan \alpha} \quad (\text{applying some trigonometry})$$

$$\text{Then we have } \frac{1}{\tan \alpha} = \frac{z}{\sqrt{x^2 + y^2}} \Leftrightarrow \tan \alpha = \frac{\sqrt{x^2 + y^2}}{z}$$



Warning: only half cone shown here

EXAMPLES OF SURFACES

Example of surface of revolution: cone

What is a cone? Let's start from the *implicit equation*

A right circular cone with *apex point* v , *axis* ℓ (a line through v), and *aperture angle* 2α (for $0 < \alpha < \pi/2$) is....

... the set of all lines through v forming angle α with ℓ .



Example: apex is $(0, 0, 0)$, axis is $0z$, an equation is:

$$\tan(\pi/2 - \alpha) = \frac{z}{\sqrt{x^2 + y^2}} \quad (\text{slope of orange segment})$$

$$\tan(\pi/2 - \alpha) = \frac{1}{\tan \alpha} \quad (\text{applying some trigonometry})$$

$$\text{Then we have } \frac{1}{\tan \alpha} = \frac{z}{\sqrt{x^2 + y^2}} \Leftrightarrow \tan \alpha = \frac{\sqrt{x^2 + y^2}}{z}$$

$$x^2 + y^2 = z^2 \tan^2 \alpha \Leftrightarrow x^2 + y^2 - z^2 \tan^2 \alpha = 0$$



Warning: only half cone shown here

EXAMPLES OF SURFACES

Example of surface of revolution: cone

What is a cone? Let's start from the *implicit equation*

A right circular cone with *apex point* v , *axis* ℓ (a line through v), and *aperture angle* 2α (for $0 < \alpha < \pi/2$) is....

... the set of all lines through v forming angle α with ℓ .

Example: apex is $(0, 0, 0)$, axis is $0z$, an equation is:



$$\tan(\pi/2 - \alpha) = \frac{z}{\sqrt{x^2 + y^2}} \quad (\text{slope of orange segment})$$

$$\tan(\pi/2 - \alpha) = \frac{1}{\tan \alpha} \quad (\text{applying some trigonometry})$$

$$\text{Then we have } \frac{1}{\tan \alpha} = \frac{z}{\sqrt{x^2 + y^2}} \Leftrightarrow \tan \alpha = \frac{\sqrt{x^2 + y^2}}{z}$$

$$x^2 + y^2 = z^2 \tan^2 \alpha \Leftrightarrow x^2 + y^2 - z^2 \tan^2 \alpha = 0$$

implicit equation of the cone




Warning: only half cone shown here

EXAMPLES OF SURFACES

Example of surface of revolution: cone

Now as a surface of revolution...




EXAMPLES OF SURFACES

Example of surface of revolution: cone

Now as a surface of revolution...

Question: what is the generatrix curve?




EXAMPLES OF SURFACES

Example of surface of revolution: cone

Now as a surface of revolution...

Question: what is the generatrix curve?

Line making angle α with $0z$



EXAMPLES OF SURFACES


Example of surface of revolution: cone

Now as a surface of revolution...

Question: what is the generatrix curve?

Line making angle α with $0z$

$P(u) = u\vec{v}$, for $u \in \mathbb{R}$, and some \vec{v} in \mathbb{R}^3



EXAMPLES OF SURFACES

Example of surface of revolution: cone


Now as a surface of revolution...

Question: what is the generatrix curve?

Line making angle α with $0z$

$P(u) = u\vec{v}$, for $u \in \mathbb{R}$, and some \vec{v} in \mathbb{R}^3

$\vec{v} = (1, \text{slope of line})$



EXAMPLES OF SURFACES

Example of surface of revolution: cone


Now as a surface of revolution...

Question: what is the generatrix curve?

Line making angle α with $0z$

$P(u) = u\vec{v}$, for $u \in \mathbb{R}$, and some \vec{v} in \mathbb{R}^3

$\vec{v} = (1, \text{slope of line}) = (1, \tan(\pi/2 - \alpha)) = (1, \frac{1}{\tan \alpha})$



EXAMPLES OF SURFACES

Example of surface of revolution: cone

Now as a surface of revolution...


Question: what is the generatrix curve?

Line making angle α with $0z$

$P(u) = u\vec{v}$, for $u \in \mathbb{R}$, and some \vec{v} in \mathbb{R}^3

$\vec{v} = (1, \text{slope of line}) = (1, \tan(\pi/2 - \alpha)) = (1, \frac{1}{\tan \alpha})$

$P(u) = u(1, \frac{1}{\tan \alpha}) = (u, \frac{u}{\tan \alpha})$



EXAMPLES OF SURFACES

Example of surface of revolution: cone

Now as a surface of revolution...

Question: what is the generatrix curve?

Line making angle α with $0z$

$P(u) = u\vec{v}$, for $u \in \mathbb{R}$, and some \vec{v} in \mathbb{R}^3


$\vec{v} = (1, \text{slope of line}) = (1, \tan(\pi/2 - \alpha)) = (1, \frac{1}{\tan \alpha})$

$P(u) = u(1, \frac{1}{\tan \alpha}) = (u, \frac{u}{\tan \alpha})$

Now, apply rotation around z -axis (based on matrix very similar to $T_y(w)$):

$C(u) = P(u)T_z(w) = (x(u) \cos w, x(u) \sin w, y(u))$

for $u \in \mathbb{R}$, $w \in [0, 2\pi]$



EXAMPLES OF SURFACES

Example of surface of revolution: cone

Now as a surface of revolution...

Question: what is the generatrix curve?

Line making angle α with $0z$

$P(u) = u\vec{v}$, for $u \in \mathbb{R}$, and some \vec{v} in \mathbb{R}^3

$\vec{v} = (1, \text{slope of line}) = (1, \tan(\pi/2 - \alpha)) = (1, \frac{1}{\tan \alpha})$


$P(u) = u(1, \frac{1}{\tan \alpha}) = (u, \frac{u}{\tan \alpha})$

Now, apply rotation around z -axis (based on matrix very similar to $T_y(w)$):

$C(u) = P(u)T_z(w) = (x(u) \cos w, x(u) \sin w, y(u))$

$= \left(u \cos w, u \sin w, \frac{u}{\tan \alpha} \right)$ for $u \in \mathbb{R}, w \in [0, 2\pi]$

parametric equation of the cone



LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane

Let S be a surface parametrized as

$S(u, v) = (x(u, v), y(u, v), z(u, v))$ for (u, v) in some domain

LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane

Let S be a surface parametrized as

$S(u, v) = (x(u, v), y(u, v), z(u, v))$ for (u, v) in some domain

If the functions x, y, z have partial derivatives, we consider

$$\begin{cases} \frac{\partial S}{\partial u} = \left(\frac{\partial x}{\partial u}, \frac{\partial y}{\partial u}, \frac{\partial z}{\partial u} \right) \\ \frac{\partial S}{\partial v} = \left(\frac{\partial x}{\partial v}, \frac{\partial y}{\partial v}, \frac{\partial z}{\partial v} \right) \end{cases}$$

LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane

Let S be a surface parametrized as

$S(u, v) = (x(u, v), y(u, v), z(u, v))$ for (u, v) in some domain

If the functions x, y, z have partial derivatives, we consider
$$\begin{cases} \frac{\partial S}{\partial u} = (\frac{\partial x}{\partial u}, \frac{\partial y}{\partial u}, \frac{\partial z}{\partial u}) \\ \frac{\partial S}{\partial v} = (\frac{\partial x}{\partial v}, \frac{\partial y}{\partial v}, \frac{\partial z}{\partial v}) \end{cases}$$

A point P is called *regular* if $\frac{\partial S}{\partial u}(P)$ and $\frac{\partial S}{\partial v}(P)$:

- exist
- are continuous at P
- their cross product is not zero

If one of these conditions does not hold for P , it is called a *singular* point

A surface is called *regular* if all its points are regular

LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane

Let S be a surface parametrized as

$S(u, v) = (x(u, v), y(u, v), z(u, v))$ for (u, v) in some domain

If the functions x, y, z have partial derivatives, we consider
$$\begin{cases} \frac{\partial S}{\partial u} = (\frac{\partial x}{\partial u}, \frac{\partial y}{\partial u}, \frac{\partial z}{\partial u}) \\ \frac{\partial S}{\partial v} = (\frac{\partial x}{\partial v}, \frac{\partial y}{\partial v}, \frac{\partial z}{\partial v}) \end{cases}$$

A point P is called *regular* if $\frac{\partial S}{\partial u}(P)$ and $\frac{\partial S}{\partial v}(P)$:

- exist
- are continuous at P
- their cross product is not zero

If one of these conditions does not hold for P , it is called a *singular* point

A surface is called *regular* if all its points are regular

Normal vector

LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane

Let S be a surface parametrized as

$S(u, v) = (x(u, v), y(u, v), z(u, v))$ for (u, v) in some domain

If the functions x, y, z have partial derivatives, we consider
$$\begin{cases} \frac{\partial S}{\partial u} = (\frac{\partial x}{\partial u}, \frac{\partial y}{\partial u}, \frac{\partial z}{\partial u}) \\ \frac{\partial S}{\partial v} = (\frac{\partial x}{\partial v}, \frac{\partial y}{\partial v}, \frac{\partial z}{\partial v}) \end{cases}$$

A point P is called *regular* if $\frac{\partial S}{\partial u}(P)$ and $\frac{\partial S}{\partial v}(P)$:

- exist
- are continuous at P
- their cross product is not zero

If one of these conditions does not hold for P , it is called a *singular* point

A surface is called *regular* if all its points are regular

Normal vector

If P is regular, then $\vec{N} = \frac{\partial S}{\partial u}(P) \times \frac{\partial S}{\partial v}(P)$ (normal vector to S at P)

LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane

Let S be a surface parametrized as

$S(u, v) = (x(u, v), y(u, v), z(u, v))$ for (u, v) in some domain

If the functions x, y, z have partial derivatives, we consider
$$\begin{cases} \frac{\partial S}{\partial u} = (\frac{\partial x}{\partial u}, \frac{\partial y}{\partial u}, \frac{\partial z}{\partial u}) \\ \frac{\partial S}{\partial v} = (\frac{\partial x}{\partial v}, \frac{\partial y}{\partial v}, \frac{\partial z}{\partial v}) \end{cases}$$

A point P is called *regular* if $\frac{\partial S}{\partial u}(P)$ and $\frac{\partial S}{\partial v}(P)$:

- exist
- are continuous at P
- their cross product is not zero

If one of these conditions does not hold for P , it is called a *singular* point

A surface is called *regular* if all its points are regular

Normal vector

If P is regular, then $\vec{N} = \frac{\partial S}{\partial u}(P) \times \frac{\partial S}{\partial v}(P)$ (normal vector to S at P)

Tangent plane

The plane tangent to S at P is given by the plane defined by P and \vec{N}

LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane

$\frac{\partial S}{\partial u}(P)$ and $\frac{\partial S}{\partial v}(P)$ are tangent vectors in the u and v directions




Figure from oreilley.com

LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane

$\frac{\partial S}{\partial u}(P)$ and $\frac{\partial S}{\partial v}(P)$ are tangent vectors in the u and v directions

If nothing “goes wrong”, they define the tangent plane at P




Figure from oreilley.com

LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane

$\frac{\partial S}{\partial u}(P)$ and $\frac{\partial S}{\partial v}(P)$ are tangent vectors in the u and v directions

If nothing “goes wrong”, they define the tangent plane at P

Exceptions:

- Surface does not have a tangent plane at P
- Parametrization is irregular at P





Figure from oreilley.com

LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane: example

The apex of a cone is a **singular point**




LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane: example

The apex of a cone is a **singular point**

Consider the cone $x^2 + y^2 = z^2$, which can be parametrized as:

$(u \cos v, u \sin v, \frac{u}{\tan \alpha})$, for $\alpha = \pi/4$ ($\tan \pi/4 = 1$) so we have,



LOCAL PROPERTIES OF SURFACES


Normal vector, tangent plane: example

The apex of a cone is a **singular point**

Consider the cone $x^2 + y^2 = z^2$, which can be parametrized as:

$(u \cos v, u \sin v, \frac{u}{\tan \alpha})$, for $\alpha = \pi/4$ ($\tan \pi/4 = 1$) so we have,

$S(u, v) = (u \cos v, u \sin v, u)$, for $u \in \mathbb{R}$ and $v \in [0, 2\pi)$



LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane: example

The apex of a cone is a **singular point**

Consider the cone $x^2 + y^2 = z^2$, which can be parametrized as:


$(u \cos v, u \sin v, \frac{u}{\tan \alpha})$, for $\alpha = \pi/4$ ($\tan \pi/4 = 1$) so we have,

$S(u, v) = (u \cos v, u \sin v, u)$, for $u \in \mathbb{R}$ and $v \in [0, 2\pi)$

The tangent vectors are as follows:

$$\frac{\partial S}{\partial u} = (\cos v, \sin v, 1)$$

$$\frac{\partial S}{\partial v} = (-u \sin v, u \cos v, 0)$$



LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane: example

The apex of a cone is a **singular point**

Consider the cone $x^2 + y^2 = z^2$, which can be parametrized as:

$(u \cos v, u \sin v, \frac{u}{\tan \alpha})$, for $\alpha = \pi/4$ ($\tan \pi/4 = 1$) so we have,


$S(u, v) = (u \cos v, u \sin v, u)$, for $u \in \mathbb{R}$ and $v \in [0, 2\pi)$

The tangent vectors are as follows:

$$\frac{\partial S}{\partial u} = (\cos v, \sin v, 1)$$

$$\frac{\partial S}{\partial v} = (-u \sin v, u \cos v, 0)$$

$$\vec{N}(u, v) = \frac{\partial S}{\partial u}(u, v) \times \frac{\partial S}{\partial v}(u, v) = (-u \cos v, -u \sin v, u)$$



LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane: example

The apex of a cone is a **singular point**

Consider the cone $x^2 + y^2 = z^2$, which can be parametrized as:

$(u \cos v, u \sin v, \frac{u}{\tan \alpha})$, for $\alpha = \pi/4$ ($\tan \pi/4 = 1$) so we have,

$S(u, v) = (u \cos v, u \sin v, u)$, for $u \in \mathbb{R}$ and $v \in [0, 2\pi)$

The tangent vectors are as follows:


$$\frac{\partial S}{\partial u} = (\cos v, \sin v, 1)$$

$$\frac{\partial S}{\partial v} = (-u \sin v, u \cos v, 0)$$

$$\left. \begin{array}{l} \frac{\partial S}{\partial u} = (\cos v, \sin v, 1) \\ \frac{\partial S}{\partial v} = (-u \sin v, u \cos v, 0) \end{array} \right\} \vec{N}(u, v) = \frac{\partial S}{\partial u}(u, v) \times \frac{\partial S}{\partial v}(u, v) = (-u \cos v, -u \sin v, u)$$

For the apex, $P = (0, 0, 0)$, we get $\vec{N}(0, 0) = 0$

(thus, the apex is a singular point)



LOCAL PROPERTIES OF SURFACES


Normal vector, tangent plane: example

Consider the north hemisphere of the unit sphere

LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane: example

Consider the north hemisphere of the unit sphere




LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane: example

Consider the north hemisphere of the unit sphere

Parametrization 1

$$S(u, v) = (u, v, \sqrt{1 - u^2 - v^2})$$



LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane: example


Consider the north hemisphere of the unit sphere

Parametrization 1

$$S(u, v) = (u, v, \sqrt{1 - u^2 - v^2})$$

$$\frac{\partial S}{\partial u} = \left(1, 0, \frac{-u}{\sqrt{1 - u^2 - v^2}}\right)$$

$$\frac{\partial S}{\partial v} = \left(0, 1, \frac{-v}{\sqrt{1 - u^2 - v^2}}\right)$$



LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane: example

Consider the north hemisphere of the unit sphere


Parametrization 1

$$S(u, v) = (u, v, \sqrt{1 - u^2 - v^2})$$

$$\frac{\partial S}{\partial u} = (1, 0, \frac{-u}{\sqrt{1 - u^2 - v^2}})$$

$$\frac{\partial S}{\partial v} = (0, 1, \frac{-v}{\sqrt{1 - u^2 - v^2}})$$

$$\vec{N}(u, v) = \frac{\partial S}{\partial u}(u, v) \times \frac{\partial S}{\partial v}(u, v) = (\frac{u}{\sqrt{1 - u^2 - v^2}}, \frac{v}{\sqrt{1 - u^2 - v^2}}, 1)$$



LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane: example

Consider the north hemisphere of the unit sphere

Parametrization 1


$$S(u, v) = (u, v, \sqrt{1 - u^2 - v^2})$$

$$\frac{\partial S}{\partial u} = (1, 0, \frac{-u}{\sqrt{1 - u^2 - v^2}})$$

$$\frac{\partial S}{\partial v} = (0, 1, \frac{-v}{\sqrt{1 - u^2 - v^2}})$$

$$\vec{N}(u, v) = \frac{\partial S}{\partial u}(u, v) \times \frac{\partial S}{\partial v}(u, v) = (\frac{u}{\sqrt{1 - u^2 - v^2}}, \frac{v}{\sqrt{1 - u^2 - v^2}}, 1)$$

For the north pole, $u = v = 0$, we have $\vec{N} = (0, 0, 1)$ (**regular point**)



LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane: example

Consider the north hemisphere of the unit sphere

Parametrization 1

$$S(u, v) = (u, v, \sqrt{1 - u^2 - v^2})$$


$$\frac{\partial S}{\partial u} = (1, 0, \frac{-u}{\sqrt{1 - u^2 - v^2}})$$

$$\frac{\partial S}{\partial v} = (0, 1, \frac{-v}{\sqrt{1 - u^2 - v^2}})$$

$$\vec{N}(u, v) = \frac{\partial S}{\partial u}(u, v) \times \frac{\partial S}{\partial v}(u, v) = (\frac{u}{\sqrt{1 - u^2 - v^2}}, \frac{v}{\sqrt{1 - u^2 - v^2}}, 1)$$

For the north pole, $u = v = 0$, we have $\vec{N} = (0, 0, 1)$ (**regular point**)

Parametrization 2



LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane: example

Consider the north hemisphere of the unit sphere

Parametrization 1

$$S(u, v) = (u, v, \sqrt{1 - u^2 - v^2})$$

$$\frac{\partial S}{\partial u} = (1, 0, \frac{-u}{\sqrt{1 - u^2 - v^2}})$$

$$\frac{\partial S}{\partial v} = (0, 1, \frac{-v}{\sqrt{1 - u^2 - v^2}})$$


$$\vec{N}(u, v) = \frac{\partial S}{\partial u}(u, v) \times \frac{\partial S}{\partial v}(u, v) = (\frac{u}{\sqrt{1 - u^2 - v^2}}, \frac{v}{\sqrt{1 - u^2 - v^2}}, 1)$$

For the north pole, $u = v = 0$, we have $\vec{N} = (0, 0, 1)$ (**regular point**)

Parametrization 2

$$S(\theta, \varphi) = (\cos \varphi \cos \theta, \cos \varphi \sin \theta, \sin \varphi)$$

$$\text{for } \theta \in [0, 2\pi) \text{ and } \varphi \in [0, \pi/2)$$



LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane: example

Consider the north hemisphere of the unit sphere

Parametrization 1

$$S(u, v) = (u, v, \sqrt{1 - u^2 - v^2})$$

$$\frac{\partial S}{\partial u} = (1, 0, \frac{-u}{\sqrt{1-u^2-v^2}})$$

$$\frac{\partial S}{\partial v} = (0, 1, \frac{-v}{\sqrt{1-u^2-v^2}})$$

$$\vec{N}(u, v) = \frac{\partial S}{\partial u}(u, v) \times \frac{\partial S}{\partial v}(u, v) = (\frac{u}{\sqrt{1-u^2-v^2}}, \frac{v}{\sqrt{1-u^2-v^2}}, 1)$$

For the north pole, $u = v = 0$, we have $\vec{N} = (0, 0, 1)$ (**regular point**)


Parametrization 2

$$S(\theta, \varphi) = (\cos \varphi \cos \theta, \cos \varphi \sin \theta, \sin \varphi) \quad \text{for } \theta \in [0, 2\pi) \text{ and } \varphi \in [0, \pi/2)$$

$$\frac{\partial S}{\partial \theta} = (-\cos \varphi \sin \theta, \cos \varphi \cos \theta, 0)$$

$$\frac{\partial S}{\partial \varphi} = (-\sin \varphi \cos \theta, -\sin \varphi \sin \theta, \cos \varphi)$$

$$\vec{N}(\theta, \varphi) = \cos \varphi (\cos \varphi \cos \theta, \cos \varphi \sin \theta, \sin \varphi)$$



LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane: example

Consider the north hemisphere of the unit sphere

Parametrization 1

$$S(u, v) = (u, v, \sqrt{1 - u^2 - v^2})$$

$$\frac{\partial S}{\partial u} = (1, 0, \frac{-u}{\sqrt{1-u^2-v^2}})$$

$$\frac{\partial S}{\partial v} = (0, 1, \frac{-v}{\sqrt{1-u^2-v^2}})$$

$$\vec{N}(u, v) = \frac{\partial S}{\partial u}(u, v) \times \frac{\partial S}{\partial v}(u, v) = (\frac{u}{\sqrt{1-u^2-v^2}}, \frac{v}{\sqrt{1-u^2-v^2}}, 1)$$

For the north pole, $u = v = 0$, we have $\vec{N} = (0, 0, 1)$ (**regular point**)

Parametrization 2


$$S(\theta, \varphi) = (\cos \varphi \cos \theta, \cos \varphi \sin \theta, \sin \varphi) \quad \text{for } \theta \in [0, 2\pi) \text{ and } \varphi \in [0, \pi/2)$$

$$\frac{\partial S}{\partial \theta} = (-\cos \varphi \sin \theta, \cos \varphi \cos \theta, 0)$$

$$\frac{\partial S}{\partial \varphi} = (-\sin \varphi \cos \theta, -\sin \varphi \sin \theta, \cos \varphi)$$

$$\vec{N}(\theta, \varphi) = \cos \varphi (\cos \varphi \cos \theta, \cos \varphi \sin \theta, \sin \varphi)$$

north pole: $\varphi = \pi/2$ (any θ) $\Rightarrow N(\theta, \pi/2) = (0, 0, 0)$
singular point



LOCAL PROPERTIES OF SURFACES

Normal vector, tangent plane: example

Consider the north hemisphere of the unit sphere

Parametrization 1


$$S(u, v) = (u, v, \sqrt{1 - u^2 - v^2})$$

$$\frac{\partial S}{\partial u} = (1, 0, \frac{-u}{\sqrt{1-u^2-v^2}})$$

$$\frac{\partial S}{\partial v} = (0, 1, \frac{-v}{\sqrt{1-u^2-v^2}})$$

$$\vec{N}(u, v) = \frac{\partial S}{\partial u}(u, v) \times \frac{\partial S}{\partial v}(u, v) = (\frac{u}{\sqrt{1-u^2-v^2}}, \frac{v}{\sqrt{1-u^2-v^2}}, 1)$$

For the north pole, $u = v = 0$, we have $\vec{N} = (0, 0, 1)$ (**regular point**)



Parametrization 2

$$S(\theta, \varphi) = (\cos \varphi \cos \theta, \cos \varphi \sin \theta, \sin \varphi) \quad \text{for } \theta \in [0, 2\pi) \text{ and } \varphi \in [0, \pi/2)$$

$$\frac{\partial S}{\partial \theta} = (-\cos \varphi \sin \theta, \cos \varphi \cos \theta, 0)$$

$$\frac{\partial S}{\partial \varphi} = (-\sin \varphi \cos \theta, -\sin \varphi \sin \theta, \cos \varphi)$$

$$\vec{N}(\theta, \varphi) = \cos \varphi (\cos \varphi \cos \theta, \cos \varphi \sin \theta, \sin \varphi)$$

north pole: $\varphi = \pi/2$ (any θ) $\Rightarrow N(\theta, \pi/2) = (0, 0, 0)$
singular point

→ no issue with surface, but with parametrization