


Historical and Projected Water Balance Report for BRCA

Janelle Christensen, Mike Tercek, David Thoma, John Gross

8/3/2020

Water balance is the mathematical accounting of water input and movement through the environment. It accounts for water storage in different phases (ice, liquid) and loss to the atmosphere (gas) via evapotranspiration. Water balance integrates the interactions of temperature and precipitation to estimate water availability, movement to streams and ground water and use by plants. Water balance accounts for heat load on different aspects and water holding capacity of different soil types. Water balance represents the modification of regional weather and climate to represent the biophysical conditions at local scales. For these reasons, water balance is usually more strongly correlated with natural resource response (plant growth, stream flow, fire danger) than temperature or precipitation.



Modified from McCabe and Markstrom 2007

This report is created from temperature and precipitation data that is input to a water balance model. The historical temperature and precipitation data are from gridMET (<http://www.climatologylab.org/gridmet.html>). The future climate projections, in mcm4 RCP 4.5 and MIROC5 RCP 8.5, are from the CMIP5 experiments (<https://www.wcrp-climate.org/wgcm-cmip/wgcm-cmip5>). The water balance model is maintained by the National Park Service (Mike Tercek, David Thoma and John Gross). Collectively this report summarizes historical and projected biophysical environmental conditions from 1980-2100 at the single grid cell level (gridMET = 4km; projections = 1km). A link to the NPS water balance gridded products can be found here (<http://www.yellowstone.solutions/thredds/catalog.html>).

Selection of the Models

The models used in this analysis were chosen based upon which two climate futures best bracket the range of possibilities. For this region, the model selection was based off of the latitude and longitude for each park centroid, plotted on the scatterplot visualization of future projections from the MACA website. (https://climate.northwestknowledge.net/MACA/vis_scatterplot.php). Models were selected to represent a “drier and hotter” climate future (the bottom right of the scatterplot) and a “wetter and hotter” climate future (the upper left of the scatterplot).

Annual values through time

The variables of runoff, rain, potential evapotranspiration (PET), actual evapotranspiration (AET), and deficit are summed to show annual values through time. Accumulated snow water equivalent (SWE) shows the max value for the year regardless of the month it occurs in. Accumulated growing degree days (AGDD) starts accumulating on October 1 of the previous year, and the max value is taken at its highest point. Soil water shows the mean value for the year.

The reason for the different summary statistics is because hydrologist use these variables in different ways. Water variables that don't have an upper limit like rain, runoff, evapotranspiration and deficit can accumulate indefinitely within a calendar year, whereas soil moisture is limited by the soil water holding capacity so it is averaged on an annual basis. Temperature variables are averaged by convention and growing degree days are summed to account for the seasonal accumulation of heat that controls biological metabolism and thus growth rates and phenology when water is available. For the variables that are accumulated, we take the max of the variable because it is already a summation of the total through the year.




Figure 1. Annual averages over time. The gray line represents historical data modeled from gridMET climate data, the turquoise line represents inmcm4 rcp45 and the yellow line represents MIROC5 rcp85

Deficit v. Actual Evapotranspiration (AET)

Deficit (D) and actual evapotranspiration (AET) represent unmet water needs of vegetation and vegetation water use. They are good predictors of vegetation type, and are useful variables in understanding climate stress on vegetation and how vegetation may transition in the future (Stephenson, 1998). If the clustering of AET and deficit change in climate space, vegetation assemblages will respond and changes in assemblages will eventually reflect

the new climate space. The figure below (from Stephenson) shows how AET v. D differs across bioregions in the U.S.




Figure 2. Stephenson (1998)






Figure 3. Modeled Deficit vs. AET. The points are annual sums of daily observations in each year. The green points represent 1980-2019; the yellow points 2020-2059; and the turquoise points 2060-2099. The contours around each set of points encircle the climate space for inmcm4 rcp45 and MIROC5 rcp85.

Soil Water


Soil Water Seasonality inmcm4
RCP 4.5




Annual Soil Water inmcm4 RCP
4.5




Soil Water inmcm4 RCP 4.5




Soil Water Seasonality MIROC5
RCP 8.5



Annual Soil Water MIROC5 RCP
8.5




Soil Water MIROC5 RCP 8.5




Runoff


Runoff Seasonality inmcm4 RCP 4.5




Annual Runoff inmcm4 RCP 4.5




Runoff inmcm4 RCP 4.5




Runoff Seasonality MIROC5 RCP 8.5



Annual Runoff MIROC5 RCP 8.5




Runoff MIROC5 RCP 8.5




Rain


Rain Seasonality inmcm4 RCP 4.5




Annual Rain inmcm4 RCP 4.5




Rain inmcm4 RCP 4.5




Rain Seasonality MIROC5 RCP 8.5



Annual Rain MIROC5 RCP 8.5




Rain MIROC5 RCP 8.5




Accumulated Snow Water Equivalent (SWE)

Accumulated SWE Seasonality
inmcm4 RCP 4.5




Annual Accumulated SWE inmcm4
RCP 4.5

— gridMET — inmcm4 RCP 4.5




Accumulated SWE inmcm4 RCP 4.5

● 1980-2019 ● 2020-2059 ● 2060-2099




Accumulated SWE Seasonality
MIROC5 RCP 8.5




Annual Accumulated SWE MIROC5
RCP 8.5

— gridMET — MIROC5 RCP 8.5




Accumulated SWE MIROC5 RCP 8.5

● 1980-2019 ● 2020-2059 ● 2060-2099




Potential Evapotranspiration (PET)

Potential Evapotranspiration
Seasonality inmcm4 RCP 4.5




Annual PET inmcm4 RCP 4.5

— gridMET — inmcm4 RCP 4.5




Potential Evapotranspiration
Seasonality MIROC5 RCP 8.5




Annual PET MIROC5 RCP 8.5

— gridMET — MIROC5 RCP 8.5




Actual Evapotranspiration (AET)


Actual Evapotranspiration
Seasonality inmcm4 RCP 4.5




Actual Evapotranspiration
Seasonality MIROC5 RCP 8.5




Annual AET inmcm4 RCP 4.5




Annual AET MIROC5 RCP 8.5



AET inmcm4 RCP 4.5




AET MIROC5 RCP 8.5




Deficit


Deficit Seasonality inmcm4 RCP 4.5




Annual Deficit inmcm4 RCP 4.5




Deficit inmcm4 RCP 4.5




Deficit Seasonality MIROC5 RCP 8.5



Annual Deficit MIROC5 RCP 8.5



Deficit MIROC5 RCP 8.5



Works Cited

Stephenson, N. (1998). Actual evapotranspiration and deficit: biologically meaningful correlates of vegetation distribution across spatial scales. *Journal of biogeography*, 25(5), 855-870.