


This report is created from temperature and precipitation data that is input to a water balance model. The historical temperature and precipitation data are from gridMET (<http://www.climatologylab.org/gridmet.html>). The future climate projections, MRI-CGCM3 RCP 4.5 and NorESM1-M RCP 8.5, used in this report are from the CMIP5 experiments (<https://www.wcrp-climate.org/wgcm-cmip/wgcm-cmip5>) .

The models used in this analysis were chosen based on two climate futures that bracket a wide range of plausible future climate conditions represented by the Global Circulation Models (GCM) in the CMIP5 experiments. The experiments include the effects of anthropogenic warming caused by changes in carbon dioxide concentration in the atmosphere. Representative Concentration Pathways (RCP) describe scenarios of carbon dioxide emission, where RCP 8.5 is the high-emission scenario and RCP 4.5 is a lower emission scenario. Model selection for this report is based on a temperature v. precipitation scatterplot of future climate models and scenarios at mid-century. Users are cautioned that the two models presented in this report are intended to represent a range of conditions, but other projections are available and may need to be considered for a full accounting of possible futures. The spatial resolution of climate data used in the water balance model may not be representative of site-specific conditions, especially in areas of high relief or high topographic complexity, and high variation in soil water holding capacity.

Changes in climate means centered on 2040 relative to historical period (1950-2000) by GCM



Change in annual values through time

Runoff, rain, potential evapotranspiration (PET), actual evapotranspiration (AET), and deficit are summed to show annual values through time. Peak snow water equivalent (SWE) is the max value for the year regardless of the month in which it occurs. Soil water is the mean annual water content in the top meter soil profile. Additionally, it is important to note that precipitation includes rain plus snow, whereas rain is only liquid precipitation. Rain is specifically precipitation that does not fall as snow. If total precipitation stayed the same but temperatures increased, the data here would display that as an increase in rain.

The reason for the different summary statistics is because water variables that don't have an upper limit like rain, runoff, evapotranspiration and deficit can accumulate indefinitely within a calendar year, whereas soil moisture is limited by the soil water holding capacity, so it is averaged on an annual basis. Temperature variables are averaged by convention and growing degree days are summed to account for the seasonal accumulation of heat that controls biological metabolism and thus growth rates and phenology when water is available. Snow water equivalent is reported as a maximum value because snow accumulates during winter and the maximum water content of the snowpack is useful for predicting stream flow rate and reservoir storage needs.





Figure 1. Annual values over time. The gray line represents historical data modeled from gridMET, the turquoise line represents MRI-CGCM3 rcp45 and the yellow line represents NorESM1-M rcp85.

Deficit v. Actual Evapotranspiration

Actual evapotranspiration (AET) is an estimate of soil water used by plants plus evaporation. Deficit (D) is an estimate of unmet water need. That is, it represents the amount of water vegetation would use if the water was available and hence is an estimate of vegetation drought stress. Together, AET and D are good predictors of vegetation condition and are useful variables for understanding climate stress on vegetation and how vegetation may transition in the future (Stephenson, 1998). If the clustering of AET and D change in climate space, vegetation assemblages will respond and changes in assemblages will eventually reflect the new climate space. The figure below (Tercek et al., 2021) shows how AET v. D, calculated using the water balance model used for this report, differs across bioregions in the U.S.



Western CONUS

Figure 2. Ecoregions in the Western United states based on AET and Deficit (CWD). (Tercek et al., 2021)
















Figure 3. Modeled Deficit vs. AET. The points are annual sums of daily observations in each year. The green points represent 1980-2019; the yellow points 2020-2059; and the turquoise points 2060-2099. The contours around each set of points delineate shifts in climate space over time.

Water Balance Projections







Soil Water









Runoff









Rain




Peak Snow Water Equivalent (SWE)



Potential Evapotranspiration (PET)




Actual Evapotranspiration (AET)




Deficit


Deficit Seasonality
MRI-CGCM3 RCP 4.5




Deficit Seasonality
NorESM1-M RCP 8.5




AnnualDeficit
MRI-CGCM3 RCP 4.5




Annual Deficit
NorESM1-M RCP 8.5



Annual Deficit
MRI-CGCM3 RCP 4.5



Annual Deficit
NorESM1-M RCP 8.5



Works Cited

Stephenson, N. (1998). Actual evapotranspiration and deficit: biologically meaningful correlates of vegetation distribution across spatial scales. *Journal of biogeography*, 25(5), 855-870.

Tercek, M., Thoma, D., Sherrill, K., Kagone, S., Senay, G., Gross, J. 2021 Historical changes in plant water use and need in the continental United States. In review.