

INLA Introduction

King Abdullah University of
Science and Technology



May 2024



Outline

1 Introduction to INLA

2 Posterior inference with INLA

3 Examples

4 Discussion



BayesComp group at KAUST





Bayesian inference

Data \mathbf{y} (with covariates \mathbf{Z}), depend on \mathbf{X} and $\boldsymbol{\theta}$ such that, $E[Y] = h(\mathbf{A}(\mathbf{Z})\mathbf{X})$.

Bayes' theorem:

$$\begin{aligned} q(\mathbf{X}, \boldsymbol{\theta}) &\propto L(\mathbf{y}|\mathbf{X}, \boldsymbol{\theta})p(\mathbf{X}, \boldsymbol{\theta}) \\ \text{Posterior} &\propto \text{Likelihood} \times \text{Prior} \end{aligned}$$



Computational aspects

- Analytical methods - conjugacy (pre-computer era)
- Approximate methods - Laplace (can be inaccurate)
- Exact methods - MCMC (very slow for complex models or large data)

Now, due to computing resources approximate methods are gaining popularity - INLA, VB, EP etc.

INLA - 2009 [Rue et al., 2009]

2021+ [Van Niekerk et al., 2023]

HPC [Gaedke-Merzhäuser et al., 2023]



Model definition - GAMM

Suppose we have response data $\mathbf{y}_{n \times 1}$ (conditionally independent) with density function $\pi(y|\mathbf{X}, \boldsymbol{\theta})$ and link function $h(\cdot)$, that is linked to some covariates \mathbf{Z} through linear predictors

$$\boldsymbol{\eta}_n = \beta_0 + \mathbf{Z}_\beta \boldsymbol{\beta} + \sum f^k(\mathbf{Z}_f) = \mathbf{A}\mathbf{X}$$

The inferential aim is to estimate the latent field $\mathbf{X}_m = \{\beta_0, \boldsymbol{\beta}, \mathbf{f}\}$, and $\boldsymbol{\theta}$.



GAMM → LGM

Assume

$$\boldsymbol{X}|\boldsymbol{\theta} \sim N(\boldsymbol{0}, \boldsymbol{Q}(\boldsymbol{\theta})^{-1})$$

where $\boldsymbol{Q}(\boldsymbol{\theta})$ is a sparse matrix (\boldsymbol{X} is a GMRF).

$p(\boldsymbol{X}, \boldsymbol{\theta}) = p(\boldsymbol{X}|\boldsymbol{\theta})p(\boldsymbol{\theta})$ and $p(\boldsymbol{\theta})$ can be non-Gaussian.



Why is INLA so accurate and so fast?

- LGM structure
- Sparse precision matrix
- Specialized matrix algebra for sparse matrices
- NEW: VB (low-rank) correction [van Niekerk and Rue, 2024]

Use precision matrix instead of covariance matrix → natural occurrence



How common are sparse $Q(\theta)$?

Consider an AR(1) model..



AR(1) example



AR(1) example



Posterior approximations by INLA

$$\begin{aligned}\pi(\boldsymbol{X}, \boldsymbol{\theta}, \boldsymbol{y}) &= \pi(\boldsymbol{\theta})\pi(\boldsymbol{X}|\boldsymbol{\theta}) \prod_{i=1}^n \pi(y_i | (\boldsymbol{AX})_i, \boldsymbol{\theta}) \\ \tilde{\pi}(\boldsymbol{\theta}|\boldsymbol{y}) &\propto \frac{\pi(\boldsymbol{X}, \boldsymbol{\theta}, \boldsymbol{y})}{\pi_G(\boldsymbol{X}|\boldsymbol{\theta}, \boldsymbol{y})} \Big|_{\boldsymbol{X}=\mu(\boldsymbol{\theta})} \\ \tilde{\pi}(\theta_j|\boldsymbol{y}) &= \int \tilde{\pi}(\boldsymbol{\theta}|\boldsymbol{y}) d\boldsymbol{\theta}_{-j} \\ \tilde{\pi}(\boldsymbol{X}_j|\boldsymbol{y}) &= \int \tilde{\pi}(\boldsymbol{X}_j|\boldsymbol{\theta}, \boldsymbol{y}) \tilde{\pi}(\boldsymbol{\theta}|\boldsymbol{y}) d\boldsymbol{\theta},\end{aligned}$$

$\tilde{\pi}(\boldsymbol{X}_j|\boldsymbol{\theta}, \boldsymbol{y})$ depends on the approximation used, for Gaussian it is straightforward for the Laplace approximation we do another Gaussian approximation to $\tilde{\pi}(\boldsymbol{X}_{-j}|\boldsymbol{\theta}, \boldsymbol{y})$.



Modern INLA

The Gaussian approximation $\pi_G(\mathbf{X}|\boldsymbol{\theta}, \mathbf{y})$ to $\pi(\mathbf{X}|\boldsymbol{\theta}, \mathbf{y})$ is calculated from a second order expansion of the likelihood around the mode of $\pi(\mathbf{X}|\boldsymbol{\theta}, \mathbf{y})$, $\mu(\boldsymbol{\theta})$ as follows

$$\begin{aligned}\log(\pi(\mathbf{X}|\boldsymbol{\theta}, \mathbf{y})) &\propto -\frac{1}{2}\mathbf{X}^\top \mathbf{Q}(\boldsymbol{\theta})\mathbf{X} + \sum_{i=1}^n \left(b_i(\mathbf{AX})_i - \frac{1}{2}c_i(\mathbf{AX})_i^2 \right) \\ &= -\frac{1}{2}\mathbf{X}^\top (\mathbf{Q}(\boldsymbol{\theta}) + \mathbf{A}^\top \mathbf{D}\mathbf{A})\mathbf{X} - \mathbf{b}^\top \mathbf{AX}\end{aligned}$$

where \mathbf{b} is an n -dimensional vector with entries $\{b_i\}$ and \mathbf{D} is a diagonal matrix with n entries $\{c_i\}$. Note that both \mathbf{b} and \mathbf{D} depend on $\boldsymbol{\theta}$, so the Gaussian approximation is for a fixed $\boldsymbol{\theta}$.



Modern INLA

The process is iterated to find \boldsymbol{b} and \boldsymbol{D} that gives the Gaussian approximation at the mode, $\mu(\boldsymbol{\theta})$, so that

$$\boldsymbol{X}|\boldsymbol{\theta}, \boldsymbol{y} \sim N\left(\boldsymbol{\mu}(\boldsymbol{\theta}), \boldsymbol{Q}_{\boldsymbol{X}}^{-1}(\boldsymbol{\theta})\right).$$

The graph of the Gaussian approximation consists of two components,

- ① \mathcal{G}_p : the graph obtained from the prior of the latent field through $\boldsymbol{Q}(\boldsymbol{\theta})$
- ② \mathcal{G}_d : the graph obtained from the data based on the non-zero entries of $\boldsymbol{A}^\top \boldsymbol{A}$



Modern INLA

Next, the marginal conditional posteriors of the elements of \boldsymbol{X} is calculated from the joint Gaussian approximation as

$$\boldsymbol{X}_j | \boldsymbol{\theta}, \boldsymbol{y} \sim N \left((\boldsymbol{\mu}(\boldsymbol{\theta}))_j, (\boldsymbol{Q}_{\boldsymbol{X}}^{-1}(\boldsymbol{\theta}))_{jj} \right).$$

and the marginals

$$\tilde{\pi}(\boldsymbol{X}_j | \boldsymbol{y}) = \int \pi_G(\boldsymbol{X}_j | \boldsymbol{\theta}, \boldsymbol{y}) \tilde{\pi}(\boldsymbol{\theta} | \boldsymbol{y}) d\boldsymbol{\theta} \approx \sum_{k=1}^K \pi_G(\boldsymbol{X}_j | \boldsymbol{\theta}_k, \boldsymbol{y}) \tilde{\pi}(\boldsymbol{\theta}_k | \boldsymbol{y}) \delta_k.$$



Conditional posterior of η_i

In order to calculate $\tilde{\pi}(\eta_i|\mathbf{y})$, we first calculate $\tilde{\pi}(\eta_i|\boldsymbol{\theta}, \mathbf{y})$. We postulate a Gaussian density for $\eta_i|\boldsymbol{\theta}, \mathbf{y}$ such that $\tilde{\pi}(\eta_i|\boldsymbol{\theta}, \mathbf{y}) = \pi_G(\eta_i|\boldsymbol{\theta}, \mathbf{y})$, with mean

$$E(\boldsymbol{\eta}|\boldsymbol{\theta}, \mathbf{y}) = \mathbf{A}E(\mathbf{X}|\boldsymbol{\theta}, \mathbf{y}) = \mathbf{A}\boldsymbol{\mu}(\boldsymbol{\theta})$$

and covariance matrix

$$\text{Cov}(\boldsymbol{\eta}|\boldsymbol{\theta}, \mathbf{y}) = \mathbf{A}\text{Cov}(\mathbf{X}|\boldsymbol{\theta}, \mathbf{y})\mathbf{A}^\top,$$




VB corrected marginal posterior of η_i

$$\begin{aligned}\eta_j | \boldsymbol{\theta}, \mathbf{y} &\sim N(\mu_j(\boldsymbol{\theta}), \sigma_j^2(\boldsymbol{\theta})) \\ \mu_j(\boldsymbol{\theta}) &= (\mathbf{A}\boldsymbol{\mu}^*(\boldsymbol{\theta}))_j \\ \tilde{\pi}(\eta_j | \mathbf{y}) &\approx \sum_{k=1}^K \pi_G(\eta_j | \boldsymbol{\theta}_k, \mathbf{y}) \tilde{\pi}(\boldsymbol{\theta}_k | \mathbf{y}) \delta_k.\end{aligned}$$



Example (small data)

$$Y_i | \beta_0 = -1, \beta_1 = -0.5 \sim \text{Poisson}(\exp(\beta_0 + \beta_1 X_i))$$





Example (large data)

$$Y_i | \beta_0 = -1, \beta_1 = -0.5 \sim \text{Poisson}(\exp(\beta_0 + \beta_1 X_i))$$





Figure: Marginal posterior of β_0 (center) and β_1 (right) from the Laplace method (points), VBC (solid line) and INLA (broken line) approximations



Tokyo example

The Tokyo dataset in the R INLA library contains information on the number of times the daily rainfall measurements in Tokyo was more than 1mm on a specific day t for two consecutive years. In order to model the annual rainfall pattern, a stochastic spline model with fixed precision is used to smooth the data.

$$\begin{aligned}y_i | \mathcal{X} &\sim \text{Bin}\left(n_i, p_i = \frac{\exp(\alpha_i)}{1 + \exp(\alpha_i)}\right) \\(\alpha_{i+1} - 2\alpha_i + \alpha_{i-1}) | \tau &\stackrel{\text{iid}}{\sim} N(0, \tau^{-1}),\end{aligned}$$

where $i = 1, 2, \dots, 366$ on a torus, and $n_{60} = 1$ else $n_i = 2$.



Results





Figure: Posterior mean of α (left) (zoomed for the first two months (right)) from the Laplace method (points), VBC (solid line) and INLA (broken line)



Spatial survival example

Consider the R dataset Leuk that features the survival times of 1043 patients with acute myeloid leukemia (AML) in Northwest England between 1982 to 1998.





Cox spatial model

$$h(t, s) = h_0(t) \exp(\beta \mathbf{X} + \mathbf{u}(s)),$$

with

$$\eta_i(s) = \beta_0 + \beta_1 \text{Age}_i + \beta_2 \text{WBC}_i + \beta_3 \text{TPI}_i + u(s).$$

which implies a latent field of size $m = 39158$.



Results




Figure: Marginal posteriors from the Laplace method (points), VBC (solid line) and INLA (broken line)



Results




Figure: Marginal posteriors from the Laplace method (points), VBC (solid line) and INLA (broken line)



Results




Figure: Marginal posteriors from the Laplace method (points), VBC (solid line) and INLA (broken line)



Cox proportional hazards model

We simulate survival data for n patients using the following very simple Cox proportional hazards model

$$h_i(t) = h_0(t) \exp(\beta x_i) = 1.2t^{0.2} \exp(0.1x_i), \quad i = 1, 2, \dots, n,$$

where x is a scaled and centered continuous covariate, and the baseline hazard, $h_0(t)$ is estimated using a scaled random walk order one model with 50 bins. We also consider four different values of n which are $n = 10^2$, to 10^5 .



Cox proportional hazards model

n	Augmented size	classic INLA (s)	modern INLA (s)
10^2	1 327	1.6	0.1
10^3	12 657	1.3	0.4
10^4	131 807	10.2	2.3
10^5	1 302 413	113.3	22.5

Table: Results from simulation of Cox proportional hazards model



cs-fMRI model

Functional magnetic resonance imaging (fMRI) is a noninvasive neuro-imaging technique used to localize regions of specific brain activity during certain tasks.

For T timepoints and N vertices per hemisphere resulting in data $\mathbf{y}_{TN \times 1}$ with the latent Gaussian model as follows:

$$\begin{aligned}
 \mathbf{y} | \boldsymbol{\beta}, \mathbf{b}, \boldsymbol{\theta} &\sim N(\boldsymbol{\mu}_y, \mathbf{V}), \quad \boldsymbol{\mu}_y = \sum_{k=0}^K \mathbf{x}_k \boldsymbol{\beta}_k + \sum_{j=1}^J \mathbf{z}_j \mathbf{b}_j \\
 \boldsymbol{\beta}_k &= \boldsymbol{\Psi}_k \mathbf{w}_k \quad (\text{SPDE prior on } \boldsymbol{\beta}_k) \\
 \mathbf{w}_k | \boldsymbol{\theta} &\sim N(\mathbf{0}, \mathbf{Q}_{\tau_k, \kappa_k}^{-1}) \\
 \mathbf{b}_j &\sim N(\mathbf{0}, \delta \mathbf{I}) \quad (\text{Diffuse priors for } \mathbf{b}_j) \\
 \boldsymbol{\theta} &\sim \pi(\boldsymbol{\theta}),
 \end{aligned}$$

where we have K task signals and J nuisance signals.



cs-fMRI model

The data consists of a 3.5-min fMRI for each subject, consisting of 284 volumes, where each subject performs 5 different motor tasks interceded with a 3 second visual cue. Each hemisphere of the brain contained 32492 surface vertices. From these, 5000 are resampled to use for the analysis. This results in a response data vector \mathbf{y} of size **2 523 624**, with an SPDE model defined on a mesh with 8795 triangles.

The inference based on the modern formulation of INLA was computed in 148 seconds.



cs-fMRI model




Figure: Activation areas for the different tasks in the left hemisphere - visual cue, right hand motor, right foot motor, tongue motor task (from left to right)



Further details

www.r-inla.org

New default setting in INLA (VB) (previously `inla.mode = "experimental"`)

- INLA can fit many different statistical models and complex models can be built using multiple "building blocks"/random effects.
- Remove the linear predictors from the latent field → accurate posterior inference with VB correction (I - VB - LA)
- New applications that aren't feasible with INLA 1.0

-  **Gaedke-Merzhäuser, L., van Niekerk, J., Schenk, O., and Rue, H. (2023).**
Parallelized integrated nested Laplace approximations for fast Bayesian inference.
Statistics and Computing, 33(1):25.
-  **Rue, H., Martino, S., and Chopin, N. (2009).**
Approximate Bayesian inference for latent Gaussian models by using integrated nested laplace approximations.
Journal of the Royal Statistical Society: Series B (Statistical Methodology), 71(2):319–392.
-  **Van Niekerk, J., Krainski, E., Rustand, D., and Rue, H. (2023).**
A new avenue for Bayesian inference with INLA.
Computational Statistics & Data Analysis, 181:107692.
-  **van Niekerk, J. and Rue, H. (2024).**
Low-rank variational Bayes correction to the Laplace method.
Journal of Machine Learning Research, 25(62):1–25.



شكراً • Thank you



جامعة الملك عبد الله
للغعلوم والتكنولوجية
King Abdullah University of
Science and Technology