

# **An overview of the SPDE approach**

Elias Teixeira Krainski

King Abdullah University of Science and Technology (KAUST)

May 2024

# Outline




Introduction

The SPDE approach

The log Gaussian Cox process model


# **Introduction**

# Spatial statistics data




# A process over space

- ▶ (est.) Maximum temperature in US mainland, in 2022-07-20



# A process over space


- ▶ (est.) Maximum temperature in US mainland, in 2022-07-20



- ▶ The spatial domain,  $S$ , is continuous.
  - ▶ E.g.  $S \in \mathbb{R}^2$
  - ▶ E.g.  $S \in \mathbb{S}^2$  (sphere)

# A process over space


- ▶ (est.) Maximum temperature in US mainland, in 2022-07-20



- ▶ The spatial domain,  $S$ , is continuous.
  - ▶ E.g.  $S \in \mathbb{R}^2$
  - ▶ E.g.  $S \in \mathbb{S}^2$  (sphere)
- ▶  $u(\mathbf{s})$  is a stochastic process,  $\mathbf{s} \in S$

# Real world data: at a finite number of locations

At a finite set of  $n$  locations



$$u(\mathbf{s}_1), \dots, u(\mathbf{s}_n)$$



# Stochastic process $u(\mathbf{l})$

- Covariance

$$V(\mathbf{l}, \mathbf{l}') = \text{Cov}(u(\mathbf{l}), u(\mathbf{l}'))$$

- Spectral density

$$u(\mathbf{l}) = \int_{-\infty}^{\infty} e^{i\mathbf{w}\mathbf{l}} du(\mathbf{w})$$

- Kernel convolution

$$u(\mathbf{l}) = \int k(\mathbf{l} - \mathbf{u}) \mathcal{W}(\mathbf{l}) d\mathbf{u}$$

- Stochastic Partial Differential Equation - SPDE


$$\mathcal{L}_1 u(\mathbf{l}) = \mathcal{L}_2 \mathcal{W}(\mathbf{l})$$

- Conditional distributions ( DISCRETE DOMAIN !!! )

$$u_{\mathbf{l}} | u_{\text{neighbourhood}} \sim \mathbb{P}(\cdot)$$

# Whittle-Matérn covariance, Matérn (1960)


$$\Sigma_{ij} = \frac{\sigma^2 (\kappa \|\mathbf{s}_i - \mathbf{s}_j\|)^\nu K_\nu(\kappa \|\mathbf{s}_i - \mathbf{s}_j\|)}{\Gamma(\nu + d/2) (4\pi)^{d/2} \kappa^{2\nu} 2^{\nu-1}}. \text{ If } d = 2 \text{ and } \nu = 1: \text{ Whittle (1954)}$$




$$\text{practical range} = r = \sqrt{8\nu}/\kappa, \text{ corr}(r) \approx 0.13$$

# Simulations, 1D, $\sigma^2 = 1$


$v = 0.5, r = 1$




$v = 0.5, r = 4$




$v = 1.5, r = 1$




$v = 1.5, r = 4$




$v = 5, r = 1$



$v = 5, r = 4$




# Consider a set of locations



- ▶ Consider the distribution at these locations
  - ▶  $\pi(u(\mathbf{s}_1), \dots, u(\mathbf{s}_n) | \theta) = \pi(u_1, \dots, u_n | \theta) = \pi(\mathbf{u} | \theta)$

# Consider a set of locations




- ▶ Consider the distribution at these locations
  - ▶  $\pi(u(\mathbf{s}_1), \dots, u(\mathbf{s}_n) | \theta) = \pi(u_1, \dots, u_n | \theta) = \pi(\mathbf{u} | \theta)$
- ▶ If it is Gaussian with some covariance  $\Sigma(\theta)$


$$\pi(\mathbf{u} | \theta) = (2\pi)^{-n/2} |\Sigma(\theta)|^{-1/2} \exp\left(-\frac{1}{2} \mathbf{u}^T \Sigma(\theta)^{-1} \mathbf{u}\right)$$

## **The SPDE approach**

# SPDE framework



# SPDE framework



- ▶ It avoids specifying covariance!
  - ▶ D. Simpson, Lindgren, and Rue (2011)
  - ▶ D. Simpson, Lindgren, and Rue (2012)



# The Matérn's SPDE

- ▶ Whittle (1954), Whittle (1963):
  - ▶ Fields with Matérn covariance are solutions to the following Stochastic Partial Differential Equation (SPDE)

$$\tau(\kappa^2 - \Delta)^{\alpha/2} u(\mathbf{s}) = \mathcal{W}(\mathbf{s})$$

- ▶  $\kappa > 0$ : scale parameter
- ▶  $\alpha = \nu + d/2$ : smoothness
- ▶  $\Delta$  is the Laplacian

$$\Delta = \sum_{i=1}^d \frac{\partial^2}{\partial s_i^2}$$

- ▶ Discretization
  - ▶ sparse precision matrix:
  - ▶  $\mathbf{Q}_\alpha(\tau, \kappa)$ , for  $\alpha \in \{1, 2, \dots\}$ .

- ▶ Discretization
  - ▶ sparse precision matrix:
  - ▶  $\mathbf{Q}_\alpha(\tau, \kappa)$ , for  $\alpha \in \{1, 2, \dots\}$ .
- ▶  $\alpha$ 
  - ▶  $\alpha = 1$ :  $\tau \mathbf{K}_1(\kappa) = \tau(\kappa^2 \mathbf{C} + \mathbf{G})$

- ▶ Discretization

- ▶ sparse precision matrix:

- ▶  $\mathbf{Q}_\alpha(\tau, \kappa)$ , for  $\alpha \in \{1, 2, \dots\}$ .

- ▶  $\alpha$

- ▶  $\alpha = 1$ :  $\tau \mathbf{K}_1(\kappa) = \tau(\kappa^2 \mathbf{C} + \mathbf{G})$

- ▶  $\alpha = 2$ :  $\tau(\kappa^4 \mathbf{C} + 2\kappa^2 \mathbf{G} + \mathbf{G}\mathbf{C}^{-1}\mathbf{G})$

- ▶  $\alpha = 2, 3, 4, \dots$ :  $\tau \mathbf{K}_1(\kappa) \mathbf{C}^{-1} \mathbf{K}_{\alpha-2}(\kappa) \mathbf{C}^{-1} \mathbf{K}_1(\kappa)$

# Lindgren, Rue, and Lindström (2011)

- ▶ Discretization
  - ▶ sparse precision matrix:
  - ▶  $\mathbf{Q}_\alpha(\tau, \kappa)$ , for  $\alpha \in \{1, 2, \dots\}$ .
- ▶  $\alpha$ 
  - ▶  $\alpha = 1$ :  $\tau \mathbf{K}_1(\kappa) = \tau(\kappa^2 \mathbf{C} + \mathbf{G})$
  - ▶  $\alpha = 2$ :  $\tau(\kappa^4 \mathbf{C} + 2\kappa^2 \mathbf{G} + \mathbf{G} \mathbf{C}^{-1} \mathbf{G})$
  - ▶  $\alpha = 2, 3, 4, \dots$ :  $\tau \mathbf{K}_1(\kappa) \mathbf{C}^{-1} \mathbf{K}_{\alpha-2}(\kappa) \mathbf{C}^{-1} \mathbf{K}_1(\kappa)$
- ▶ Equivalent models (discretized): Whittle (1954), Besag (1974), Besag (1981), Besag and Kooperberg (1995) and Besag and Mondal (2005).

## $Q_\alpha(\tau, \kappa)$ : grid and piecewise linear basis

- ▶  $\alpha = 1$ :  $\tau \mathbf{K}_1(\kappa) = \tau(\kappa^2 \mathbf{C} + \mathbf{G})$ 
  - ▶  $d=1$ ,  $u_1, u_2, \dots, u_n$ , two neighbours

$$\tau^2 \begin{bmatrix} 1 + \kappa^2 & -1 & & & \\ -1 & 2 + \kappa^2 & -1 & & \\ & & \ddots & & \\ & & -1 & 2 + \kappa^2 & -1 \\ & & & -1 & 1 + \kappa^2 \end{bmatrix}$$


## $Q_\alpha(\tau, \kappa)$ : grid and piecewise linear basis

- ▶  $\alpha = 1$ :  $\tau \mathbf{K}_1(\kappa) = \tau(\kappa^2 \mathbf{C} + \mathbf{G})$ 
  - ▶  $d=1$ ,  $u_1, u_2, \dots, u_n$ , two neighbours

$$\tau^2 \begin{bmatrix} 1 + \kappa^2 & -1 & & & \\ -1 & 2 + \kappa^2 & -1 & & \\ & & \ddots & & \\ & & & -1 & 2 + \kappa^2 & -1 \\ & & & & -1 & 1 + \kappa^2 \end{bmatrix}$$


- ▶  $d = 2$ ,  $\mathbf{C} = \mathbf{I}$ ,  $\mathbf{G} = \text{Laplacian}$  (4 neighbours)

# Piecewise linear basis, Finite Element Method (FEM): 1d








# Piecewise linear basis, Finite Element Method (FEM): 1d





- ▶  $u(\mathbf{s}) \approx \sum_{k=1}^m \psi_k(\mathbf{s}) u_k = \mathbf{A}(\mathbf{s}, \mathbf{s}_0) u(\mathbf{s}_0),$ 
  - ▶  $\psi_k$ : basis functions evaluated at data locations  $\mathbf{s}$
  - ▶  $u_k$ : the process at the discretization points  $\mathbf{s}_0$



## Piecewise linear basis, Finite Element Method: 2d




# Piecewise linear basis, Finite Element Method: 2d




- ▶  $u(\mathbf{s}) \approx \sum_{k=1}^m \psi_k(\mathbf{s}) u_k = \mathbf{A}(\mathbf{s}, \mathbf{s}_0) u(\mathbf{s}_0),$ 
  - ▶  $\psi_k$ : basis functions evaluated at data locations  $\mathbf{s}$
  - ▶  $u_k$ : the process at the discretization points  $\mathbf{s}_0$

# Piecewise linear basis, FEM matrices


Mesh nodes



Dual mesh




Data locations




## Piecewise linear basis, FEM matrices


## Mesh nodes




## Dual mesh




## Data locations




C



**G**



A



## **The log Gaussian Cox process model**

# Point pattern and intensity

- ▶ Given a set of locations on a domain  $\mathcal{D}$
- ▶ One interest is to estimate the intensity function
  - ▶  $\lambda(\mathbf{l}), \lambda(\mathbf{l}) \geq 0, \mathbf{l} \in \mathcal{D}$ .

# Point pattern and intensity

- ▶ Given a set of locations on a domain  $\mathcal{D}$
- ▶ One interest is to estimate the intensity function
  - ▶  $\lambda(\mathbf{l}), \lambda(\mathbf{l}) \geq 0, \mathbf{l} \in \mathcal{D}$ .
  - ▶ number of events in  $\mathcal{R} \subset \mathcal{D}$ :  $y_{\mathcal{R}} \sim \text{Poisson}(n_{\mathcal{R}})$
  - ▶  $n_{\mathcal{R}} = \int_{\mathcal{R}} \lambda(\mathbf{l}) d\mathbf{l}$



# Point pattern and intensity

- ▶ Given a set of locations on a domain  $\mathcal{D}$
- ▶ One interest is to estimate the intensity function
  - ▶  $\lambda(\mathbf{I}), \lambda(\mathbf{I}) \geq 0, \mathbf{I} \in \mathcal{D}$ .
  - ▶ number of events in  $\mathcal{R} \subset \mathcal{D}$ :  $y_{\mathcal{R}} \sim \text{Poisson}(n_{\mathcal{R}})$
  - ▶  $n_{\mathcal{R}} = \int_{\mathcal{R}} \lambda(\mathbf{I}) d\mathbf{I}$
- ▶ Cox process (CP):  $\lambda(\cdot)$  is assumed to be a random function
  - ▶  $\lambda(\mathbf{I})$  is a random variable

# Point pattern and intensity

- ▶ Given a set of locations on a domain  $\mathcal{D}$
- ▶ One interest is to estimate the intensity function
  - ▶  $\lambda(\mathbf{I}), \lambda(\mathbf{I}) \geq 0, \mathbf{I} \in \mathcal{D}$ .
  - ▶ number of events in  $\mathcal{R} \subset \mathcal{D}$ :  $y_{\mathcal{R}} \sim \text{Poisson}(n_{\mathcal{R}})$
  - ▶  $n_{\mathcal{R}} = \int_{\mathcal{R}} \lambda(\mathbf{I}) d\mathbf{I}$
- ▶ Cox process (CP):  $\lambda(\cdot)$  is assumed to be a random function
  - ▶  $\lambda(\mathbf{I})$  is a random variable
- ▶ Log Gaussian Cox Process (LGCP)
  - ▶  $\log(\lambda(\cdot)) = u(\cdot)$  is a Gaussian process - GP, Møller, Syversveen, and Waagepetersen (1998) .
  - ▶  $u(\cdot|\theta), \theta$  are GP parameters

# LGCP inference

- ▶ The log-likelihood function:

$$l(\Lambda, \theta | \mathcal{Y}) = c - \int_{\mathcal{D}} \lambda(l) \partial l + \sum_{i=1}^n \log(\lambda(\mathbf{l}_i))$$

# LGCP inference

- ▶ The log-likelihood function:

$$l(\Lambda, \theta | \mathcal{Y}) = c - \int_{\mathcal{D}} \lambda(l) \partial l + \sum_{i=1}^n \log(\lambda(\mathbf{l}_i))$$

- ▶ The log-likelihood function direct approximation

$$\begin{aligned} l(\Lambda, \theta | \mathcal{Y}) &\approx c - \sum_{j=1}^m w_j \lambda(l) + \sum_{i=1}^n \log(\lambda(\mathbf{l}_i)) \\ &= c - \sum_{j=1}^m w_j \exp(\eta(l)) + \sum_{i=1}^n \eta(\mathbf{l}_i) \end{aligned}$$

approximated with  $m$  integration points.

- ▶ SPDE approach for easier computations, D. P. Simpson et al. (2016)
- ▶ more complex Point Process models using INLA in inlabru

# References

- Besag, J. 1974. "Spatial Interaction and the Statistical Analysis of Lattice Systems." *JRSS-B* 36 (2): 192–236.
- . 1981. "On a System of Two-Dimensional Recurrence Equations." *JRSS-B* 43 (3): 302–9.
- Besag, J., and C. Kooperberg. 1995. "On Conditional and Intrinsic Autoregression." *Biometrika* 82 (4): 733–46.
- Besag, J., and D. Mondal. 2005. "First-Order Intrinsic Autoregressions and the de Wijs Process." *Biometrika* 92 (4): 909–20.
- Lindgren, F., H. Rue, and J. Lindström. 2011. "An Explicit Link Between Gaussian Fields and Gaussian Markov Random Fields: The Stochastic Partial Differential Equation Approach (with Discussion)." *JRSS-B* 73 (4): 423–98.
- Matérn, B. 1960. "Spatial Variation-Stochastic Models and Their Application to Some Problems in Forest Surveys and Other Sampling Investigations. Meddelanden Fran Statens Skogsforskningsinstitut, Almaenna Foerlaget, Stockholm. (1986), 49 (5)." Springer, Berlin.
- Møller, J., A. R. Syversveen, and R. P. Waagepetersen. 1998. "Log Gaussian Cox Processes." *SJS* 25: 451–82.
- Simpson, D. P., J. B. Illian, F. Lindgren, S. H. Sørbye, and H. Rue. 2016. "Going Off Grid: Computationally Efficient Inference for Log-Gaussian Cox Processes." *Biometrika* 103 (1): 49–70.
- Simpson, D., F. Lindgren, and H. Rue. 2011. "In Order to Make Spatial Statistics Computationally Feasible, We Need to Forget about the Covariance Function." *Environmetrics*, no. 23: 65–74. <https://doi.org/https://doi.org/10.1002/env.1137>.
- . 2012. "Think Continuous: Markovian Gaussian Models in Spatial Statistics." *Spatial Statistics* 1: 16–29. <https://doi.org/http://dx.doi.org/10.1016/j.spasta.2012.02.003>.
- Whittle, P. 1954. "On Stationary Processes in the Plane." *Biometrika* 41 (3/4): 434–49.
- . 1963. "Stochastic-Processes in Several Dimensions." *Bulletin of the International Statistical Institute* 40 (2): 974–94.