

Differential Equations in Geophysical Fluid Dynamics

XVI. Wave: surface gravity wave and wave equation

Jang-Geun Choi

Center for Ocean Engineering
University of New Hampshire

Oct, 2025

This seminar is supported by mathematics community EM (maintained by Prof. Gunhee Cho)
and oceanography community COKOAA.

Recap


We discussed wind driven circulation governed by

Sverdrup (1947):

$$-f\bar{v} = -g \frac{\partial \eta}{\partial x} + \frac{\tau_x^s}{\rho_0 h} \quad (1a)$$

$$f\bar{u} = -g \frac{\partial \eta}{\partial y} + \frac{\tau_y^s}{\rho_0 h} \quad (1b)$$

$$h \left(\frac{\partial \bar{u}}{\partial x} + \frac{\partial \bar{v}}{\partial y} \right) = 0 \quad (1c)$$



Recap


We discussed wind driven circulation governed by

Stommel (1948):

$$-f\bar{v} = -g \frac{\partial \eta}{\partial x} + \frac{\tau_x^s}{\rho_0 h} - \frac{\gamma}{h} \bar{u} \quad (2a)$$

$$f\bar{u} = -g \frac{\partial \eta}{\partial y} + \frac{\tau_y^s}{\rho_0 h} - \frac{\gamma}{h} \bar{v} \quad (2b)$$

$$h \left(\frac{\partial \bar{u}}{\partial x} + \frac{\partial \bar{v}}{\partial y} \right) = 0 \quad (2c)$$



Recap


We discussed wind driven circulation governed by

Munk (1950):



$$-f\bar{v} = -g \frac{\partial \eta}{\partial x} + \frac{\tau_x^s}{\rho_0 h} \quad (3a)$$

$$f\bar{u} = -g \frac{\partial \eta}{\partial y} + \frac{\tau_y^s}{\rho_0 h} + A_h \frac{\partial^2 \bar{v}}{\partial x^2} \quad (3b)$$

$$h \left(\frac{\partial \bar{u}}{\partial x} + \frac{\partial \bar{v}}{\partial y} \right) = 0 \quad (3c)$$



A few asides



Henry Stommel, Walter Munk, and part of a letter from Stommel to Munk (left to right).

Introduction

But all assumed “steady-state ($\partial/\partial t \approx 0$)”!

The fluid surfaces found in nature keep fluctuating ($\partial/\partial t \neq 0$), resulting in what we refer to as “**waves**”.



Niagara River, New York, photo by J.-G. Choi.

Hydrostatic surface gravity wave

$$\frac{\partial \bar{u}}{\partial t} + \vec{u} \cdot \nabla \bar{u} - f \bar{v} = -g \frac{\partial \eta}{\partial x} + A_h \nabla^2 \bar{u} + \frac{\tau_x^s}{\rho_0 h} - \frac{\gamma}{h} \bar{u} \quad (4a)$$

$$\frac{\partial \bar{v}}{\partial t} + \vec{u} \cdot \nabla \bar{v} + f \bar{u} = -g \frac{\partial \eta}{\partial y} + A_h \nabla^2 \bar{v} + \frac{\tau_y^s}{\rho_0 h} - \frac{\gamma}{h} \bar{v} \quad (4b)$$

$$\frac{\partial \eta}{\partial t} + h \left(\frac{\partial \bar{u}}{\partial x} + \frac{\partial \bar{v}}{\partial y} \right) = 0 \quad (4c)$$

Substituting $\partial(4a)/\partial x$ and $\partial(4b)/\partial y$ into $\partial(4c)/\partial t$ yields

$$\frac{\partial^2 \eta}{\partial t^2} = gh \left(\frac{\partial^2 \eta}{\partial x^2} + \frac{\partial^2 \eta}{\partial y^2} \right) \quad (5)$$

that is referred to as the “**wave equation**”.


Wave equation

Let us consider one-dimensional problem ($\partial/\partial y \approx 0$ or $\bar{v} \approx 0$) and then conduct scale analysis ($t = Tt^*$, $x = Lx^*$, and $\eta = \phi\eta^*$):

$$\frac{\partial^2 \eta}{\partial t^2} = gh \frac{\partial^2 \eta}{\partial x^2}. \quad (6)$$

Unit of velocity!

$$\frac{L}{T} \sim \pm \sqrt{gh} \quad (7)$$



https://jang-geun.github.io/vis_wave1d.gif

Wave equation

Assume $\eta = \eta_0 e^{i(kx-wt)}$ and then substituting it to the governing equation yields

Wave equation

$$\frac{\partial^2 \eta}{\partial t^2} = A \frac{\partial^2 \eta}{\partial x^2}$$

Unit of velocity!

$$w = \pm \sqrt{Ak}, \quad \left(\frac{w}{k} = \pm \sqrt{A} \right)$$
$$\therefore \eta = C_1 e^{i(k(x-\sqrt{At}))} + C_2 e^{i(k(x+\sqrt{At}))}$$

Heat equation

$$\frac{\partial \eta}{\partial t} = A \frac{\partial^2 \eta}{\partial x^2}$$
$$w = -iAk^2$$
$$\therefore \eta = C_1 e^{ikx} e^{-Ak^2 t}$$

The equation showing relationship between wavenumber k and frequency w is referred to as "**dispersion relation**". This is already a solution for the differential equation.

Wave equation

Do not forget where we come from:

$$\frac{\partial \bar{u}}{\partial t} = -g \frac{\partial \eta}{\partial x} \quad (8a)$$

$$\frac{\partial \eta}{\partial t} + h \frac{\partial \bar{u}}{\partial x} = 0 \quad (8b)$$

As we assumed $\eta = \eta_0 e^{i(kx-wt)}$, we can assume $\bar{u} = \bar{u}_0 e^{i(kx-wt)}$.¹
Substituting these in to system (8) yields:

$$-iw\bar{u}_0 e^{i(kx-wt)} + gik\eta_0 e^{i(kx-wt)} = 0 \quad (9a) \qquad \underbrace{\begin{bmatrix} -w & gk \\ hk & -w \end{bmatrix}}_{\vec{M}} \begin{bmatrix} \bar{u}_0 \\ \eta_0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$-iw\eta_0 e^{i(kx-wt)} + hik\bar{u}_0 e^{i(kx-wt)} = 0 \quad (9b) \qquad (10)$$

¹We can easily see that the k and w in \bar{u} should be identical to those in η based on the equation.

Energy conservation of wave equation

The wave equation satisfies that

$$\frac{\partial}{\partial t} \left(\underbrace{\frac{1}{2} h \bar{u}^2}_{\text{Kinetic energy (KE)}} + \underbrace{\frac{1}{2} g \eta^2}_{\text{Potential energy (PE)}} \right) + gh \frac{\partial(\bar{u}\eta)}{\partial x} = 0 \quad (11)$$

Integrating over spaces with closed boundary conditions yields

$$\frac{\partial}{\partial t} \left(\underbrace{\int_0^L E dx}_{E \text{ is conserved!}} \right) = 0 \quad (12)$$

where $E = KE + PE$. Using $\bar{u} = u_0 e^{i(kx-wt)}$, $\eta = \eta_0 e^{i(kx-wt)}$, and dispersion relation $w/k = \pm\sqrt{gh}$, (11) can be written as

$$\frac{\partial E}{\partial t} \pm \underbrace{\sqrt{gh}}_{E \text{ is transported by wave!}} \frac{\partial E}{\partial x} = 0. \quad (13)$$

Conclusion

1. By substituting $\eta = \eta_0 e^{i(kx-wt)}$ into the governing equation of the wave problem, we can obtain relation between wavenumber k and frequency w , so-called dispersion relation. In case of the surface gravity wave, it is given by

$$w^2 - Ak^2 = 0 \quad (14)$$

2. $w/k (\equiv c)$ is the wave velocity that can be estimated by the dispersion relation (e.g., $c = \pm\sqrt{gh}$).
3. Energy is conserved in the wave problems and transported by the waves.

Assignment

Governing equation of Rossby wave in low frequency limit is given by

$$\frac{\partial \eta}{\partial t} - g \frac{gh}{f^2} \frac{\partial \eta}{\partial x} = 0 \quad (15)$$

Using wave-like solution $\eta = \eta_0 e^{i(kx-wt)}$, find (1) dispersion relation and (2) wave velocity.

References I

-  Munk, Walter H (1950). "On the wind-driven ocean circulation". In: *Journal of Atmospheric Sciences* 7.2, pp. 80–93.
-  Stommel, Henry (1948). "The westward intensification of wind-driven ocean currents". In: *Eos, Transactions American Geophysical Union* 29.2, pp. 202–206.
-  Sverdrup, Harald Ulrich (1947). "Wind-driven currents in a baroclinic ocean; with application to the equatorial currents of the eastern Pacific". In: *Proceedings of the National Academy of Sciences* 33.11, pp. 318–326.