Differential Equations in Geophysical Fluid Dynamics

XXII. Wave of quasi-geostrophic equation

Jang-Geun Choi

Center for Ocean Engineering
University of New Hampshire

Feb, 2026

This seminar is supported by mathematics community EM (maintained by Prof. Gunhee Cho) and oceanography commuinity COKOAA.

Recap

Govering equations of the quasi-geostrophic model are given by

$O(1)$ balances (predominant geostrophic balance):

$$\bar{v}_0 = \frac{g}{f_0}\frac{\partial \eta}{\partial x} \quad \text{(1a)} \qquad \bar{u}_0 = -\frac{g}{f_0}\frac{\partial \eta}{\partial y} \quad \text{(1b)} \qquad \frac{\partial \bar{u}_0}{\partial x} + \frac{\partial \bar{v}_0}{\partial y} = 0 \quad \text{(1c)}$$

$O(Ro)$ balances (small perturbations):

$$\bar{v}_1 = \frac{1}{f_0}\left(\frac{\partial \bar{u}_0}{\partial t} + \bar{u}_0\frac{\partial \bar{u}_0}{\partial x} + \bar{v}_0\frac{\partial \bar{u}_0}{\partial y} - \beta y \bar{v}_0\right) \qquad \text{(2a)}$$

$$\bar{u}_1 = -\frac{1}{f_0}\left(\frac{\partial \bar{v}_0}{\partial t} + \bar{u}_0\frac{\partial \bar{v}_0}{\partial x} + \bar{v}_0\frac{\partial \bar{v}_0}{\partial y} + \beta y \bar{u}_0\right) \qquad \text{(2b)}$$

$$\frac{\partial \eta}{\partial t} + \frac{\partial (h\bar{u}_1)}{\partial x} + \frac{\partial (h\bar{v}_1)}{\partial y} + \frac{\partial (\eta\bar{u}_0)}{\partial x} + \frac{\partial (\eta\bar{v}_0)}{\partial y} = 0 \qquad \text{(2c)}$$

Recap

Those can be rewritten as the vorticity conservation equation given by

$$\frac{\partial q_0}{\partial t} + \bar{u}_0 \frac{\partial q_0}{\partial x} + \bar{v}_0 \frac{\partial q_0}{\partial y} = 0 \tag{3a}$$

$$q_0 = \xi_0 + f - \frac{f_0}{h}\eta \tag{3b}$$

Relative vorticity of geostrophic current

$\nabla^2 \eta$

$$\xi_0 = \frac{\partial \bar{v}_0}{\partial x} - \frac{\partial \bar{u}_0}{\partial x} = \frac{g}{f_0}\left(\frac{\partial^2 \eta}{\partial x^2} + \frac{\partial^2 \eta}{\partial y^2}\right) \tag{3c}$$

Using stream function, $\bar{u}_0 = -\partial\psi/\partial y$ and $\bar{v}_0 = \partial\psi/\partial x$, it can be written as

$$\frac{\partial q_0}{\partial t} + J(\psi, q_0) = 0 \tag{4a}$$

Jacobian operator

$$J(\psi, q_0) = \frac{\partial \psi}{\partial x}\frac{\partial q_0}{\partial y} - \frac{\partial \psi}{\partial y}\frac{\partial q_0}{\partial x} \tag{4b}$$

$$q_0 = \nabla^2 \psi + f - \frac{1}{Rd^2}\psi \tag{4c}$$

Consideration of varying bathymetry

$$\frac{\partial \eta}{\partial t} + \frac{\partial((h+\eta)\bar{u})}{\partial x} + \frac{\partial((h+\eta)\bar{v})}{\partial y} = 0 \tag{5}$$

Let us decompose not only velocity component ($\bar{u} = \bar{u}_0 + \bar{u}_1$ and $\bar{v} = \bar{v}_0 + \bar{v}_1$) but also water depth $h = h_0 + h'(x, y)$ where h_0 and h' indicate constant and variation components of h.

$O(Ro)$

$$\begin{aligned}
&\frac{\partial \eta}{\partial t} + h_0\left(\frac{\partial \bar{u}_0}{\partial x} + \frac{\partial \bar{v}_0}{\partial y}\right) + h_0\left(\frac{\partial \bar{u}_1}{\partial x} + \frac{\partial \bar{v}_1}{\partial y}\right) \qquad O\left(Ro\frac{H'}{h_0}\right) \\
&+ \left(\frac{\partial (h'\bar{u}_0)}{\partial x} + \frac{\partial (h'\bar{v}_0)}{\partial y}\right) + \left(\frac{\partial (h'\bar{u}_1)}{\partial x} + \frac{\partial (h'\bar{v}_1)}{\partial y}\right) \\
&O\left(\frac{H'}{h_0}\right) + \left(\frac{\partial (\eta\bar{u}_0)}{\partial x} + \frac{\partial (\eta\bar{v}_0)}{\partial y}\right) + \left(\frac{\partial (\eta\bar{u}_1)}{\partial x} + \frac{\partial (\eta\bar{v}_1)}{\partial y}\right) = 0
\end{aligned} \tag{6}$$

Consideration of varying bathymetry

What if $Ro \ll H'/h_0$? Note that we already assumed $Ro \ll 1$. So, if $H'/h_0 \approx 1$, zeroth order balance becomes

$$\frac{\partial(h\bar{u}_0)}{\partial x} + \frac{\partial(h\bar{v}_0)}{\partial y} = 0 \tag{7}$$

that can be written as

$$\frac{\partial h}{\partial t} + \bar{u}_0 \frac{\partial h}{\partial x} + \bar{v}_0 \frac{\partial h}{\partial y} = 0 \tag{8}$$

$$\left(\because \frac{\partial h}{\partial t} = 0, \quad \frac{\partial \bar{u}_0}{\partial x} + \frac{\partial \bar{v}_0}{\partial y} = 0 \right)$$

that is equivalent to the **Taylor–Proudman theorem**!

Consideration of varying bathymetry

Quasi-geostrophic model chooses $H'/h_0 \approx Ro$. In this case, $O(Ro)$ balance of the continuity equation is given by

$$\frac{\partial \eta}{\partial t} + \frac{\partial((h' + \eta)\bar{u}_0)}{\partial x} + \frac{\partial((h' + \eta)\bar{v}_0)}{\partial y} + h_0 \left(\frac{\partial \bar{u}_1}{\partial x} + \frac{\partial \bar{v}_1}{\partial y} \right) = 0 \quad (9)$$

Note that there is no difference in momentum equations. Vorticity equation using (1), (2), and (9) is given by

$$\frac{\partial q_0}{\partial t} + \bar{u}_0 \frac{\partial q_0}{\partial x} + \bar{v}_0 \frac{\partial q_0}{\partial y} = 0 \quad (10a)$$

$$q_0 = \xi_0 + f - \frac{f_0}{h_0} (h + \eta) \quad (10b)$$

Planetary and topographic β-effect

$$\frac{\partial q_0}{\partial t} + \bar{u}_0 \frac{\partial q_0}{\partial x} + \bar{v}_0 \frac{\partial q_0}{\partial y} = 0 \tag{11a}$$

$$q_0 = \xi_0 + f - \frac{f_0}{h_0}(h + \eta) \tag{11b}$$

In case of constant depth with β-plane:

$$q_0 = \xi + \beta y - \frac{f_0}{h_0}\eta \tag{12}$$

In case of f-plane with varying topography $h = \alpha y + h_0$:

$$q_0 = \xi + \frac{\alpha f_0}{h_0} y - \frac{f_0}{h_0}\eta \tag{13}$$

So, they can do same role!

Wave-like solution to quasi-geostrophic model

Let us consider quasi-geostrophic equations written by the stream function:

$$\frac{\partial q_0}{\partial t} + \underbrace{\frac{\partial \psi}{\partial x}\frac{\partial q_0}{\partial y} - \frac{\partial \psi}{\partial y}\frac{\partial q_0}{\partial x}}_{J(\psi,\, q_0)} = 0 \tag{14a}$$

$$q_0 = \nabla^2 \psi + f - \frac{1}{Rd^2}\psi \tag{14b}$$

that can be rewritten by

$$\frac{\partial}{\partial t}\left(\nabla^2\psi - \frac{1}{Rd^2}\psi\right) + J\left(\psi, \nabla^2\psi - \frac{1}{Rd^2}\psi\right) + \beta\frac{\partial \psi}{\partial x} = 0 \tag{15}$$

Once "**monochromatic**" wave-like solution ($\psi = \psi_0 e^{i(kx+ly-wt)}$) is assumed, we can see

$$J\left(\psi, \nabla^2\psi - \frac{1}{Rd^2}\psi\right) = 0 \tag{16}$$

so it becomes equation of the linear Rossby wave perturbed by inertia!

Role of nonlinearity

Let us consider wave-like solution with two differen modes:

$$\psi = A_1 e^{\underbrace{i(k_1 x + l_1 y - w_1 t)}_{\equiv \theta_1}} + A_2 e^{\underbrace{i(k_2 x + l_2 y - w_2 t)}_{\equiv \theta_2}} \tag{17}$$

In this case, the Jacobian operator is non-zero:

$$J\left(\psi, \nabla^2\psi - \frac{1}{Rd^2}\psi\right) = A_1 A_2 (k_1 l_2 - k_2 l_1)({K_2}^2 - {K_1}^2) e^{\theta_1 + \theta_2} \tag{18}$$

where $K_n = \sqrt{{k_n}^2 + {l_n}^2}$.

Because of its nonlinearity, principle of superposition does not work!

Role of nonlinearity

Predominant (Big)!

$$\underbrace{\frac{\partial}{\partial t}\left(\nabla^2\psi - \frac{1}{Rd^2}\psi\right)}_{1/(\beta LT)} + \underbrace{J\left(\psi, \nabla^2\psi - \frac{1}{Rd^2}\psi\right)}_{\Psi/(\beta L^3)\text{: small}} + \underbrace{\beta\frac{\partial\psi}{\partial x}}_{1} = 0 \quad (19)$$

Choosing $1/(\beta LT) = 1$ and $\Psi/(\beta L^3) \ll 1$ and applying perturbation method yields below.

$O(1)$ balance:

$$\frac{\partial}{\partial t}\left(\nabla^2\psi_0 - \frac{1}{Rd^2}\psi_0\right) + \beta\frac{\partial\psi_0}{\partial x} = 0 \quad (20)$$

$O(\Psi/(\beta L^3))$ balance:

$$\frac{\partial}{\partial t}\left(\nabla^2\psi_1 - \frac{1}{Rd^2}\psi_1\right) + \beta\frac{\partial\psi_1}{\partial x} = -J\left(\psi_0, \nabla^2\psi_0 - \frac{1}{Rd^2}\psi_0\right) \quad (21)$$

Role of nonlinearity

Solution to (21) is given by

$$\psi_1 = \frac{A_1 A_2 (k_1 l_2 - k_2 l_1)({K_2}^2 - {K_1}^2)}{i(w_3({k_3}^2 + {l_3}^2 + Rd^{-2}) + \beta k_3)} e^{\theta_3} \tag{22}$$

where $\theta_3 = i(k_3 x + l_3 y - w_3 t)$, $k_3 = k_1 + k_2$, $l_3 = l_1 + l_2$, and $w_3 = w_1 + w_2$, so it is new third mode depending on the original two modes.

If $w_3({k_3}^2 + {l_3}^2 + Rd^{-2}) + \beta k_3) = 0$,[1] the solution becomes

$$\psi_1 = -\frac{A_1 A_2 (k_1 l_2 - k_2 l_1)({K_2}^2 - {K_1}^2)}{{k_3}^2 + {l_3}^2 + Rd^{-2}} t e^{\theta_3}. \tag{23}$$

It grows as time!

that was discussed by (Longuet-Higgins and Gill, 1967).

[1] Note that this is dispersion relation of the Rossby wave perturbed by inertia

References I

Longuet-Higgins, Michael Selwyn and Adrian Edmund Gill (1967). "Resonant interactions between planetary waves". In: *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences* 299.1456, pp. 120–144.