
Josh Andrews - ECE 498 - HW #7

Table of Contents

Question 1	1
Question 2	1
Question 3	1

Question 1

Finding roots in interval

```
clear all; clc;

f = inline('0.05*x - sin(x)');
x0 = [-5 -1];
x(1) = fzero(f, x0);
x0 = [-1 1];
x(2) = fzero(f, x0);
x0 = [1 5];
x(3) = fzero(f, x0);
disp('The roots with x [-5 5] are:')
disp(x)
```

```
The roots with x [-5 5] are:
-2.9915      0      2.9915
```

Question 2

Finding roots of polynomial

```
p = [1, 2, -7, -8, 10, 0, 8, 0];
r = roots(p);
disp('The roots of polynomial p are:')
disp(r)
```

```
The roots of polynomial p are:
0.0000 + 0.0000i
-3.0529 + 0.0000i
-2.0000 + 0.0000i
2.0000 + 0.0000i
1.2835 + 0.0000i
-0.1153 + 0.7051i
-0.1153 - 0.7051i
```

Question 3

solve set of equations

```
x0 = [1,1,1];
%nle(x0);
x = fsolve('nle', x0);
disp('X = ')
disp(x(1))
disp('Y = ')
disp(x(2))
disp('Z = ')
disp(x(3))
```

Equation solved.

fsolve completed because the vector of function values is near zero as measured by the default value of the function tolerance, and the problem appears regular as measured by the gradient.

```
X =
    2.0000
```

```
Y =
    2.0000
```

```
Z =
    2.0000
```

Published with MATLAB® R2017b